



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

BUHR B



a39015 00008545 9b

PROPERTY OF
*University of
Michigan
Library*

1817

ARTES SCIENTIA VERITAS

BULLETIN No. 3.

1902.

MYCOLOGICAL SERIES, No. 1

BULLETIN
of the
LLOYD LIBRARY
of
BOTANY, PHARMACY AND
MATERIA MEDICA

J. U. & C. G. LLOYD
CINCINNATI, OHIO

MYCOLOGICAL SERIES, No. 1

THE GENERA
OF
GASTROMYCETES

Illustrated with 49 Figures

By C. G. LLOYD

QK
600
.L79
no. 1-6

INTRODUCTION.



Fig. 1.

a—A basidium, bearing spores.
b—An ascus, containing spores.

Fungi, the larger fungi, are divided into two classes, 1st, the Basidiomycetes, which have the spores borne free on a basidia; 2nd, the Ascomycetes, which have the spores borne in a sack called an ascus. In this pamphlet we have to deal only with part of the first class.

The Basidiomycetes can in turn be divided into two very natural classes, 1st, the Hymenomycetes, those that have the spores exposed and free from the beginning, or at least from a very early state; 2nd, the Gastromycetes, those that develop the spores in cavities or chambers *within the tissue of the plant*. We are aware that these divisions are not in keeping with the very latest authorities which primarily divide the Basidiomycetes into sections based respectively on septate or nonseptate basidia, but we believe that these latter divisions while possibly theoretically correct, tend only to confuse matters excepting to the advanced and expert student.

It should not be inferred from the above that in order to recognize the Gastromycetes it is necessary to study the nature of the basidia, or make other minute anatomical examination. As a matter of fact, the merest tyro soon learns to recognize on sight the various phalloids, bird-nest fungi, and various kinds of "puff-balls" constituting the Gastromycetes and they were well classified before their anatomical structures were known.

Terms used in the description of the Gastromycetes.

PERIDIUM.

The shell or hull, enclosing the spore mass of a gastromycete is called the peridium. In many of the different genera, the simplest type is a simple, uniform layer such as surrounds the spore mass in the accompanying cut of Scleroderma. (Fig. 2.) Usually however, the peridium consists of two distinct layers, called the outer peridium or exoperidium and the inner peridium or endoperidium. In Geaster, the outer peridium is thick and when the plant (*) ripens it splits in a stellate manner separating from the inner peridium and becoming more or less reflexed.



Fig. 2
Section of a Scleroderma.

(*) In speaking of the "plants" it will be observed that we do not use precise language for what we call the "plants" are really the fruit bodies, compound sporophores, of the fungi corresponding to the fruit of flowering plants, but it seems more natural in a work intended largely for general distribution to call a "puff-ball" or a "toad-stool" a plant than a fruit body. The vegetative portions of fungi, corresponding to the stem of flowering "plants" are thread-like growths called the mycelium, that permeate the soil or rotten wood, and which in reality bear the fruit bodies, or sporophores, that we have chosen here to call plants.

(See Plate 8, figures.) In *Lycoperdon*, *Bovista*, and many genera, the outer peridium is a thin, friable coat, often bearing spines or warts.




Fig. 3.

A *Lycoperdon* with the cortex peeling off. As the plant matures this membrane usually peels off and disappears. (See Fig. 3.) When the outer peridium is of this nature it is called the cortex. In the genus *Mitremyces* the outer peridium is a thin membrane covered with a thick gelatinous mass. (See Plate 5, Fig. 29.) As the plant ripens this thin membrane breaks into little pieces which curl up and fall off carrying the thick gelatinous coat with them. All phalloids are in the young state enclosed in a thick, gelatinous membrane corresponding to a peridium, and called the volva. The outer peridium of *Mitremyces* is also usually called the volva. The phalloid is only enclosed in its volva during its young or "egg" state. (See Plate 1, Fig. 16.) When the plant grows the volva is ruptured at the apex, and remains as a cup at the base of the plant. (See Plate 1, Fig. 15.)

THE STEM OR STALK.

Many genera of *Gastromycetes*, (*Lycoperdon*, *Bovista*, etc.) are entirely destitute of any stalk or stem, but other genera (*Tylostoma*, *Queletia*, etc.) are characterized by having the peridium borne on a distinct stalk. The base of the peridium of *Lycoperdon*, (see Plate 10, Fig. 45,) or *Calvatia*, is often contracted into stalk-like appearance, but must not be confused with the true stem of such genera as *Tylostoma*. Stalked *gastromycetes* are readily divided into two tribes: *Tylostomeae* in which the stalk is entirely distinct from the peridium and *Podaxineae* in which the stalk is continuous, forming an axis reaching the apex of the peridium.

THE GLEBA.

The inside of an immature puff-ball is filled with a white fleshy mass of soft cellular matter called the gleba.

Our study of the *Gastromycetes* has been confined to the mature specimens in our collection, but it will not be amiss to give the developments as recorded by De Bary, (whom we have for the most part copied in some sentences literally) Tulasne, Corda, Berkeley and others to whom we are indebted for our knowledge of the minute structure of the gleba. At first it is simply a cellular mass, but as the plant grows it gradually assumes the form of a tissue of minute chambers. The chambers of the gleba are in countless numbers, usually too small to be seen by the naked eye, and are narrow, irregularly curved, branched cavities, separated from one another by their curved plates

of tissue which anastomose with one another in every direction. In the accompanying cut (Fig. 4) the chambers of the gleba can be seen with the eye (if the printer does justice to the cut); the chambers of the sterile base are very large and evident. The walls of the



Fig. 4.
Section of a young Lycoperdon.



Fig. 5.
An ideal enlarged chamber of the gleba of Geaster.

chambers consist of layers of branched hyphæ bearing a hymenial layer on both surfaces which line the interior walls of the cavities. The hymenial hyphæ terminate in basidia bearing usually four spores. The figure which we give herewith (Fig. 5) taken from Engler & Prantl, (originally from Tulasne) of an enlarged gleba chamber illustrates this structure. In this figure the hyphæ constitute the thread-like tissue forming the walls of the chamber, the basidia are seen to bear four sessile spores.

In *Scleroderma*, *Geaster hygrometricus*, *Polysaccum*, and in certain other genera, all the hyphæ which enter a chamber are elongating, copiously branched, and woven together into a loose mass filling the chamber. (Fig. 6.) Plants possessed of this structure form




Fig. 6.
Basidia (enlarged) of *Scleroderma*.

the order Plectobasidiaceae of Fischer, but we think even if theoretically correct, it is not a matter of policy to classify plants by minute

anatomical differences which only an expert microscopist can trace and concerning which the ordinary student knows nothing excepting that which he reads.

BASIDIA.




Fig. 7. Basidia.
a—Tylostoma. c—Bovista.
b—Geaster. d—Mitromyces.

If we believed in the German scheme of classification we would consider the basidia the most important part about a *Gastromyces* and all other characters subservient to them. We are told that the basidia are of various shapes and that in some plants they form a lining to the gleba chambers. The way the spores are borne on the basidia is also characteristic; in *Geaster* they are almost sessile; in *Bovista* the spores are borne on long stalks called sterigmata; in *Tylostoma* and *Mitromyces* they are almost sessile and lateral. The number of spores also vary from four in *Lycoperdon* to a dozen or more in *Mitromyces* as shown in our cuts. These cuts copied from Engler & Prantl were originally from several authors. (*)

THE RIPENING OF PUFF-BALLS.

One of the most curious phenomena in connection with these plants is the change that takes place when the spores ripen. As the young plant grows the interior is a solid, white, firm, fleshy mass. When it reaches full size and ripens the tissues deliquesce, become moist, discolored, the tissues of the tramal chambers are absorbed and disappear, and finally the water dries away, leaving the peridium filled with a dry, dusty mass, usually consisting of slender threads and countless multitudes of ripe spores. This is now called the spore mass and the threads capillitium. The phenomenon of ripening in all *Gastromycetes* I believe is attended with deliquescence and absorption of more or less of the hyphal elements of the gleba, but the walls of the chambers do not in all genera disappear.

(*) We have given thus the detail of the minute structure of *Gastromycetes* as it is the basis of modern classification. Personally we do not approve of it. Assuming that it is the correct theory the time is not ripe for it. The basidial structure of comparatively few species is known. With by far the greater part of them and many genera the basidial structure is only conjectural. It seems to be the tendency of some writers to select the most obscure and difficult points on which to base classification. This has one advantage, it gives an air of greater learning. For our part we feel that a system based on points of difference of the mature plant obvious to the student, is more satisfactory and rational. To our mind there is no room in any *Natural* system of classification for the *Nidulariaceae* between *Astræus* (admitting the genus for argument) and *Geaster*, no matter what their basidial structure may be.

PERIDIOLES.

In the Nidulariaceæ or "bird-nest fungi" the walls thicken and each chamber remains as a separate, little seed-like body enclosing the spores. This is called a peridiole. (See Fig. 8.) In *Arachnion* the




Fig. 8.
Section (enlarged) of a *Cyathus*,
showing peridioles.



Fig. 9.
Polysaccum with upper portion broken off,
showing peridioles.

"puff-ball" is filled with sack-like peridioles appearing to the eye as grains of sand. In *Polysaccum* (see Fig. 9) the peridioles are large and only partially separated from each other, the interior of a broken plant having the appearance of being honeycombed. In *Scleroderma* the walls of the gleba chambers are more or less permanent in the different species. In some specimens of *S. bovista* they remain almost perfect and approximate *Polysaccum*. In most species of *Scleroderma* however, only fragments of the walls are mixed with the spores.

CAPILLITIUM.

The threads that are contained in the spore mass of various *Gastromycetes*, though absent in many genera, are characteristic in each genus that has them, and are important factors in classification. How much longer I do not know, but certainly as far back as 1876, the peculiarities of the capillitium of the different genera were described and illustrated by Hesse.

There are two distinct types of capillitium threads. 1st, the threads are long hair-like strands, simple or more or less branched and interwoven, (see Fig. 11) proceeding from the inner walls of the peridium or the columella of the plant. 2nd, the threads are relatively short and branched, each entirely separate and distinct from the other (see Fig. 10), though the branches are usually interwoven, and have no connection with the peridium or columella. The latter type is characteristic of *Bovista*, *Bovistella* and *Mycenastrum*. Threads of the first type are usually broken into short fragments in the ripe spore mass, but are readily distinguished from those of the second type by the blunt ends of the fracture. Threads of the 2nd type when perfect run out in all directions into sharp pointed branches. (*)



Fig. 10.

A Capillitium thread (magnified) of *Bovista*.



Fig 11.

Capillitium of *Tylostoma*. (magnified.)

Capillitium threads have varying character in different genera. In *Calvatia* they are long, branched, and interwoven. In *Catostoma*, mostly broken in short fragments. In *Tylostoma* often septate. In *Mycenastrum* they bear little spiny processes. Usually they are colored, sometimes hyaline.

The hyphal strands that persist as capillitium are shown by *Tulasia* as penetrating and passing through the walls and chambers of the gleba. We can readily understand this structure in such genera as *Lycoperdon* where they are attached to the peridium or columella, but the exact attachment is obscure to our mind where they are "separate threads."

(*) Capillitium threads are relatively large microscopic objects, often visible to the naked eye and readily examined under low (1 inch lens) magnifying power. By simply pressing a fragment of spore mass on a slide the nature of the threads can usually be readily made out. To secure separate threads of *Bovista* type put a little spore mass in a small vial half filled with alcohol. Agitate violently and pour the alcohol over a clean slide, and separate threads can be readily floated out.

SPORES.

The ripe spores of *Gastromycetes* are readily examined under a microscope ($\frac{1}{4}$ inch power is the best) and afford characters useful in distinguishing species. Some spores are smooth, some spinulose; more are globose but some are oblong or oval. They vary also in size and in color. Some spores are borne on the basidia on long sterigmata (see fig. 7c) which as the spores ripen persist attached to the spore and are known then as pedicels. (See Fig. 14.) The value of pedicellate



Fig. 12.
Globose spores of *Mitromyces*
lutescens.



Fig. 13.
Oblong spores of *Mitromyces*
cinnabarinus.



Fig. 14.
Pedicellate spores of *Bovista*

spores as a specific character is a disputed question. I am convinced that in some species the persistence of the pedicel depends on the stage of development when the plant is collected as I have found pedicellate and non-pedicellate spores in different plants of same species and of same collection. Masee says the pedicels of spores of old herbarium specimens are always broken off, hence the character of no value. We know however, that there are certain species such as *Bovistella* *Ohiense*, *Bovista plumbea*, in which the pedicellate spores are constant and persist for years, and we feel that in such cases the character is of value, even if it does disappear with age.

STERILE BASE.

In *Lycoperdon*, *Calvatia* and other genera, all portions of the gleba are not fertile and spore bearing. The lower portion called the "sterile base" consists simply of sterile cells or threads. The "sterile base" has been made a character to distinguish genera, *Globaria* of recent writers being simply *Lycoperdon* devoid of a sterile base. While it is a good primary character to divide the genus *Lycoperdon*, there are all shades of development from species with none at all, through species with very little, to species with it strongly developed, and I feel that it alone should not be held of generic importance. Several species shed their spores but the sterile bases persist through the winter and are often picked up for perfect plants. Bosc described and illustrated *Calvatia cyathiformis* on such a remnant, thinking it was a perfect plant.

HISTORICAL.

As previously stated, we do not believe in the recent German classification of *Gastromycetes* based on minute anatomical differences such as basidia. A natural system drawn from characters found in the mature plants has been evolved gradually, can be readily understood, and plants can be identified by anyone with little trouble. There is no department of mycology where there is so much confusion as in the *Gastromycetes*, and it is a most puzzling task to try and trace the species through the writings of the various authors. While it is necessary for us to study the history of the plants, we do not attach the importance to solving these old time puzzles that we do to the study of the plants themselves. Saccardo's *Gastromycetes* is probably the poorest compilation of all his volumes. *Sclerodermas*, *Mycenastrums* and *Bovistellas*; *Bovistas*, *Catastomas* and *Globalarias*,* are all jumbled indiscriminately together and often the same species appears under two or three different names. Absence of illustrations, or crude attempts at it on the part of authors, are responsible for much of this trouble. Endeavors on the part of authors, such as Fries and Persoon, to classify species that they know nothing of, on these crude illustrations further contributed to the confusion. In our pamphlet we shall make no attempt to compile genera or species that we have not in our collection or have not seen and studied. Most of our specimens have been submitted to Bresadola and Patouillard, in our opinion the best authorities in the world.

In our country there have been three important workers with the *Gastromycetes*, Peck, Trelease and Morgan. Prof. Peck wrote an account of the New York species of *Lycoperdon* which appeared in the 32nd Report (1879). This is a very plain description of the species that he had seen and studied as they grew, and is one of the best accounts that has appeared. Those who live in the Eastern section of our country, can take this old monograph and make out most all the *Lycoperdons* that they find. Practically the same paper, to which was added a compilation of species described which he had not met, was published in the Transactions of the Albany Institute under the title of "United States Species of *Lycoperdon*." A paper on "The morels and puff-balls of Madison (Wisc.)," by Prof. Trelease, appeared (1889) in Transactions of Wisconsin Academy of Sciences. This article gives evidence of great study and research, and the conclusions Prof. Trelease reached are mostly maintained at the present day. Unfortunately, the paper is illustrated by most miserable figures.

Prof. Morgan has probably done more work on the *Gastromycetes* in this country than any other man, wrote four papers on the subject that were published in the Journal of the Cincinnati Society of Natural History (1889 to 1892). The field was not completely covered as the work was not completed. Morgan made a critical study of the internal structure, especially the capillitium of "puff-balls," and established several new genera that are universally recognized.

(*) Using the term for convenience for *Lycoperdon* without sterile base.

CLASSIFICATION.

Gastromycetes can be readily divided into four families widely differing from each other as to the nature of the mature plants.

FAM. 1.—PHALLOIDAE.—PHALLOIDS. Plant fleshy, enclosed in a gelatinous volva when young. The gleba deliquescent and becoming a mucilaginous (generally foetid) mass.

Phalloids are noted for the foetid odor that they have and for their bizarre shapes. They force themselves to the attention of the most unobserving and are often called such appropriate names as Stink Fungus, Stink Horns, Dead Mens' Fingers. Our most common species are *Phallus duplicatus*, and *Phallus Ravenelii*. In the South, *Clathrus columnatus*.

FAM. 2.—NIDULARIACEAE.—NEST FUNGI. Plants shaped like little cups, opening at the top, and containing a number of little seed-like bodies (peridioles). They look something like little birds' nests and are often called "Birds-nest fungi." *Crucibulum vulgare* and *Nidularia striatus* are our most common species.

FAM. 3.—HYMENOGASTRACEAE.—HYPOGEAL FUNGI. Peridium indehiscent; gleba cavities permanent, not resolved into a mass of spores; capillitium absent.

This family is mostly subterranean like the true tubers or truffles. Harkness has recorded many species from the Pacific Coast, and we have seen three from the section east of the Mississippi. It is probably that many occur but have been overlooked on account of their subterranean habits.

FAM. 4.—LYCOPERDACEAE.—PUFF BALLS.—Ripe peridium enclosing a mass of dry spores, often mixed with capillitium. Sometimes the gleba walls persist forming peridioles, but in those cases the peridioles are filled with a mass of dry powdery spores.

The largest and most frequent tribe of Gastromycetes and embracing all the families known as "Puff-balls."

GENERA OF LYCOPERDACEAE.

For the time being we will pass over the genera embraced in the first three families and enumerate the genera with which we are familiar, of the "puff-ball" family. We would divide the plants into four tribes.

TRIBE 1.—TYLOSTOMEAE.—Plant stalked. Stalk distinct from the peridium. Capillitium present.

Peridium opening by an apical mouth,

Volva indistinct, adherent, TYLOSTOMA.

Volva cup-like, CHLAMYDOPUS.

Peridium circumscissal, BATTARREA.

Peridium opening irregularly,

Volva none, QUELETIA.

Volva thick, permanent, DICTYOCEPHALOS.

TRIBE 2.—PODAXINEAE.—Plant stalked. Stalk continuous to the apex of the peridium forming an axis.

- Gleba with irregular persistent chambers,
 - Peridium, club-shaped, CAULOGLOSSUM.
 - Peridium, round or conical, SECOTIUM.
- Gleba with sinuate, lamellate plates, GYROPHRAGMIUM.
- Walls of the gleba chambers not persistent, PODAXON.

TRIBE 3.—SCLERODERMEAE.—Plant not stalked, or stalk short, confluent with the peridium. Capillitium none.*

- Peridium of a single layer,
 - Walls of the gleba chambers persistent forming peridioles, POLYSACCUM.
 - Walls of the gleba chambers most disappearing or only partially persistent, SCLERODERMA.
- Peridium, double,
 - Outer peridium, thin (a cortex) ARACHNION.
 - Outer peridium, thick, gelatinous, MITREMYCES.

TRIBE 4.—LYCOPERDEAE.—Plant not stalked. Spore mass, dry spores mixed with capillitium.

TRIBAL ALLIANCE 2.—GEASTRAE,—EARTH STARS.—Peridium double, outer peridium thick, persistent, splitting into segments and recurving.

- Mouth, one, GEASTER.
- Mouths, several, MYRIOSTOMA.

TRIBAL ALLIANCE 2.—BOVISTAE,—TUMBLERS.—Outer peridium thin (cortex mostly peeling off). Inner peridium firm or papery. Mature plant loosened from place of growth.

- Capillitium of separate threads, with slender pointed branches, BOVISTA.
- Capillitium of separate threads bearing spiny points, MYCENASTRUM.
- Capillitium threads broken into short fragments with blunt ends, CATASTOMA.

TRIBAL ALLIANCE 3.—LYCOPERDAE.—TRUE PUFF BALLS.—Outer peridium thin (cortex, mostly disappearing.) Inner peridium usually flaccid. Plants normally remaining attached to place of growth.

- Capillitium of separate threads with slender pointed branches, BOVISTELLA.
- Capillitium long threads more or less broken in fragments,
 - Peridium, opening by definite mouth, LYCOPERDON.
 - Peridium, irregularly ruptured, no lining membrane CALVATIA.
 - Peridium, irregularly ruptured, furnished with a lining membrane, HYPOBLEMA.

*Basing the Tribe thus for convenience on the absence of capillitium, it embraces widely diverging genera, but we prefer to do this at least for the present rather than to multiply the tribes. Mitremyces is the type of a good tribe, Arachnion perhaps of another.

ILLUSTRATIONS.

We present herewith eleven plates illustrating the various genera. We expect to publish from time to time, pamphlets describing and illustrating the species of each genera.

Acknowledgement of Sources of Illustrations.

- Fig. 1, 5, 6, 7, 8, copied from Engler & Prantl, (originally from various sources.)
- Fig. 20, copied from drawing by V. S. White.
- Fig. 10, 11, 12, 13, 14, 42, Microphotographs by Dr. Edward Thompson.
- Fig. 9, 36, 37, Specimens from Mrs. Delia Sams, New Smyrna, Florida.
- Fig. 17, 18, 35, Specimens from Simon Davis, Boston, Mass.
- Fig. 21, 40, Specimens from A. P. Morgan, Preston, Ohio.
- Fig. 22, Specimen in collection of L. M. Underwood.
- Fig. 23, Specimen from Dr. Wm. Herbst, Trexlertown, Pa.
- Fig. 24, 26, 32, 49, Specimens in Ellis' Collection, New York Botanical Garden.
- Fig. 25, Specimen from C. V. Piper, Pullman, Washington.
- Fig. 27, Specimen from L. A. Greata, Los Angeles, Cal.
- Fig. 30, 46, Specimens from Fred. J. Braendle, Washington, D. C.
- Fig. 31, Specimen from Prof. A. J. McClatchie, Phoenix, Arizona.
- Fig. 38, Specimen from Caroline A. Burgin, Philadelphia, Pa.
- Fig. 39, Photograph from Fred. J. Braendle, Washington, D. C.
- Fig. 41, Specimen from E. Bartholomew, Rockport, Kan.
- Fig. 43, Specimen from Mrs. Eugene Wright, Hubbard Lake, Mich.
- Fig. 4, 48, Specimens from Geo. E. Morris, Waltham, Mass.
- Fig. 2, 3, 15, 16, 19, 28, 29, 33, 34, 44, 45, 47, Specimens collected by author.

PLATE 1.



Fig. 15.
Mutinus elegans.



Fig. 16.
Section of egg. *Mutinus elegans.*



Fig. 17.
Section of *Rhizopogon luteolus.*



Fig. 18.
Rhizopogon luteolus

PLATE 2.



Fig. 19. *Nidularia striatus*.



Fig. 20.
Chlamydopus clavatus.



Fig. 21.
Tylostoma verrucosum.



Fig. 22.
Battarreia Griffithii.

PLATE 3.



Fig. 23.
Queletia mirabilis



Fig. 24.
Cauloglossum transversarium



Fig. 25.
Scootium acuminatum

PLATE 4



Fig. 26.

Dictyocephalos curvatus. (Reduced one-third.)

PLATE 5.



Fig. 27.
Gyrophragmium Delilei. (Not perfect, wanting volva.)



Fig. 28.
Arachnion album. a—plant. b—section.



Fig. 29.
Mitromyces cinnabarinus.



Fig. 30.
Mitromyces lutescens.

PLATE 6.



Fig. 31.
Podaxon Farlowii.



Fig. 32.
Catastoma subterranea.



Fig. 33.
Catastoma circumscissa.



Fig. 34.
Bovistella Ohiense.

PLATE 7.



Fig. 35.
Scleroderma verrucosum.



Fig. 36.
Polysaccum crassipes.

PLATE 8.



Fig 37. *Myriostoma coliforme*.



Fig. 39. *Geaster triplex*.



Fig. 38. *Geaster pectinatus*



Fig. 40. *Geaster hygrometricus*.

PLATE 9.



Fig. 41.
Mycenastrum spinulosum. (Section.)




Fig. 42.
Capillitium of *Bovista.* (Magnified 80 diam.)



Fig. 43.
Bovista pila. (Mature.)



Fig 44.
Bovista pila. (Young.)

PLATE 10.



Fig. 45.
Lycoperdon muscorum.



Fig. 47.
Lycoperdon eruciatum.



Fig. 46.
Lycoperdon pseudoradicans.



Fig. 48.
Calvatia elata.

PLATE II.



Fig. 49.
Hypoblema pachyderma

INDEX TO FIGURES
Illustrating Families and Genera.

Family Phalloideae, Plate 1, Fig. 15 and 16.
 Family Nidulariaceae, " 2, " 19.
 Family Hymenogastraceae, " 1, " 17 and 18.

FAMILY LYCOPERDACEAE.

Arachnion,	Plate 5,	Fig. 28.
Battarrea,	" 2,	" 22.
Bovista,	" 9,	" 42, 43 and 44.
Bovistella,	" 6,	" 34.
Calvatia,	" 10,	" 48.
Catastoma,	" 6,	" 32 and 33.
Cauloglossum,	" 3,	" 24.
Chlamydomus,	" 2,	" 20.
Dictyocephalos,	" 4,	" 26.
Geaster,	" 8,	" 38, 39 and 40.
Gyrophragmium,	" 5,	" 27.
Hypoblema,	" 11,	" 49.
Lycoperdon,	" 10,	" 45, 46 and 47,
Mitremyces,	" 5,	" 29 and 30.
Mycenastrum,	" 9,	" 41.
Myriostoma,	" 8,	" 37.
Podaxon,	" 6,	" 31.
Polysaccum,	" 7,	" 36.
Queletia,	" 3,	" 23.
Scleroderma,	" 7,	" 35.
Secotium,	" 3,	" 25.
Tylostoma,	" 2,	" 21.

QK
600
.L79

BULLETIN No. 5. JUNE, 1902. MYCOLOGICAL SERIES, No. 2

BULLETIN
of the
LLOYD LIBRARY
of
BOTANY, PHARMACY AND
MATERIA MEDICA

J. U. & C. G. LLOYD
CINCINNATI, OHIO

MYCOLOGICAL SERIES, No. 2

THE
GEASTRAE

Illustrated with 80 figures

By C. G. LLOYD

GEASTRAE.

We have classed the Geastrae as a sub-tribe of the Lycoperdeae, the essential characters of which are plants sessile, spores mixed with capillitium. (See "The Genera of Gastromycetes, p. 11.)

Geastrae differ from the other Lycoperdeae in having the outer peridium thick, permanent, and when the plant ripens the outer peridium peels away from the inner, splits into segments and becomes more or less recurved or spreading.

THE MYCELIUM.

There are two distinct types of mycelium. Most Geastrae develop under the ground and the mycelial threads proceed from every portion of the outer peridium binding it to the soil. This is the usual type of most Geaster mycelii. In some species however, (see Fig. 57) the mycelium proceeds only from the base of the plant, and has the appearance of large cord-like roots.

THE OUTER PERIDIUM.

There are three distinct layers forming the outer peridium of a Geaster and they are quite evident to anyone who will closely observe them.

- 1st, the mycelial or outer layer,
- 2nd, the fibrillose or middle layer,
- 3rd, the fleshy or inner layer, also called the Collenchyma.

THE MYCELIAL LAYER.—This derives its name from the fact that in many cases in the growing plants, mycelium threads proceed from all parts of it and bind the plant to the surrounding soil. In plants of the section *Rigidae* it is fragile, and so closely attached to the soil that as the plant expands it tears away from the mycelial layer which remains attached to the soil. In herbarium specimens (see Fig. 8) of *Geaster hygrometricus* and others of the *Rigidae*, the outer peridium appears smooth, the mycelial layer having entirely disappeared. In most Geasters however, the mycelial layer remains more or less firmly attached to the fibrillose but the degree of attachment in different specimens, otherwise the same, is of no importance, merely a condition. In *Geaster limbatus*, most of the specimens have the two layers adnate but we have specimens that have the mycelial layer only slightly attached at the extremities of the segments, and specimens also where it has entirely peeled off. In some species, (*fornicatus coronatus*, *radicans* in particular) the mycelial layer remains as a cup, the fibrillose layer separates and arches up, tearing away, except at the tips of the segments which remain attached. Species with this character are called fornicate, and as it seems to have been supposed to have been the character of only one species called *fornicatus*, several have been confused under this name. As a matter of fact quite a number of species have this character in a more or less perfect degree. All Geasters have an outer layer which

for uniformity we call the mycelial layer, though inaccurately so-called in cases like *G. radicans* where the mycelium is basal.

THE MIDDLE OR FIBRILLOSE LAYER.—This is usually the thickest and principal layer of the outer peridium and in many herbarium specimens is the only one that remains, the outer and inner layers having peeled off and disappeared. Its nature varies much in different species. In the *Rigidae* it is firm, thick, strongly incurved when dry, and strongly hygroscopic. When the plant is moist the segments reflex, and they curl in again when dried, and the process can be repeated as often as the plants are moistened and dried. Fig. 5 represents a dried plant as found in the herbarium, Fig. 6 is the same plant after having been moistened. All Geasters are to an extent hygroscopic and the simplest way to make a crushed specimen assume its normal shape is to place it a few minutes in a jet of free steam which puffs them out plump and natural. The photographs of many of the specimens we present would not be supposed to be the same specimen we received. In most species of Geaster the fibrillose layer instead of being firm as in *Rigidae* is to an extent flexible and in the only specimen we have seen of "*G. turbinatus*" it resembles parchment paper.

THE INNER OR FLESHY LAYER.—This layer differs very much from both of the preceding. When the plant opens it is thick, soft, fleshy, usually white or pinkish. As it dries it almost always turns dark reddish brown, dries down to a thin adnate layer, or splits up and peels off entirely or partially. A photograph of a Geaster taken with this layer fresh is quite different from the photograph of the dried specimen of the plant. Sometimes instead of drying down to a thin layer, if exposed to the weather it thickens, becomes spongy, torn. This is particularly the character of the fleshy layer of *G. rufescens*. In many species if specimen of the plant be dried when it first opens, the fleshy layer remains as a thin red adnate layer, whilst if left exposed to the weather the layer peels off and disappears entirely. Specimens collected in these different conditions appear like different plants. Sometimes the fleshy layer separates from the fibrillose, and remains as a kind of cup at the base of the inner peridium. This is purely an accidental character and while present in many specimens (see Fig. 47) is absent in others. It is the basis for such species as *G. triplex*, and made the *key character* in Saccardo. While we consider *G. triplex* a good species, it is on entirely different points from this feature, from which it receives its name.

Fig. 60 shows a specimen of *G. coronatus* in which a portion of the fleshy layer in peeling off has chanced to tear in a circumscissile manner and dried as a separate ring, which being too small to slip over the inner peridium remains as a loose collar at its base. It is needless to say that this is purely accidental and might never occur in another specimen.

THE INNER PERIDIUM.

The inner peridium of Geasters is generally dull, flaccid, soft. It is either globose or more or less ovate, often tapering to the base. Sometimes it is pedicellate, sometimes sessile and this feature is a primary character in several authors' classification. I do not however, feel that it is of primary importance for I think the length of the pedicel, in some species, is dependent largely on the extent that the outer peridium is reflexed or drawn away from the inner. *Geaster rufescens* I believe varies in having the inner peridium sessile or distinctly pedicellate.

THE MOUTHS OF THE INNER PERIDIUM are of three types. 1st, not defined but simply a torn aperture; 2nd, distinct, usually conical, but even. 3rd, strongly sulcate. I think Geasters are more strongly characterized by their mouths than by any other feature. In addition some Geasters have the mouths seated on a definite circular area strongly marked, and differing in shade of color from the balance of the inner peridium. Such mouths we call definite. In others the mouth is conical and distinct but is not marked with a definite area. Such we call indefinite. While the various species are characterized by having in general definite or indefinite mouths we think it is not rare that individual plants of a species usually having indefinite mouths may have a definite mouth or vice versa.

In addition to these characters above we read of "dentate" mouths especially in connection with *G. rufescens*, and such a mouth is clearly shown on Schmidel's drawing. We believe however, that it is purely in error, and as that error has been handed down in our descriptions for 150 years it is time we were rid of it.

We also read of fimbriate mouths, especially in connection with *G. fimbriatus*. Most Geasters of the even-mouthed series have appressed hairs around the mouth, and when the plant is old and weather worn these hairs become frayed and take on a fimbriate appearance, but that it is a character, I do not believe.

We have seen specimens with an even mouth, rimose, and appearing at first sight as if sulcate. That is simply the result of the way the plant dries and its occurrence is rare. It was from such a specimen Schaeffer (1761) drew his figure on which *G. coronatus* (fornicatus of many authors) was based, and hence the error that persists for 140 years that "Geaster fornicatus has a sulcate mouth." No fornicate species of Europe has to my knowledge a sulcate mouth.

"Pectinate" mouth is a term used in connection with Geasters. A pectinate mouth would be composed of narrow segments set parallel like the teeth of a comb. Such mouths are often shown in illustrations, as in Chevallier's cut of "*G. minimus*" and in Masee's beautiful but inaccurate figures of Geasters in the *Annals of Botany*. We do not think that such a mouth occurs in nature but are exaggerated conceptions of sulcate mouths. A plant with a sulcate mouth might have the divisions broken apart and thus become "pectinate," but we have never seen one and do not believe they occur.

CAPILLITIUM.

With the exception of the anomalous species, *hygrometricus*, the capillitium of Geasters consists of long unbranched threads that proceed from the columella and inner surface of the peridium.

The capillitium, in some species at least, is more firmly attached to the peridium and columella than usual in most *Gastromycetes*. Cut open a Geaster, shake out the spores, and with a hand glass abundant capillitium can be seen proceeding from both columella and peridium. Fragments of these threads are mixed with the spores, and these fragments as seen under the microscope are usually simple, cylindrical and tapering. The relative thickness of the threads as compared to the spores, we give in our descriptions as a matter of form. We place little value on it however, as the threads as well as the spores may vary in thickness.

SPORES.

With the exception of the anomalous species, *hygrometricus*, the spores of the species we have examined are very similar, all globose, all slightly warted, all about 3-5 mc. in diameter. Some are slightly larger than others, some slightly rougher than others, but the differences while evident by contrast are not sufficient to determine specific characters. Cooke describes species from Australia with "smooth" spores. We have never seen a perfectly smooth spore in a Geaster. *G. hygrometricus* can be known at once by its large rough spores 8-12 mc. in diameter.

The color of the spore mass of Geasters affords no distinction as it does in other genera. We find no species with pronounced olive or purplish spores. The usual color is a dark brown deepening to black.

COLUMELLA.

In our opinion one of the most striking points of difference between species is the shape of the columellæ, which varies from ovate, globose, or filiform. To study the columellæ however, the plant should be examined just before it expands. After the spores ripen the columellæ usually become indistinct. Vittadini seems to have been the only author who has observed and illustrated the columellæ in his plates.

SHAPE OF UNEXPANDED PLANT.

If we knew the shapes of the unexpanded plants, the best primary division of the genus would be in two sections. Plants with unexpanded forms, globose (see Fig. 41) and plants with unexpanded form, acute (see Figs. 48, 77). Unfortunately, however, we only know the unexpanded form of a few species, simply from lack of observation. We call attention of collectors especially to this point that in gathering Geasters it is particularly important to secure a few unexpanded plants or to make a note of their form. We hope should we issue a second edition of this pamphlet that we may have the data, and not be forced to admit our ignorance on this character of many of the species.

CLASSIFICATION.

The Geastrae consist of only two Genera, Myriostoma with but a single widely distributed species, and Geaster of which we are familiar with 22 species, and know imperfectly several others.

Geaster hygrometricus differs from other species widely in its internal structure. It has no columella, (neither has other species) the capillitium is branched and interwoven and in mature specimens scanty as compared to other species; the spores are larger and approximate the spores of Scleroderma, and the spore mass closely resembles to the eye that of a Scleroderma. In De Bary's Morphology (English, 1887, pp. 313 and 314,) the points are clearly brought out. Morgan (1889) proposed for it the name Astraeus. Desveaux had many years before (1809) proposed the same thing and Corda (Icones Vol. 5) elaborated it, only they retained the name Geaster for this species, proposing to change the other species to Pleastoma. We do not feel that Geaster hygrometricus ought to be separated from other species which it so closely resembles in general appearance that it was for years confused with them, and which to-day frequently requires the use of the microscope to distinguish from other species. We certainly do not think it ought to be put in a different order (we do not use the word natural) as Fischer proposes, and if we did we would not put Nidulariaceae between it and Geaster.

KEY TO GENERA.

Mouths and pedicels several Myriostoma.
 Mouth and pedicel one Geaster.

MYRIOSTOMA COLIFORMIS.

Exoperidium usually recurved, cut to about the middle to six to ten lobes; if collected and dried when first open rather firm and rigid; when exposed to weather, becoming like parchment paper by the peeling off of the inner and outer layers. Inner peridium, subglobose, sup-



Fig. 1.
Myriostoma coliformis.



Fig. 2.
Myriostoma coliformis.

ported on several, more or less confluent, pedicels. Surface minutely roughened; mouths several, appressed fibrillose, round, plain or slightly elevated; Columellae several, filiform, probably the same in number as the pedicels; spores globose, roughened, 3-6 mc.; capillitium simple, unbranched, long, tapering, about half diameter of spores.



Fig. 3.

Myriostoma coliformis (section showing columellae.)




Fig. 4.

Myriostoma coliformis, (spores magnified.)

The inner peridium with its several mouths can be, not inaptly, compared to a "pepper-box." The specific name is derived from the latin *colum*, a strainer, and the old English name we find in Berkeley "Cullenden puff-ball" refers to a cullender (or colander more modern form) now almost obsolete in English but meaning a kind of strainer. This plant is first mentioned by Doody (in the appendix to Ray's Syn. 2nd Ed., 1696.)* Dickson (†) 1785, beautifully illustrated the plant and as it is such an odd plant it has never been confused in literature. Dickson called it *Lycoperdon coliforme*. Persoon (Syn. 1801) refers it to Geaster, and Desveaux (1809) proposed for it the genus *Myriostoma*. At the present day it is generally known as *Myriostoma coliformis*, though some writers (strangely to our mind) still continue to call it *Geaster coliformis*.

Geographical Distribution.

In Europe the plant is reported from England, Holland, Germany, and Poland, and develops abundantly in the sandy plains of Hungary. In England it is a very rare plant. In this country species were sent to Chas. Peck from Colorado. We have it from Dakota and abundantly from Florida.

Specimens in our Collection.

Florida, (many specimens) Mrs. Delia Sams.

Dakota, Black Hills, Prof. T. H. McBride.

Hungary, magnificent specimens, Dr. L. Hollos.

Explanation of Figures.

Figs. 1 and 2 plants natural size; Fig. 3 Section showing columellae; Fig. 4 Spores magnified 450 diameters. Specimens all from Mrs. Delia Sams, Florida. All figures in this pamphlet are natural size unless otherwise specified. All micro-photographs are by Dr. Edward H. Thompson.

(*) The previous citations of Ray to Merrett (1667) is more probably a Geaster.

(†) Fasc. Plant, Cryp. Britanniae.

GEASTER.

Exoperidium thick, divided into sections and usually recurved away from the inner peridium. Inner peridium sessile or stipitate with a single pedicel. Mouth only one. Capillitium (mostly) simple, unbranched. Spores globose, rough. We would divide the genus primarily into two sections.

Rigidae (see following).

Non-Rigidae (see page 14).

SECTION 1. RIGIDAE.

Exoperidium rigid, strongly incurved when dry, strongly hygroscopic.

This section is a very natural division of the genus readily recognized by the rigid incurved exoperidium segments of the dried specimens. All species of Geaster are hygroscopic to a more or less extent, but these are strongly hygroscopic. The mycelium covers the entire young plants and the layer is thin. When the plant expands the mycelium layer tears off and remains as fragments attached to the soil, hence the plants of this section as found in collections are smooth externally, and entirely devoid of mycelial layers.

Spores large, (8-12 mc.)

(1) hygrometricus.

Spores small, (4-6 mc.)

Mouth indeterminate,

(2) delicatus.

Mouth strongly sulcate,

(3) Drummondii.

Mouth definite, even,

(4) mammosus.

1. GEASTER HYGROMETRICUS.

Unexpanded plant globose. Mycelium layer, thin, tearing away as the plant expands. Fibrillose layer thick, rigid, strongly hygroscopic, splitting into six to twenty segments becoming reflexed when the plant is moist; strong incurved and rigid when dry. Flesh layer thin, soon separating and often absent from herbarium specimens. Inner peridium globose, thin, opening by simply a torn aperture; columella none. Capillitium threads long, branched, about half diameter of largest spores. Spores large, globose, rough, 8-12 mc.



Fig. 5.

Geaster hygrometricus (dried specimen.)



Fig. 6.

Geaster hygrometricus (expanded specimen.)

This plant is found in sandy localities and very common in many places. It develops under the ground and is of slow growth. Young plants received from W. N. Suksdorf grew in clumps, (see fig. 10)



Fig. 7.
Geaster hygrometricus (as it grows.)




Fig. 8.
Geaster hygrometricus (unexpanded.)



Fig. 9.
Geaster hygrometricus (section, unexpanded.)



Fig. 10.
Geaster hygrometricus (unexpanded, caespitose plants.)



Fig. 11.
Geaster hygrometricus (spores magnified.)

but that this is exceptional, or usual, we do not know. The young plants are liable to be taken at first for a species of hypogaeal fungi, or on examination under a microscope for an undeveloped Scleroderma. The genus Diploderma was based on unopened specimens of this plant and Cycloderma on unopened specimens of other Geasters. The general resemblance of the spore glass to that of a Scleroderma is close, for the large rough spores are very similar, and the capillitium is so relatively scanty, that when a piece of gleba is pressed on a glass often only spores can be seen. The capillitium however can be readily floated out by method described in foot note on page 7 of "Gastromycetes Genera."

The plant ripens in late summer or fall of the year, and the thick outer peridium splits into segments, sometimes as few as four, sometimes as many as twenty. They are strongly hygroscopic and in moist weather recurve and standing on their tips lift up the inner ball. In

dry weather they closely curve in, clasping the ball, and they will repeat this as often as the conditions become moist or dry. Hence they are called "hygrometricus" and frequently by children "poor-man's weather-glass." Miss Marshall in St. Nicholas states that in the closed condition they are carried along by the wind and applies to them the name of "Fair weather travellers." Plants persist often during winter and one observing them in the spring expanding under the influence of moisture may take them for growing plants. They become "weather-worn," the inner surface of the exoperidium cracked in numerous areas, the surface of the inner peridium frayed and fibrillose. It is a weather-worn specimen that Schweinitz named "Geaster fibrillosus."

Geographical Distribution.

The plant is cosmopolitan. Common throughout Europe, it is more rare in England than on the continent. In this country it occurs from coast to coast and from Canada to Mexico. Locally however, it has never been found in the immediate vicinity of Cincinnati.

Specimens in our Collection.

Massachusetts, Miss Cora Clarke, Mrs. Chas. Cheney, Simon Davis, Walter Deane. *Connecticut*, James B. Rorer. *New York*, Ella K. Hays. *Pennsylvania*, Ellen M. Dallas. *Maryland*, C. L. Shear. *Minnesota*, Minn. Bot. Survey. *Tennessee*, S. F. Corly. *Georgia*, Roland M. Harper. *Florida*, Mrs. Delia Sams, H. C. Culbertson, P. H. Rolfs, C. G. Lloyd. *Colorado*, C. F. Baker. *Washington*, W. N. Suksdorf. *Illinois*, L. H. Watson.

France, N. Patouillard, F. Fautrey. *Tirol*, Rev. G. Bresadola. *Hungary*, Dr. L. Hollos.

Explanation of Figures.

Fig. 5. A plant of our collection from Walter Deane, Cambridge, Mass. Fig. 6. The same plant when moistened. Fig. 7. Photograph of plant in situ, by F. J. Braendle, Washington, D. C. Fig. 8. Unexpanded plant. Fig. 9. Section of same. Fig. 10. A cluster of unexpanded plants, from W. N. Suksdorf, Washington. Fig. 11. Spores magnified 450 diameters.



Fig. 12.

Geaster hygrometricus var. *giganteus*, (unexpanded)

GEASTER HYGROMETRICUS VAR. GIGANTEUS.

A large form, differing from the ordinary plant only as to size, frequently reaches us from the Western States. It is so much larger than the usual plant that we think is entitled to a distinctive name. This large plant does not grow in Europe to our knowledge.



Fig. 13.
Geaster hygrometricus var. giganteus (expanded.)

**Specimens in our
Collection.**

California, L. A. Greata.
Washington, W. N. Suksdorf.
Iowa, T. H. McBride.

**Explanation of
Figures.**

Fig. 12. Geaster hygrometricus var. giganteus, specimen, from L. A. Greata, Los Angeles, Cal.

Fig. 13. The same after expanding by moisture.

2—GEASTER DELICATUS.

Outer peridium thin, smooth, firm, hygroscopic, cut (about $\frac{2}{3}$ deep) to 8–10 segments. Spreading when moist, incurved when dry. Inner peridium subglobose, opening by a plane, indefinite aperture. Columella none. Capillitium slender, interwoven, simple or sometimes slightly branched near the end, slightly thinner than the spores. Spores globose, minutely warted, 5–6 mc.

This elegant little species is known only from the Northwest. It was described by Prof. Morgan from specimens received from Nebraska. Hollos considers this plant a synonym of *G. lageniformis* of which he sends specimens. (*) It seems to me however that the plants while *very close* are different. *Lageniformis* has a protruding mouth. *Delicatus* the mouth is indefinite, plane, merely an aperture, the same as *G. hygrometricus*. We admit that the two plants are very close, probably the same, but for the present would keep them distinct. Had Morgan had access to Vittadini's figure we should not have blamed him for describing the plant he met as a new species. The figure is an



Fig. 14.
Geaster delicatus.




Fig. 15.
Geaster delicatus (unexpanded.)

(*) See Appendix.

elongated, oval plant with a protruding mouth. *Delicatus* is a depressed globose plant with no protruding mouth.

From *G. mammosus* which this plant closely resembles in general, having the same thin hygroscopic peridium, it can be distinguished by its mouth. From small specimens of *G. hygrometricus* with which it agrees as to its mouth, it can be at once distinguished by its thin peridium and small spores.

Specimens in our Collection.

Washington, W. N. Suksdorf. *Nebraska*, Chas. E. Bessey, (given us by A. P. Morgan).

Explanation of Figures.

Fig. 14. *Geaster delicatus* expanded. Fig. 15. Same unexpanded.

a—Specimen from Chas. E. Bessey, Nebraska.

b—Specimen from W. N. Suksdorf, Washington.

3—GEASTER DRUMMONDII.

Exoperidium rigid, hygroscopic, strongly incurved when dry, cut (about $\frac{2}{3}$ deep) to usually ten linear segments. Mycelium and fleshy layers absent in all specimens I have seen. Inner peridium globose, smooth, firm, sessile, having a short, conical, strongly sulcate mouth, not seated on a definite area. Columella linear (?) (*). Capillitium simple, tapering, about thickness of spores in thickest part. Spores globose, rough, 5–7 mc.



Fig. 18. *Geaster Drummondii*.

The little plant is apparently rare. I first received it under the name *striatulus* from Dr. Hollos, Hungary. Afterwards I found it in Ellis' Exs. (No. 110) in Washington, Philadelphia and New York, labeled *mammosus*, (cfr. *Myc. Notes*, p. 71, No. 162, where however, the reference to Ellis' *exsiccatae* is given in error as 109). Hollos who is familiar with this small plant in Hungary, has examined specimens of *G. Drummondii* of Berkeley from Australia, and pronounced it the same plant, only larger specimens. We really see no essential difference in Cooke's description (save size) of the two plants in "Australian Fungi", and we believe Berkeley's illustration in Hooker's *Journal* is this plant. We think there is no question but that Kalchbrenner had the plant in view in his description of *striatulus*, (*Grev.* vol. 9, p. 3,) though he gives a wrong synonym. Henning beautifully illustrates the little plant from South Africa under the name *G. Schweinfurthii*, (*Eng. Bot. Jahrb.* Vol. 14, t. 6, f. 7.)

(*) Very indistinct in specimens examined and not clearly made out.

Geographical Distribution.

Hungary, (Hollos). Australia, (Kalchbrenner). South Africa, (Henning). New Jersey, (Ellis).

Specimens in our Collection.

Hungary, Dr. Hollos.

Florida, Specimens from A. P. Morgan, (I am in some doubt as to these specimens, they are not so typically hygroscopic as all others I have seen.)

Explanation of Figures.

Fig. 18: *Geaster Drummondii*. a—Expanded. b—Unexpanded. Specimens from Dr. L. Hollos, Hungary.

4—GEASTER MAMMOSUS.

Exoperidium thin, rigid, hygroscopic, smooth, divided almost to base into about ten linear segments, often umbilicate at the base as shown in fig. 17b. Inner peridium globose, smooth, sessile, furnished



Fig. 16.
Geaster mammosus, (expanded.)




Fig. 17.
Geaster mammosus (unexpanded.)

with a conical, even, protruding mouth seated on a definite area. Columella short, globose, evident (though indistinct in mature plants). Capillitium simple, tapering, hyaline, often flattened, slightly thinner than the spores. Spores globose, roughened, 3-7 mc. (*)

This plant differs from other hygroscopic species by its *even* conical mouth. The plant was early (1809) beautifully illustrated by Sowerby (t. 401). Fries (1829) gave the name *Geaster mammosus* to some plant, but not to this, as he describes it as drying with the exoperidium reflexed, and refers Sowerby's characteristic plate, doubtfully, to *Geaster hygrometricus*. Chevallier (1836) clearly describes and characterizes its difference from *hygrometricus* by its mouth. He is usually (and justly in our opinion) cited as the author of the name. Vittadini (1843) gave a fine figure of the plant, but strangely in his text states that it is the Friesian interpretation of the plant and "not Chevallier."

(*) Morgan states 5-6 mc. We have specimens from Canada where the spores vary in the same plant from 3 to 7 mc. in diameter. Our English specimens run more uniform, 3 to 4 mc.

Geographical Distribution.

This plant is distributed through Europe and United States. (*)

Specimens in our Collection.

Canada, Wm. Dearness. *Minnesota*, Minn. Bot. Survey. *Iowa*, W. J. Teeters. *Pennsylvania*, Dr. Wm. Herbst. *California*, L. E. Benton, (specimen from A. P. Morgan).

England, Chas. Crossland. *Hungary*, Dr. L. Hollos.

Explanation of Figures.

Fig. 16. *Geaster mammosus*, expanded. Fig. 17. Same, unexpanded. Fig. 18 and 17b. Specimens from Chas. Crossland, England. Fig. 17a. From John Dearness, Canada.

SECTION 2.—NON-RIGIDAE.

This section is readily recognized from the previous by the segments of the exoperidium not drying strongly incurved over the endoperidium. Two species which we include in this section (*Smithii* and *arenarius*) have a strong tendency toward the previous section, but the tips only of the exoperidium segments dry incurved, not the entire exoperidium. We divide the section into two subsections.

Mouths sulcate (see following)

Mouths even, (see page 22.)

SPECIES WITH SULCATE MOUTHS.—NON-RIGIDAE.

Plants of this section are distinguished from the following section by the sulcate (not even) mouths. It is a question if the same plant under different conditions cannot have a mouth that varies, sulcate or even. If that is so then *Geaster Morganii* becomes *G. lageniformis* and *Geaster arenarius* becomes *G. Smithii*. We think while it is possible it is not proven, for our observation is that plants of the same collection have mouths either all sulcate, or all even.

Omitting from discussion at present *G. Morganii* (which differs in being truly sessile and usually saccate) and *G. Smithii*, (which is unique in itself,) there remains in this section *G. pectinatus*, *G. Bryantii*, *G. Schmidelii* and *G. asper*. These four plants no doubt should be truly considered as forms of one species, but as they never run into each other so closely that there is trouble in naming them, we think it better to present them as distinct species. At the same time they have been so confused in literature it is almost a hopeless task to straighten out the tangled threads. All have strongly sulcate beaked mouths, all pedicels either short or long, all exoperidia usually revolute. All are covered partly in the text and partly in citations of Fries 'striatus' and no doubt that conglomerate species of Fries is responsible for the confusion that has since existed.

KEY TO THE SPECIES.

- Mouth long beaked, pedicel slender, inner peridium usually striate beneath,.....(5) *pectinatus*.
Mouth long beaked; pedicel slender; inner peridium with a circular groove beneath,.....(6) *Bryantii*.
Mouth short beaked; pedicel short, thick; peridium neither striated nor grooved,.....(7) *Schmidelii*.
Mouth short beaked; inner peridium short pedicellate, asperate (8) *asper*.
Mouth conical, inner peridium sessile,(9) *Morganii*.
Mouth flattened conical, depressed,.....(10) *Smithii*.

(*) Notwithstanding Masee's statement "The North American specimens under this name are certainly not the true species."

5—GEASTER PECTINATUS.

Exoperidium revolute, cut about to the middle into 8 to 10 segments. Mycelial layer generally adnate, carrying with it soil. Fleshy layer thin, finally peeling off, and partly peeled off in most specimens giving them a ragged appearance. Pedicel slender. Inner peridium subglobose but somewhat tapering into the pedicel and marked with striae at the base, either faintly or strong enough to be called ridges. Mouth strongly sulcate, beaked, or slender conical. Capillitium slightly thicker than spores. Spores globose, rough, 5-6 mc. in diameter.



Fig. 19. *Geaster pectinatus* (large plant)



Fig. 20.
Geaster pectinatus.



Fig. 21.
Geaster pectinatus.



Fig. 22.
Geaster pectinatus (small plant.)

Schmidel (1747) gave four figures (t. 37, f. 11-14) excellently illustrating this plant. Persoon (1801) called these figures *Geaster pectinatus*. Fries mixed it up with three other species under the name *G. striatus* and since Fries' day it has been so badly confused that we can only refer our readers to the references in appendix for details.

Hollos states that *G. pectinatus* is "a fungus of so rare occurrence it was quickly forgotten." It is undoubtedly a rare plant, we do not remember having seen a specimen in any of the Eastern collections, and yet we find we have five different gatherings, in our own collection. Miss Caroline A. Burgin of Philadelphia and Mrs. Delia Sams of Florida are the only collectors of the plant in this country to our knowledge.

Geographical Distribution.

Europe and the United States, rare in both countries.

Specimens in our Collection.

Pennsylvania, Miss Caroline A. Burgin. *Florida*, Mrs. Delia Sams. *Tirol*, Rev. G. Bresadola. *France*, E. Boudier. *Sweden*, L. Romell.

Explanation of Figures.

Fig. 19. Large plant from L. Romell, Sweden. Figs. 20 and 21. Specimens from Caroline A. Burgin, Pennsylvania. Fig. 22. A small plant, specimen from Mrs. Delia Sams, Florida.

6—GEASTER BRYANTII.

Exoperidium similar to preceding species. Pedicel slender. Inner peridium subglobose, or somewhat abrupt at base, marked with a circular groove at the base. Mouths sulcate, beaked. Capillitium and spores as in the preceding.



Fig. 23. *Geaster Bryantii*



Fig. 25. *Geaster Bryantii*.



Fig. 24. *Geaster Bryantii*.



Fig. 26. *Geaster Bryantii*.

This plant is so close to the preceding that I am convinced it might more properly be considered a variety of it. Its distinctive feature the groove at base of peridium, is formed by the pedicel expanding to a disk shape top supporting the inner peridium, which being smaller where it is united forms a groove. It is the original of De Candolle's *Geaster striatus*, particularly as regarding his citations, but he does not mention in his text its distinctive feature, the circular groove. Hence there is a doubt whether he had this plant or the preceding. Fries, as previously stated, confused this plant with three others under the name *Geaster striatus*. Berkeley (Eng. Flo. p. 301) apparently drawing his conclusions from Fries, applied the name *G. striatus* to the preceding plant and renamed this *G. Bryantii*, citing the same references for it that De Candolle had cited for *striatus* with the addition of one citation, (Schmidel, t. 37, f. 11, 12). The last citation is an error, Berkeley having confused a ring shown on the pedicel of the cut, in reality a remnant of the fleshy layer, with the groove that this plant properly has. Berkeley's idea of a distinctive groove, the essential feature, is the first clear conception of the plant and we adopt his name, there never having been any confusion about it. The name *Geaster striatus* which priorists will no doubt use, is subject to the objection in our mind of not having been clearly defined in the first place, and having been applied since to *six different plants* by six different authors. Our specimens show another difference between this plant and *pectinatus*. The peridium is lead color, due to a kind of pruinose covering which may be rubbed off, and usually is on the exposed parts, giving the peridium a variegated appearance as shown in our photographs. (*)

Misconception as to the value of the fleshy layer is the source of at least two species based on this plant. *Geaster orientalis* (Grev. vol. 6, pl. 98, f. 12) is the plant with fleshy layer still remaining and forming "a tube in the shape of a ring at the base of the interior peridium." *Geaster Kunzei* (Winter in Rabenhorst's Flora) is the same plant, the fleshy layer having peeled off, hence "Stiel ohne basale Scheide." I judge from literature that the species is more common than *pectinatus*, yet it has reached me more rarely and fewer specimens.

Specimens in our Collection.

Maine, H. C. Beardslee. *Texas*, W. H. Long, (specimen from C. L. Shear.)
England, E. M. Holmes, Chas. Crossland. *Sweden*, L. Romell.

Explanation of Figures.

Specimens from: Fig. 23, H. C. Beardslee, (from Maine). Fig. 24, L. Romell, Sweden. Fig. 25, Chas. Crossland, England. Fig. 26, E. M. Holmes, England.

(*) The student will note that this is exactly the reverse of statement made by Masee on same subject.

7—GEASTER SCHMIDELII.

Exoperidium revolute, cut to about the middle to usually five to seven segments. Mycelial layer usually adnate. Fleshy layer thin, usually adnate. Inner peridium with a short thick stipe or sessile. Mouth conical, sulcate. Columella large, ovate. Spores small, globose, minutely roughened, $3\frac{1}{2}$ –5 mc.



Fig. 27. *Geaster Schmidelii*.

This little plant is characterized by its small size, and short, thick pedicel. It is probably the plant covered in the text of Fries' *Geaster striatus*, but not his citations. It is the plant we think Chevallier intended to represent as *G. minimus*. (*) We have adopted the name used in the first illustration (Vittadini) that represents accurately this plant, though a large one, and although Vittadini's citations cover other species. The plant seems to be rare and has reached me but rarely.

Specimens in our Collection.

Tirol, Rev. G. Bresadola. *Hungary*, Dr. L. Hollos.
New Hampshire, C. E. Montgomery. (We have seen specimens also from *Vermont* in collection of A. E. Burt, and from *New Jersey* (unlabeled) in collection of N. Y. Bot. Gardens.

Explanation of Figures.

Specimens from: Fig. 27a, J. B. Ellis, New Jersey; b and c, C. E. Montgomery, New Hampshire; d, L. Hollos, Hungary.

8—GEASTER ASPER.

Exoperidium revolute, cut to about the middle to eight to ten segments. Both mycelial and fleshy layers are more closely adherent than in most species. Pedicel *short, thick*. Inner peridium subglobose, *verrucose*. Mouth conical, beaked, strongly sulcate, seated on a depressed zone. Columella prominent, persistent. Capillitium threads simple, long tapering. Spores globose, rough, 6 mc.

The character of this plant is the verrucose inner peridium. Under a glass of low power it appears as though the peridium was densely covered with grains of sharp sand. This plant alone has this character to our knowledge, and although it is indicated in the figures of *G. coronatus* of both Schaeffer and Schmidel, we think there it is an exaggeration of the very *minute* granular appearance *coronatus* has.

(*) Hollos refers this figure to *Geaster asper*.

Geaster asper is on the plate of the first Geasters figured (Michelius, 1729, pl. 100, f. 2), where the plant is characteristically shown, excepting the pedicel is more slender than normal. The word



Fig. 28. Geaster asper.



Fig. 29. Geaster asper.



Fig. 30. Geaster asper.

“asper” is the first descriptive adjective applied by Michelius. Fries included it in his complex striatus. It has been described as a new species in recent times by three authors; as *G. campestris* by Morgan (1887); as *G. Berkeleyi* by Masee (1889); and as *G. pseudomammosus* by Henning in 1900.

Specimens in our Collection.

Hungary, Dr. L. Hollos.
Kansas, E. Bartholomew. *Kentucky*, C. G. Lloyd. *Ohio*, A. P. Morgan,
 (type specimens of *G. campestris*).

Explanation of Figures.

Figs. 28, 29 and 30 (section). Specimens all from A. P. Morgan, Ohio, and the type of “*G. campestris*.”

9—GEASTER MORGANII.

Young plant acute. Exoperidium cut beyond the middle to seven to nine *acute* segments. In herbarium specimens usually saccate but sometimes revolute. Mycelial layer closely adherent, compared to previous species relatively smooth. (*) Fleshy layer when dry, thin closely adherent. Endoperidium globose, *sessile*. Mouth sulcate, indefinite. Columella globose-clavate. Capillitium thicker than the spores. Spores small, globose, 4 mc, almost smooth.

This plant is common around Cincinnati and was referred by Morgan to “striatus.” It is a reddish brown plant and differs widely from other species with sulcate mouths previously described in its

(*) As in the previous species the mycelium covers the young plant but is not so strongly developed so that the adhering dirt is not so evident on the mature plant.

closely sessile endoperidium. It is the same plant as *lageniformis*, indeed Bresadola so refers it, excepting that plant normally has an even mouth, and no other species to our knowledge has mouths in both the



Fig. 31. *Geaster Morganii*.



Fig. 32. *Geaster Morganii*.



Fig. 33. *Geaster Morganii*.



Fig. 34. *Geaster Morganii*.



Fig. 35. *Geaster Morganii*.
(A young plant.)



Fig. 36 *Geaster Morganii*.
(Section of a young plant.)

even and the sulcate series. Still we are convinced of the strong probability of this view and have found in a collection of sulcate mouthed specimens *a single specimen with an even mouth*. It is quite common in this immediate vicinity growing about old stumps and logs, but has never reached me from any other locality in this country or from Europe.

Specimens in our Collection.

Ohio, Mr. Spurlock, W. H. Aiken, C. G. Lloyd.

Explanation of Figures.

Figs. 31, 32 and 33. Specimens from Mr. Spurlock. Figs. 34, 35 and 36. Collected by author; all from immediate vicinity of Cincinnati. Figs. 35 and 36 from fresh specimens, others from dried specimens.

10—GEASTER SMITHII.

Young plant globose. Exoperidium subhygroscopic, cut about half way to 8 to 12 segments, partly reflexed but tips of segments drying incurved. Mycelial layer thin, usually adnate, with adhering sand. Fleishy layer drying thin, adnate. Inner peridium subpedicellate, in reality almost sessile but the outer peridium drawing away from it. Mouth *flattened conical* (or when old conical) *seated on a depressed area, regularly sulcate-striate*. Color of spore mass blacker than in most Geasters. Threads about thickness of spores. Spores globose, rough, apiculate, 4-5 mc.



Fig. 37. Geaster Smithii.

This little plant is unique as to its mouth (well shown in our figures) from all other species. Morgan refers it to *G. umbilicatus* of Fries, and if we draw our conclusions only from what is published we should so refer it. Both Patouillard and Bresadola however, say "not umbilicatus" (*) and they are in better position to know than we are.

This plant was well described and figured by W. G. Smith (in Gard. Chron. 1873, p. 469) under the erroneous name of *G. striatus*. The figures have the mouth more protruding than our cut, but that is a condition of age. His figures show the same *depressed area* characteristic of the plant. He states "the striae of the mouth are so matchlessly perfect and beautiful that no art can do them justice." We believe however, our figure will give a good idea of them.

Being unable to call this plant *umbilicatus* (as did Morgan) or *striatus* (as did Smith) we have named it in honor of Worthington G. Smith, who has done better work with Geasters of England than any other mycologist.

Specimens in our Collection.

Florida, Mrs. Delia Sams.

Explanation of Figures.

Fig. 37. Specimens from Mrs. Delia Sams, Florida.

(*) "Not umbilicatus but a species unknown to me perhaps new."—Bresadola.

"Geaster umbilicatus of modern authors, but I am not certain that it is that species of Fries, and in any case it is *not* that of Montagne, neither of L'éveillé"—Patouillard.

SPECIES WITH EVEN MOUTHS.—NON-RIGIDAE.

(See remarks on page 14 under head of "Species with sulcate mouths.")

The even-mouthed species can be divided into three subsections:

- Exoperidium recurved (not fornicate), (see following).
- Exoperidium fornicate,* (mostly quadrifid), (see p. 29).
- Exoperidium saccate, sessile, (see p. 33).

**EXOPERIDIUM RECURVED, (not fornicate.)
NON-RIGIDAE, MOUTH EVEN.**

The mycelial layer in this subsection is often disposed to separate either entirely or partly adherent (particularly in *limbatus* and *minimus*) but is never truly fornicate as in the following subsection.

KEY TO THE SPECIES.

- Large species,
 - Unexpanded plant globose,
 - reddish brown, sessile or pedunculate,..... (11) *rufescens*.
 - black, pedunculate,..... (12) *limbatus*.
 - Unexpanded plant acute, plant reddish brown, (13) *triplex*.
- Small species,
 - pedicellate, not hygrosopic, (14) *minimus*.
 - subsessile, subhygrosopic,..... (15) *arenarius*.

11—GEASTER RUFESCENS

Unexpanded plant globose. Exoperidium recurved, cut to usually eight segments to about the middle. Mycelial layer, adnate with its adhering dirt or sometimes entirely peeled off. Fleshy layer mostly adnate, *thick*, porous, cracked and having the appearance of rough reddish leather. Inner peridium sessile or usually with a short thick pedicel, somewhat tapering toward the base. Mouth in-



Fig. 38. *Geaster rufescens*.



Fig. 39. *Geaster rufescens*.

(*) The word fornicate meaning arched, as applied to a *Geaster* means *arched over the cup shape mycelial layer*.

definite, fibrillose, frequently torn. (*) Columella large, thick, globose, permanent. Threads thicker than spores. Spores globose, roughened, varying from three to six mc.



Fig. 40. *Geaster rufescens* (section).



Fig. 41. *Geaster rufescens* (unexpanded plant.)

This is the large reddish plant, the most frequent species we have in this country. It is sometimes sessile but usually has a short thick peduncle. The plant from the days of Persoon has been placed in the "sessile" section of the genus, hence when Morgan met the peduncled form he naturally referred it to *limbatus*. *Rufescens* is a reddish brown plant, *limbatus* is a black plant, otherwise they are very close, though *limbatus* has usually a longer peduncle and a different shaped inner peridium. Schaeffer's old figure of the plant shows a regularly toothed mouth and Fries no doubt basing his description largely on this figure, described it as having a toothed mouth. The mouth is often torn but no more frequently than any other species, and the idea that this species can be distinguished by its "dentate peristome" is entirely erroneous, and should be dropped from descriptions.

Specimens in our Collection.

Ohio, A. P. Morgan, (labeled *limbatus*), David L. James, Tom Bell, H. L. True, E. J. Arrick, Tom Lloyd, C. G. Lloyd. New York, Ida M. Hays. Kentucky, Sister Marie. Canada, John Dearness, (spec. tending toward *limbatus*.)

Sweden. L. Romell. England, Carleton Rea. Hungary, Dr. L. Hollos. Tirol, Rev. G. Bresadola.

Explanation of Figures.

Fig. 38. Specimens from A. P. Morgan, Ohio. Fig. 39. Specimens from David L. James, Ohio. Fig. 40. Section, showing large columella. Fig. 41. Unexpanded plant, specimen from Dr. H. L. True, Ohio.

12—GEASTER LIMBATUS.

Outer peridium recurved, cut to about the middle to eight to twelve segments. Mycelial layer usually adnate with its adhering dirt, often partially separate, and sometimes entirely peeled off. Fleishy layer drying firm, hard, and closely adnate. Inner peridium some-

(*) Hence often inaccurately described as "toothed."



Fig. 42. *Geaster limbatum*



Fig. 48. *Geaster limbatum*.



Fig. 44. *Geaster limbatum*.



Fig. 45. *Geaster limbatum*.

times globose rounded at the base (Fig. 42) but usually "slightly constricted and then swollen at the base." (Fig. 45). Pedicel usually distinct—cylindrical (Fig. 43) but sometimes very short and thick, (Fig. 45). Mouth indefinite, fibrillose. Columella indistinct (in ripe specimens at least). Threads thicker than spores. Spores globose, roughened, 4–5 mc.



Plr. 46. *Geaster limbatum* (section)

Geaster limbatum is very close to *G. rufescens*, a fact that seems to have been noted by only one author, Scherffel. (*) The writers who usually place *G. rufescens* in the "sessile" section do not realize that it is so close to *limbatum* that specimens occur that are hard to refer to either species. *G. rufescens* is a reddish brown plant. *G. limbatum* is a black plant but the color distinctions run into each other to an extent. We have never seen *G. limbatum* with the thick porous fleshy layer, usually found on *rufescens*, and we have never seen *rufescens* with the peculiar constricted inner peridium usually (not always) found on *limbatum*. We believe that the prominent, persistent columella of *rufescens* is the characteristic feature which distinguishes it from *G. limbatum*. Any one knowing only extreme forms of *limbatum* such as Fig. 42, from England, and Fig. 45, from Kansas, would be justified in calling them different plants, but our series of specimens shows all grades of connecting forms.

G. limbatum is a frequent plant in this country and in Europe.

Specimens in our Collection.

Kansas, E. Bartholomew. *Iowa*, T. H. McBride. *Wisconsin*, Steve C. Stuntz. *Massachusetts*, F. Le Roy Sargent. *England*, Carleton Rea. *Tirol*, Rev. G. Bresadola. *Hungary*, Dr. L. Hollos.

Explanation of Figures.

Fig. 42. Specimen from Carleton Rea, England. Fig. 43. Specimen from Steve C. Stuntz, Wisconsin. Fig. 44. Specimen from Dr. L. Hollos, Hungary. Fig. 45. Specimen from E. Bartholomew, Kansas. Fig. 46. Section showing indistinct columella.

13—GEASTER TRIPLEX.

Unexpanded plant acute. Exoperidium recurved (or when not fully expanded somewhat saccate at base), cut to the middle or usually two-thirds to five to eight segments. Mycelial layer adnate. Fleshy layer generally peeling off from the segments of the fibrillose layer but usually remaining partially free as a cup at base of inner peridium. Inner peridium subglobose, *closely sessile*. Mouth *definite*, fibrillose, broadly conical. Columella prominent, persistent, elongated (see Fig. 49). Threads thicker than spores. Spores globose, roughened, 3–6 mc.

(*) "*Geaster limbatum* steht dem *G. rufescens* ungewein nahe."

Geaster triplex is a reddish brown color the same as *G. rufescens* with which we think it has been much confused though in reality a very different plant. It is not recorded from England (to our knowledge) and we think English botanists have mistaken it for *rufescens*. As the early figures on which *rufescens* is based show neither of the characters by which that plant is distinguished from *triplex*, it is doubtful if the latter plant is not really the original *rufescens*. The two plants were confused evidently by all the early botanists. The character generally given to distinguish *triplex*, viz :—the remains of the fleshy layer forming a cup at base of inner peridium while usually present should be considered in the nature of an accidental feature and not an essential character of the plant. It is however, the feature from which the plant derives its name, viz :—*triplex*, three fold, three layers. The distinguishing features by which the plant can be known from *rufescens* are, the acute (not globose) young form, the definite mouth, and shape of the columella (see Figs 40 and 49.)



Fig. 47. *Geaster triplex*.



Fig. 48. *Geaster triplex*.
(Beginning to expand.)



Fig. 49. *Geaster triplex*.
(Section.)

Geaster triplex seems to be frequent both in this country and in Europe, though we have no specimens from Europe.

Specimens in our Collection.

Canada; J. Dearness. Minnesota, Minn. Bot. Survey. Ohio, A. P. Morgan. Pennsylvania, Caroline A. Burgin. Massachusetts, G. E. Morris. Connecticut, E. P. Ely.

Explanation of Figures.

Fig. 47. A typically expanded plant, showing the remains of the fleshy layer from which the plant received its name, specimen from A. P. Morgan, Ohio. Fig. 48. A fresh plant beginning to expand, specimen from E. P. Ely, Connecticut. Fig. 49. Section showing columella.

14—GEASTER MINIMUS.

Exoperidium recurved, cut to about the middle to eight to twelve segments. Mycelial layer usually adnate, usually shaggy with adhering fragments of leaves, etc., sometimes partly or entirely separating.



Fig. 50. *Geaster minimus*.



Fig. 51. *Geaster minimus*.



Fig. 52. *Geaster minimus* (section).

Fleshy layer closely adnate, very light color, usually smooth on the limb of the exoperidium but rimose on the segments. Pedicel short but distinct. Inner peridium subglobose or tapering to base, covered with minute granules, usually light colored, but sometimes almost black. Mouth definite, with well marked circular area. Columella slender. Threads slender, equal or thinner than the spores. Spores about 5 mc.

This little plant is the most common small species of *Geaster* we have in this country. It seems to be rarer in Europe where it is usually known as *G. marginatus*. Vittadini's cut accurately represents our plant and the identity of the European plant is well established. There is an earlier *G. minimus* of Chevallier but his figure is doubtful and even if it could be positively identified, it would not be advisable to replace the name so firmly established for the common American plant.

While the specimens in Schweinitz herbarium are normal, he described the plant as having a flattened base, "basi plano."

Morgan reconstructs a cut (Am. Nat. 1884, p. 967) based on this error.

Specimens in our Collection.

Florida, H. C. Culbertson, C. G. Lloyd. *Louisiana*, W. N. Clute. *North Carolina*, Hannah C. Anderson. *South Carolina*, P. H. Rolfs. *Ohio*, W. H. Aiken. *Pennsylvania*, Caroline A. Burgin, Dr. Wm. Herbst. *Michigan*, B. O. Longyear. *Iowa*, T. H. McBride. *Canada*, John Dearness. *France*, E. Boudier. *Tirol*, Rev. G. Bresadola.

Explanation of Figures.

Fig. 50. Specimens from W. H. Aiken, Ohio. Fig. 51. Specimens from Dr. Wm. Herbst, Pennsylvania. Fig. 51. Section.

15—GEASTER ARENARIUS.

Exoperidium subhygroscopic, cut to five to ten segments; drying usually with segments incurved. Mycelial layer closely adnate with adhering sand. (*) Fleshy layer closely adnate, light color, not



Fig. 53. *Geaster arenarius*.



Fig. 54. *Geaster arenarius*.

rimose. Inner peridium subglobose, with a very short but distinct pedicel in some specimens, in others appearing sessile. Mouth even, conical, acute, definite and usually darker colored than remainder of inner peridium. Columella indistinct. Spores globose, rough, 3-4 mc.

This little plant which I collected Feb. 1895, in the sand at Jupiter, Florida, I have never succeeded in getting named. My correspondents have suggested "*G. saccatus*" and "*G. floriformis*," but I am sure it is neither of these. It is very close to *minimus*, differing in its shorter pedicel and more hygroscopic exoperidium. It is still closer to *Smithii*, excepting its mouth.

Specimens in our Collection.

Florida, H. C. Culbertson, C. G. Lloyd, (both from the sand at Jupiter, Florida.)

Explanation of Figures.

All specimens from Jupiter, Florida. The segments of the one closed are more strongly incurved than usual.

(*) One specimen alone we have with the mycelial layer peeled away except at the tips, showing its relation to the fornicate section.

EXOPERIDIUM FORNICATE.—NON-RIGIDAE, MOUTH EVEN.

The word fornicate means arched but as applied to a Geaster means *arched over the mycelial layer which separates and remains as a cup in the ground*. The first two species are thus strongly characterized, the third not to such a strong extent.

KEY TO SPECIES.

- Mouth indeterminate, plant not rooting, (16) fornicatus.
- Mouth indeterminate, plant strongly rooting, (17) radicans.
- Mouth determinate, plant small, (18) coronatus.

16—GEASTER FORNICATUS.

Outer peridium strongly and typically fornicate, the mycelial layer forming a perfect cup at base of plant. Fibrillose layer arched above the cup, to which it is attached by the tips of the segments, cut into four (rarely five) long segments. Fleshy layer partly adherent. Inner peridium distinctly urn shape as shown in our figures (not globose as Masee depicts) tapering below into a short thick peduncle. Mouth indefinite. Columella. (*) Spores globose, almost smooth, 4 mc.

While the very early botanists (Persoon and Buxbaum) distinguished the plant from coronatus, as varieties of same plant, from the day when Fries made his confusing compilation (1829) up to last year, these two plants, so widely different (see figs. 56 and 61) that even the crude cuts are readily distinguished, have been confused by authors in general under the name "fornicatus." We have concluded to retain it (†) for this plant for two reasons. 1st, Hudson who first gave the name to a species of Geaster while confusing as to his citations, evidently knew only this plant, as evidence all tends to the fact that the other (coronatus) probably does *not occur in England*. Every English illustration, (Bryant, Blackstone, Sowerby, Smith, Masee,) represents this plant. It is the only one we have received from England and English botanists advise us it is the only one they know. 2nd, The idea of a "fornicate" species is so strongly connected with the genus Geaster that it should be perpetuated in nomenclature, and applied to the plant that typically represents the idea. This plant which grows only in deciduous woods is much rarer in continental Europe than coronatus that grows common in pine woods, hence the latter plant is the usual species that has been distributed in exsiccatae under the name "fornicatus" These two plants are so distinct that it is strange to us how they could ever have been considered



Fig. 56. Geaster fornicatus.

(*) We do not wish to mutilate by cutting the few specimens we have of this plant.
 (†) This is a reversal of our decision last year (see Myc. Notes, p. 71).

varieties of the same plant much less confused under the same name. Fries not content by including in "fornicatus" two distinct species, further adds to the confusion by ascribing to it a sulcate mouth, a



FIG 56 *Geaster fornicatus*.

character which neither plant has. We do not think that *Geaster fornicatus* has ever been found in this country, and *Geaster coronatus* but rarely. The specimen preserved in the Schweinitz herbarium is neither of these species. (*) We do not know what it is.

Specimens in our Collection.

Hungary, Dr. L. Hollos. *England*, Carleton Rea.

Explanation of Figures.

Fig. 55. Specimen from Dr. L. Hollos, Hungary.

Fig. 56. Specimen from Carleton Rea, England.

(*) We state (*Myc Notes*, p. 77) that this is probably *radicans*, but a re-examination since of the specimen convinces us that it is not.

17—GEASTER RADICANS

Exoperidium typically fornicate, the outer layer separating and remaining as cup at the base, not having mycelium except at the base



Fig. 57. *Geaster radicans*.

where it is strongly developed in a cluster of root-like fibers. Fibrillose layer arched, cut to five (or usually four probably) segments. Fleishy layer thin, dark reddish, closely adherent. Inner peridium subglobose but tapering to the base. Mouth indefinite. Spores globose, almost smooth, 4 mc.

This plant related to *fornicatus*, is strongly different in the basal mycelium, and in the cup having lobes. It enjoys the unique distinction of being the only American species that has never been claimed by any one to grow in Europe. The only specimens we have seen are Rav. exsic. No. 103, and in the collection of Division of Veg. Pathology of Washington, where it was labeled "*fornicatus*." It grew on "a cedar log in Florida," but the collector's name not preserved. All its recorded stations are Southern and we believe it does not grow in our Northern States.

Specimens in our Collection.

Florida, (Kindness of Mrs. Patterson from the Washington collection).

Explanation of Figures.

Fig. 57. Specimen as above.

18—GEASTER CORONATUS.

Exoperidium fornicate, the mycelial layer forming an imperfect cup to which the arched segments of the fibrillose layer are loosely attached at the tips. The cup is not perfect however, as in the two previous species, but the mycelium is so strongly developed that adhering dirt and pine-needles represent an irregular mass rather than a definite cup. Segments of the arched fibrillose layer usually four, sometimes five, deeply cut, but relatively short as compared to the segments of *fornicatus*. Fleishy layer light colored, partially adherent or sometimes entirely peeled off. Inner peridium oblong, tapering to a short pedicel at the base and to an acute mouth at the apex, *covered with minute granular particles*. Mouth definite. Spores globose, roughened, 4 mc.

It is not necessary to repeat here what we have said under *forficatus* in regard to the confusion of these two plants. This plant is much closer to the *minimus* than to *forficatus*. Indeed, its inner pe-



Fig. 58. *Geaster coronatus*.



Fig. 59. *Geaster coronatus*.



Fig. 60. *Geaster coronatus*.



Fig. 61. *Geaster coronatus*.

ridium is the same as *minimus* and specimens, as often found in collections devoid of the mycelial layer, might be referred to *minimus* if attention were not directed to its fewer and deeper lobes of the exoperidium. There is really no name in use that we can apply to this plant free from all objections. Both *coronatus* used by Schaeffer and Scopoli and *quadrifidus* by Persoon, include two plants in the citations. We have adopted the earlier name of Schaeffer because it is quite appropriate, (the plant is not inaptly compared to a crown) and there is no question as to Schaeffer's figure being intended to represent this plant. This species is very common in continental Europe and fre-

quent in collections (usually under the name *fornicatus*). Romell writes me that it is the most common Geaster of Sweden and hence must have been known to Fries, though why he describes the mouth as "sulcate" is strange if he had observed the plant instead of Schaeffer's inaccurate figure. We have never seen but one collection of the plant from this country made by G. E. Morris, of Waltham, Mass

Specimens in our Collection.

Tirol, Rev. G. Bresadola. *Hungary*, Dr. L. Hollos. *France*, F. Fautrey. *Sweden*, L. Romell.
Massachusetts, G. E. Morris.

Explanation of Figures.

Fig. 58. Specimen from G. E. Morris, Massachusetts. Fig. 59. Specimen from F. Fautrey, France. Fig. 60. Specimen from Rev. G. Bresadola, Tirol. Fig. 61. Specimen from L. Romell, Sweden. The collar shown in this figure is an accidental remnant of the fleshy layer and might never occur in another specimen.

EXOPERIDIUM SACCAE.—MOUTH EVEN.

In all the previous species with even mouths the exoperidium when expanded is revolute away from the inner peridium, but in this subsection the base remains as a cup holding the inner peridium. We can readily conceive however, that this would not hold true in all cases, but it is the usual condition that we find in specimens. *Fimbriatus* of Europe while *saccate* in all our specimens is not put into the *saccate* section by Fries. *Lageniformis* while we have never seen specimens not *saccate*, we have of the closely related plant *Morganii* and conceive that if perfectly expanded this would become *revolute* (as Fig. 32). *Velutinus* and *saccatus* are however truly *saccate* species.

KEY TO SPECIES.

- Unexpanded plant globose,
 - Exoperidium splitting into two layers,
 - velutinate, (19) *velutinus*.
 - smooth, (20) *fimbriatus*.
 - Exoperidium not separating, (21) *saccatus*.
- Unexpanded plant acute, (22) *lageniformis*.

19—GEASTER VELUTINUS.

Unexpanded plants globose, sometimes slightly pointed at apex. *Mycelium basal*. Outer layer rigid, membranaceous, firm, light color in the American plant; dark, almost black in the Samoan. Surface covered with short, dense, appressed velumen in the American plant so short that to the eye the surface appears simply dull and rough, but its nature is readily seen under a glass of low power. In the Samoan plant the velumen is longer and plant appears to the eye as densely tomentose. The outer layer separates from the inner as the plant expands and in mature specimens is usually partly free. The thickness and texture of the two layers is about the same. Fleshy layer dark reddish brown when dry, a thin adnate layer. Inner peridium sessile, dark colored, subglobose with a broad base and pointed mouth. Mouth even, marked with a definite circular light-colored basal zone. Columella elongated, clavate. Spores globose, almost smooth, small, 2½-3½ mc.



Fig. 62.



Fig. 63.



Fig. 64.



Fig. 65.



Fig. 69.



Fig. 71.



Fig. 68.



Fig. 70.

GEASTER VELUTINUS.

Explanation of Figures.

Figs. 62, 63 and 64. Expanded plant dried. Fig. 65. Just opening, showing the way two exoperidium layers separate. Figs. 66 and 67. Inner and outer view of a fresh expanded plant. Figs. 68, 69 and 71. Unexpanded plant. Fig. 70. Section of same.

Figs. 62, 63, 64 and 65. Specimens from Hugo Bilgram, Philadelphia. Figs. 66, 67 and 68. Photographs of fresh plants from Samoa. Figs. 69 and 70. From Cincinnati. Fig. 71. Specimen from A. P. Morgan and *type* of "Cycloderma Ohiensis."

This plant has a strange history. As far as we know it was first collected by Morgan in an unexpanded form and sent to Cooke,



Fig. 66. *Geaster velutinus*.



Fig. 67. *Geaster velutinus*.

who hailed it with delight as a re-discovery of the long-lost genus "Cycloderma" (see Grevillea 1882, p. 95) and named it *Cycloderma Ohiensis*. We have "type specimens" of this plant given us by Morgan. We first collected it in the same condition and determined and distributed it as above. In 1898 Hugo Bilgram of Philadelphia, sent us a fine lot of a *Geaster*, new to us but mixed with a number of unexpanded specimens that we recognized as "*Cycloderma Ohiensis*." Comparison with "type" specimen leaves no question. When Morgan found the *expanded* plant he did not recognize the "*Cycloderma Ohiensis*" he had sent Cooke, but described as a new species *Geaster velutinus*. During a trip I made to Samoa (winter of 1899) I gathered a *Geaster* and sent it to Bresadola, which was described in *Myc. Notes*, p. 50, as "*Geaster Lloydii*." The plant was very dark colored, almost black, and densely velutinate, and the mouth is not definite, but a comparison of the specimens now with our American, leaves no doubt in my mind as to their being the same species. We are glad we are not priorists and therefore do not have to adopt the name "*Geaster Ohiensis*" for this plant, although we might write "*Lloyd*" after it; for "*Ohiensis*" was based on a mistake in the first place and is a local name not fitting to a plant that grows in Samoa.

Specimens in our Collection.

Canada, Wm. Dearness. Pennsylvania, Hugo Bilgram. North Carolina, H. C. Beardslee. Ohio, C. G. Lloyd, A. P. Morgan, (Type of *Cycloderma Ohiensis*)
Samoa, C. G. Lloyd, (Type of *Geaster Lloydii*).

GEASTER VELUTINUS VAR. CAESPITOSUS,

A little plant growing densely caespitose, we collected and photographed at Crittenden, Ky. several years ago. We have lost our specimens but have no doubt it was but a small caespitose form of velutinus. The fresh plants were much darker color than the ordinary form, approximating in that respect the plants we collected in Samoa.



Fig. 72. *Geaster velutinus* var. *caespitosus*.
(Unexpanded.)



Fig. 73. *Geaster velutinus* var. *caespitosus*.
(Expanded.)

Explanation of Figures.

Figs. 72 and 73. From fresh plants, Kentucky.

20—GEASTER FIMBRIATUS.

Mycelium universal. Exoperidium cut to six to eight segments about half way, the limb shallow saccate. (*) Outer layer membranaceous, usually separating partially from the inner, the two layers



Fig. 74. *Geaster fimbriatus*.

being very similar as to texture and thickness as in the preceding plant. Fleshy layer when dry, thin, adnate. Inner peridium sessile globose, with an indeterminate fibrillose mouth. Spores globose, almost smooth, 4 mc.

(*) Fries who established the species did not describe it as saccate though if we can depend on the specimens we have, and the figure from Europe it belongs in this section.

This plant which I only know from European specimens I am convinced is practically the same plant as our *saccatus*. (*) With the exception of the indeterminate mouth, and the tendency of the exoperidium to split into two layers I can see no other difference. The idea that *fimbriatus* can be known by its "fimbriate" mouth is an error. The mouth does not differ from several other species with indeterminate mouths. The plant is recorded several times from this country, but I think determinations are based on *saccatus*.

Specimens in our Collection.

France, E. Boudier. *Hungary*, Dr. L. Hollos. *Tirol*, Rev. G. Bresadola.

Explanation of Figures.

Fig. 74a, c and d. Specimens from Dr. L. Hollos, Hungary.

Fig. 75b. From Rev. G. Bresadola, Tirol.

21—GEASTER SACCATUS.

Unexpanded plant globose. Mycelium universal. Exoperidium cut to six to twelve segments about half way, the limb deeply saccate. Mycelial layer adnate to fibrillose. Fleshy layer when dry, thin, adnate. Inner peridium sessile, globose, with a determinate fibrillose mouth. Spores globose, almost smooth, 4 mc.




Fig. 75. *Geaster saccatus*.

Although the plant differs in being more deeply saccate and having a determinate mouth, I believe it is only the American expression of *G. fimbriatus* of Europe. It is a very common little plant in this section, growing gregarious over rich soil and decaying leaves in woods. *Geaster saccatus* is a name given to a South American plant by Fries and applied to our species by apparently universal consent. I do not know however, that anyone really knows that it is Fries' plant. It certainly is not the plant that Spegazzini distributed from South America as *saccatus*.

Specimens in our Collection.

Florida, Mrs. Delia Sams. *Missouri*, N. M. Glatfelter. *Minnesota*, Minn. Bot. Survey. *Illinois*, L. H. Watson. *Ohio*, A. D. Selby, W. H. Aiken. *Kentucky*, C. G. Lloyd. *Pennsylvania*, Ellen M. Dallas. *Mexico*, E. W. D. Holway.

Explanation of Figures.

Fig. 75a. Expanded plant from fresh specimens. Fig. 74b. From dried specimens. Fig. 75c. Reverse view of expanded specimen. All from collection of author.

(*) Bresadola says not.

22—GEASTER LAGENIFORMIS.

Unexpanded plant acute, ovate, (compared to shape of a flask). Mycelium mostly basal. Exoperidium usually saccate. (*) Mycelial layer generally closely adnate, sometimes disposed to separate, often split into parallel lines. (†) Fleshly layer thin, usually peeling off from the segments but remaining on the limb of the exoperidium. Endoperidium subglobose, closely sessile. Mouth conical, definite. (‡) Columella elongated, in dried ripe specimens somewhat subglobose. Spores globose, rough, 5–6 mc.



Fig. 76. *Geaster lageniformis*.



Fig. 77.
Geaster lageniformis.
(Unexpanded, dried.)

The entire plant is a reddish brown. Morgan (in conversation) suggests that it is a depauperate form of *G. triplex*, a view that is not improbable. The expanded plant can with difficulty be told from *G. saccatus*, though segments are more acute. The distinction is in the form of the unexpanded plant.

Specimens in our Collection.

Pennsylvania, Dr. Herbst. *Florida*, Mrs. Delia Sams. *Connecticut*, E. P. Ely.
Minnesota, Minn. Bot. Survey. *Washington*, W. N. Suksdorf.
Germany, P. Magnus.

Explanation of Figures.

Fig. 76. Expanded plant from dried specimens in N. Y. Bot. Garden.

Fig. 77. Unexpanded plant from Minn. Bot. Survey; the shape is no doubt more abruptly acute than the fresh plant would be.

(*) In all our specimens and in Vittadini's figure. Smith (Gard. Chronicle 1873, p. 608) shows it recurved and it probably is so in fully expanded fresh plants.

(†) Mentioned by Morgan as *G. vittatus*.

(‡) In some specimens the entire mouth is lighter color than remainder of endoperidium, in other the mouth is dark but has a light color basal line.

APPENDIX 1.

REFERENCES.

These references are to plants and not to authorities for names of plants. They represent our views of the classification of *plants*. We do not present reference to the ownership or authority for names, as many authors do. Thus our citation under Geaster asper of "Geaster granulatus Cragin in Bull. Washburn" does not indicate that Cragin named a plant "Geaster granulatus." Whether he did, or did not, is of no possible interest to anyone save possibly to Mr. Cragin. The fact however, that he recorded a plant as "Geaster granulatus" which plant is *G. asper* is of interest to every student of Geasters and these facts alone we have endeavored to cite.

We give the names applied to plants since the adoption of the binomial system, and the fact that the *same name* has been applied to so many *different plants* by various authors we think should impress upon the student the importance of turning his attention to the study of plants, rather than the study of names. Previous to the adoption of the binomial system, we have cited no "names" as we consider the polyglot adjectives applied by the pre-Linnaean botanists in the nature of descriptions rather than names. We have given a few references on the authority of Rev. Bresadola (kindly communicated to us in letters), and some on the published work of Dr. Hollos.

Except when stated however, these references represent our views. We have cited very few references save where the plant is illustrated, or where we have seen specimens, for the citations of many authors are so conflicting that it is impossible to state what *plant* they have in view. Where an author gives an illustration of a plant that can be recognized, we accept that figure as representing the plant he had, though it may be in direct conflict to citations that he has made. We have given no bibliography in explanation of these references, and refer those interested to the excellent bibliography of the Gastromycetes given by Masee in Vol. 4 of Annals of Botany.

We feel and hope that most of our readers will study the plants that they meet, and that few will care to puzzle over these references. Those however, who study names of plants, or rather, who study misnames of plants, should be prepared to interpret these references without the aid of a "bibliography."

MYRIOSTOMA COLIFORMIS.

Doody in Ray. Syn. 2nd Ed. App. p. 340,—*Lycoperdon coliforme*, Dickson Fasc. 1, t. 3, f. 4, (good); Sowerby t. 313 (fine); *Gastrum coliforme*, Pers. Syn. p. 131,—*Geaster coliformis*. Smith in Gard. Chron. 1873, p. 469, f. 86; (Reproduced Grev. Vol. 2, t. 15, fig. 1); Masee Monog. Brit. Gast. fig. 66; Fischer in Eng. & Prantl, p. 321, fig. A.

GEASTER ASPER.

Michelius t. 100, f. 2 (more distinctly pedicellate but quite characteristic); Gleditsch Meth. t. 6, (copied from Michelius).—*Lycoperdon stellatum*. Purton Midland Flora Vol. 3, t. 20, (a splendid figure and rarely cited).—*Geaster asper*, Myc. Notes, No. 151; Hollos Term. Füzetek, (1902) p. 120; *Geaster Berkeleyi*, Masee Mon. Brit. Gast. t. 2, f. 41 (poor);—*Geaster campestris*, Morgan's Flora, p. 14; Ellis N. A. F. Exs. No. 1940; Hollos "Kül. a Term. Köz." p. 23, f. 9;—*Geaster granulatus*, Cragin Washburn Bull., p. 40;—*Geaster pseudomammosus*, Henning Hedw. Vol. 39, p. 54, (teste Hollos);—*Geaster pseudostriatus*, Hollos Math. Term. Ert. (1901), p. 505, (Specimen examined, see Appendix p. 43).

GEASTER BRYANTII.

Geaster Bryantii, Berk. Eng. Flo. p. 300; Mass. Mong. Brit. Gast. t. 4, f. 56; Smith Gard. Chron. 1873, p. 505, f. 94; Reproduced Grev. Vol. 2, t. 16, f. 2.—*Gastrum coronatum* var. *Woodwardii*, Pers. Syn. p. 132.—*Geaster calyculatus*, Fuckel Symbolae, t. 5, f. 3; Zopff & Sydow Exs. No. 6; Rabenhorst Exs. No. 2639.—*Geaster Bryantii* form *a fallax*, Scherffel Ber. Deut. Bot. Ges. 1896, t. 19, f. 3 (only);—*Geaster Rabenhorstii*, Haszl. Grev. Vol. 6, t. 98, f. 11.—*Geaster Kunzei*, Winter

Rab. Flora, p. 911.—*Geaster orientalis*, Haszl. Grev. Vol. 6, t. 98, f. 12.—*Geaster fornicatus* var. *multifidus*, Karsten (Spec. in N. Y. Bot. Garden).—Greville states "It is well figured in new series of Flora Londinensis." I have found no other references to this figure.

GEASTER CORONATUS.

Schmidel, t. 37, f. 1 and 2, (mouth not good in either, but both evidently this plant); Buxbaum, t. 28, f. 2, (teste Hollos); *Geaster quadrifidum* var. *minus*. Pers. Syn., p. 133;—*Lycoperdon coronatum*, Schaeffer, t. 183, (figure inaccurate but evident);—*Geaster fornicatus*, Thümen Myc. Univ. Exs. No. 526; Zopff & Sydow, Myc. Marc. Exs. No. 53; Kunze Exs. No. 11; Rabenhorst Exs. No. 2013b; Krieger Fungi Sax. Exs. No. 272; Roumeguere Exs. No. 3635; Winter's Rab. Flora, p. 896, f. 5; Hahn Pilzsammler, t. 29, f. 156; Myc. Notes, No. 153.—*Gastrum quadrifidum*, Pers. Comm., p. 75; Nees Pilze, t. 12, f. 128. (copied from Schmidel);—*Geaster quadrifidus* var. *minor*, Hollos Term. Füzetek, 1902, p. 116. (*)

GEASTER DELICATUS.

Geaster delicatus, Morgan's Flora, p. 17; Ellis' N. A. F. Exs. 2nd Series, No. 1941.

GEASTER DRUMMONDII.

Geaster Drummondii, Berk. in Hooker's Journal, 1845, t. 1, f. 4.—*Geaster striatulus*, Kalch. Grev. Vol. 9, p. 3; Myc. Notes, No. 152.—*Geaster Schweinfurthii*, Eng. Bot. Jahrb. Vol. 14, t. 6, f. 7, (fine);—*Geaster mammosus*, Ellis N. A. F. Exs. No. 110.

GEASTER FIMBRIATUS.

Geaster fimbriatus, Fries' Syst., p. 16 (exc. cit.); Smith Gard. Chron., 1873, p. 543, f. 104; Reproduced Grev. Vol. 2, t. 17, f. 2; Roumeguere Exs. No. 510 and No. 2317; Thümen Myc. Univ. Exs. No. 411; Kunze Fung. Exs. No. 8; Desmazieres' Exs. No. 956; Rabenhorst's Exs. No. 2010b.

GEASTER FORNICATUS.

Battarreia Fung. t. 39, (characteristic): Buxbaum t. 28, f. 1 (teste Hollos). *Lycoperdon fornicatum*, Huds. Fl. Eng., p. 644; Sowerby t. 198, (fine, but segments of exoperidium not relatively long enough); Bryant f. 14-17 (teste Hollos).—*Geaster fornicatus*, Masee Mon. Brit. Gast. t. 2, f. 42, (subject to same criticism as Sowerby's figure); Smith Gard. Chron., 1873, p. 469, f. 87; Reproduced Grev. Vol. 2, t. 15, f. 2.—*Lycoperdon fenestratum*, Batsch Elen. t. 29, f. 168 a. b. (teste Hollos).—*Geaster fenestratus*, Myc. Notes, No. 150.—*Gastrum quadrifidum* var. *fenestratum*, Pers. Syn., p. 133.—*Geaster quadrifidus* var. *major*, Hollos Term. Füzetek (1902) p. 116. (*)—*Geaster Marchicus*, Fischer in Eng. & Prantl, p. 321, fig. B.—*Pleostoma fornicatum*, Corda Icon. Vol. 5, t. 4, f. 43.—*Geaster MacOwani*, Kalch, in Grev. Vol. 10, p. 108.

GEASTER HYGROMETRICUS.

Schmidel t. 28; Michelius t. 100, f. 4, 5 and 6, (the last the best); Gleditsch Meth. t. 6, (copied from Michelius).—*Geastrum hygrometricum*, Pers. Syn. p. 135; Schweinitz Fung. Car. No. 329; Nees Pilze t. 12, f. 127, (copied from Schmidel).—*Geaster hygrometricus*, Fries Syst. p. 19; Smith Gard. Chron. 1873, p. 577, t. 112; Reproduced Grev. t. 13, f. 2; Trelease Trans. Wis. Acad. Vol. 7, t. 7, f. 1, (poor); Winter Rab. Flora, p. 895, f. 1-3;—*Geaster vulgaris*, Corda Icones, Vol. 5, t. 4, f. 42;—*Astraeus stellatus*, Fischer in Eng. & Prantl, p. 341, fig. A, B and C;—*Astraeus hygrometricus*, Morgan's Flora, p. 19;—*Gastrum fibrillosum*, Schweinitz Syn. Car. No. 330, (we have examined the specimen and it is unquestionably an o'd weather-worn specimen of hygrometricus).

(*) The names adopted by Hollos seem very strange in view of the statement in the text "These two fungi are *no varieties* but are two different, independent species."

GEASTER LAGENIFORMIS.

Boccone Mus. t. 301, f. 6; (section of young plant);—*Geaster lageniformis*, Vitt. Monog. Lyc. t. 1, f. 2; Myc. Notes, No. 167;—*Geaster saccatus*, Morgan's Flora, p. 18 (exc. of illustration); Smith, Gard. Chron. 1873, p. 1275, f. 266; Reproduced Grev. Vol. 2, t. 20, (We think the plant Smith took for lageniformis fig. 116, is a form of the plant but not so typically as the plant he called *saccatus*); Trelease Trans. Wis. Acad. Vol. 7, t. 7, f. 2.—*Geaster minutus*, Henning Hedw. Vol. 39, p. 54 (teste Hollos).

GEASTER LIMBATUS.

Schmidel t. 46, (mouths too strongly defined); Ray Syn. 3rd Ed. t. 1, (poor);—*Lycoperdon stellatum*, Sowerby t. 312, (good); *Geaster limbatus*, Fries Syst. p. 15; Hussey Brit. Myc. t. 2, (splendid and shows both slender and thick peduncled forms); Zopff & Sydow Exs. No. 103; Myc. Notes, No. 154.—*Gastrum coronatum*, Pers. Syn. p. 132;—*Gastrum multifidum* var. B—"Pers. Disp. meth. p. 6"—*Geaster pseudolimbatus*, Hollos Math. Term. Ert. 1901, p. 507, (specimens examined, see Appendix p. 43).

GEASTER MAMMOSUS.

Michelius t. 100, f. 3;—*Geaster mammosus*, Chevallier Flo. Paris, p. 359; Morgan's Flora, p. 16; Smith Gard. Chron. 1873, p. 543, f. 105; Reproduced Grev. Vol. 2, t. 19, f. 1; Vitt. Monog. Lyc. t. 1, f. 9, (fine);—*Lycoperdon reoli-gens*, Sowerby, t. 401, (fine).—*Geaster hygrometricus*, Masee, Monog. Brit. Gast. t. 4, f. 70, (His text of *hygrometricus* is correct but his figure is that of *mamosus*);—*Gastrum hygrometricum* var. *anglicum*, Pers. Syn. p. 135.

GEASTER MINIMUS.

Gastrum minimum, Schweinitz Fung. Car. No. 327, (confirmed by examination of his specimen).—*Geaster minimus*, Fries' Syst. p. 16; Morgan's Flora, p. 15; Ravenel Car. Exs. No. 74; Ravenel Amer. Exs. No. 472; Ellis N. A. F. Exs. No. 109; Roumeguere Exs. No. 4549; Thümen Myc. Univ. Exs. No. 13; Myc. Notes, No. 146.—*Geaster marginatus*, Vitt. Monog. Lyc. t. 1, f. 6, (a small but correct figure of the plant);—*Geaster granulosis*, Fuckel (teste Bresadola), "I have just examined original specimens of *G. granulosis* Fuck. and it is *G. marginatus* 'tout à fait.'—Bresadola.—*Geaster Schmidelii*, Roumeguere Exs. No. 3828.—*Geaster Queletii*, Hazsl. (teste Bresadola in letter).—*Geaster Cesatii*, Rabenhorst (teste Bresadola in letter.)

GEASTER MORGANII.

Geaster Morganii, Myc. Notes, No. 168.—*Geaster striatus*, Morgan's Flora, p. 17; Ellis' N. A. F. 2nd series, No. 2736.—*Geaster saccatus*, Morgan's Flora, Plate 1, f. C.

GEASTER PECTINATUS.

Schmidel t. 37, figs. 11, 12, 13, 14, (the "rings" shown in fig. 11 has caused this figure to be referred, (erroneously) to *Bryantii*);—*Gastrum pectinatum*, Pers. Syn., p. 132;—*Gastrum multifidum* var. a, "Pers. Disp. Meth. p. 6."—*Geaster limbatus*, Smith Gard. Chron. 1873, p. 504, f. 95; Reproduced Grev. Vol. 2, t. 17, f. 1.—*Geaster Schmidelii*, Masee Mon. Brit. Gast. t. 4, f. 74; Winter Rab. Flora, p. 910;—*Geaster Bryantii forma fallax*, Scherffel Ber. Deut. Bot. Ges. 1896, t. 19, f. 1, 2 and 4, (not 3); *Geaster tenuipes*, Myc. Notes, No. 155.

GEASTER RADICANS.

Geaster radicans, Ravenel Exs. No. 103; Myc. Notes, No. 159.

GEASTER RUFESCENS.

Schmidel t. 43 (mucellial layer inaccurately shown; the "dentate" mouths of this figure are responsible for this erroneous idea in connection with the species). Schmidel t. 43 (cont. on t. 50).—*Gastrum rufescens*, "Pers. Disp. meth.

p. 6"; Pers. Comm. p. 74; Pers. Syst. p. 134; Schweinitz' Fung. Car. No. 328. (the specimen in his collection is typical but sessile).—*Geaster rufescens*, Fries' Syst. p. 18; Smith Gard. Chron. 1873, p. 577, f. 111, (Reproduced Grev. Vol. 2, t. 19, f. 2).—*Lycoperdon recolligens*, Sowerby t. 80, (Usually here referred but I think more probably fimbriatus).—*Lycoperdon stellatum*, Sowerby in index to same figure; Schaeffer t. 182, (mouth very poor).—*Lycoperdon sessile*, Sowerby in text under t. 401, (referring to fig. t. 80).—*Geaster multifidum*, Grev. Flo. t. 306, (the expanded plant has the fleshy layer gone and endoperidium distinctly peduncled, the unexpanded plant is *globose*).—*Geaster limbatum*, Morgan's Flora, p. 15, plate 1, f. B.; Ellis' N. A. F. Exs. No. 1309.—*Geaster mammosus*, Rabenhorst's Exs. No. 814.—*Geaster Schaefferi*, Vitt. Monog. Lyc. t. 1, f. 1, (a small plant).

GEASTER SACCATUS.

Geaster saccatus, Ellis & Ev. Fung. Col. Exs. No. 1217; Mvc. Notes, No. 162.—*Geaster lageniformis*, Morgan's Flora, p. 19.—*Geaster capensis*, Thümen Myc. Univ. Exc. No. 715; Roumeguere Exs. No. 4548.

GEASTER SCHMIDELII.

Geaster Schmidelii, Vitt. Monog. Lyc. t. 1, f. 7.—*Geaster Rabenhorstii*, Trelease Trans. Wis. Acad. Vol 7, t. 7, f. 3; Kunze Exs No. 10; Rabenhorst Exs. No. 2011; Zopff & Sydow Exs. No. 7.—*Geaster striatus*, Peck's 38th Rep. p. 94, (teste Trelease).

GEASTER SMITHII.

Geaster striatus, Smith Gard. Chron. 1873, p. 469, f. 88. (Reproduced Grev. Vol. 2, t. 16, f. 1.)—*Geaster umbilicatus*, Morgan's Flora, p. 16, (exc. reference to Ellis' Exc).

GEASTER TRIPLEX.

Michelius t. 100, f. 1, (Fries refers this to fimbriatus, Smith to Michelianus).—*Geaster triplex*, Morgan's Flora, p. 18; Ellis N. A. F. Exs. No. 2735; Thümen Exs. No. 1410.—*Geaster cryptorhynchus*, Hazslinszky Grev. Vol. 3, p. 162, t. 47.—*Geaster Pelotii*, Rose (teste Bresadola).—"Geaster stellatus Linn." Morgan in Jour. of Mycology, Vol. 8, p. 4. (*)

GEASTER VELUTINUS.

Geaster velutinus, Journ. Cin. Soc. Nat. Hist. Vol 18, p. 38; Cycloderma Ohiensis Grev. Vol. 11, p. 95.—*Geaster Lloydii*, Myc. Notes, No. 117.

(*) Linnaeus' idea of "*Lycoperdon stellatum*" was simply the genus *Geaster* as we now know it. He did not know any *species* of *Geasters* and referred to "*Lycoperdon stellatum*" every figure of a *Geaster* he found, some half dozen different species. It is absurd in our mind to attempt to replace an established name of a species of *Geaster* on the *authority of Linnaeus*, a man who had no idea of any *species* of *Geaster*. Michelius who wrote many years before Linnaeus, had definite ideas of a few *Geasters*, but Linnaeus did not know enough of the subject to avail himself of the work of Michelius. Linnaeus apparently was not acquainted with the work of Schmidel, a pre-Linnaean botanist, who well illustrated several species.

APPENDIX 2.

SPECIMENS FROM DR. HOLLOS.

Since most of this pamphlet has been in type we have received from Dr. Hollos, Hungary, three specimens of Geasters.



Fig. 78.

Fig. 78, a little plant which Dr. Hollos sends as *G. floriformis* of Vittadini and considers same as *G. delicatus* of Morgan. We consider both of these views probable but neither proven. If it is *G. delicatus* then our idea of mouth of *G. delicatus* is wrong, for Hollos' specimen has a distinct and protruding mouth as shown in our figure, and we have always supposed *G. delicatus* to have a mouth not protruding, being merely an aperture. It is possible that these views, drawn from all specimens we have seen are wrong and that the mouth of *G. delicatus* when perfect is protruding as shown in fig. 78. In two of Hollos' specimens the mouths were worn off and the specimens could well be taken for *G. delicatus*. Let us hope that Messrs. Bessey, Piper, and Suksdorf who collect this plant will notice this point particularly during the present season.

As to the plants being *G. floriformis*, they do not agree with Vittadini's figure in two particulars. They are depressed globose; Vittadini shows an elongated plant. Their mycelium is evidently universal; Vittadini shows the mycelium basal, though this to our mind is probably erroneous as we doubt if any of the *rigidae* have basal mycelium.



Fig. 79. Geaster pseudostriatus.



Fig. 80. Geaster pseudolimbatum.

Fig. 79 a plant that Dr. Hollos has recently described as a "new species," *G. pseudostriatus*. To our mind it is *G. asper* and differs but little from the form we have in this country. The longer peduncle, we consider only a condition, not an essential character.

Fig. 80 a plant that Dr. Hollos has recently described as *G. pseudolimbatum*. We should call it *G. limbatum*.

.....INDEX.....

	PAGE
Myriostoma coliformis	6
Geaster arenarius	28
“ asper	18
“ Bryantii	16
“ coronatus	31
“ delicatus	11
“ Drummondii	12
“ fimbriatus	36
“ fornicatus	29
“ hygrometricus	8
“ lageniformis	38
“ limbatus	23
“ mammosus	13
“ minimus	27
“ Morganii	19
“ pectinatus	15
“ radicans	31
“ rufescens	22
“ saccatus	37
“ Schmidelii	18
“ Smithii	21
“ triplex	25
“ velutinus	33

QK
600
.L79

BULLETIN No. 8. APRIL, 1905. MYCOLOGICAL SERIES, No. 3

BULLETIN
of the
LLOYD LIBRARY
of
BOTANY, PHARMACY AND
MATERIA MEDICA

J. U. & C. G. LLOYD
CINCINNATI, OHIO

MYCOLOGICAL SERIES, No. 3

The Lycoperdaceae of
Australia, New Zealand and
Neighboring Islands

Illustrated with 15 Plates and 49 Figures

By C. G. LLOYD

INTRODUCTORY.

Australia is the richest country in the world in Lycoperdaceae, and more strange and endemic genera are found there than in any other continent. Our knowledge of the subject is based on relatively scanty material. Probably not more than a hundred specimens have reached the museums of Europe and most of these are at Kew. A majority of the species are known only from a single collection. I do not feel that the knowledge we have of the subject is more than introductory. The work that has been done with the Lycoperdaceae of Australia is mostly sporadic, scattered descriptions of supposedly "new species" by authors who desire to attach their names to them. The only systematic work is in Cooke's Handbook of Australian Fungi, which is a very complete compilation of this sporadic work. The author of this pamphlet has spent fourteen months in the museums of Europe in a systematic study of all the material to be found there, and he has been enabled to study practically all the type specimens on which descriptions of Australian species have been based. In addition we have solicited our correspondents to send us specimens and desire to acknowledge our indebtedness to the following who have kindly forwarded specimens:

MISS JESSIE DUNN, Wellington, N. Z.
F. M. READER, Warracknabeal, Victoria.
J. T. PAUL, Grantville, Victoria.
W. R. GUILFOYLE, Melbourne.
ROBERT BROWN, Christ Church, N. Z.
R. T. BAKER, Sydney, Australia.
J. G. O. TEPPER, Norwood, S. Australia.
ROBERT M. LAING, C.
J. S. TENNANT, Ashburton.
WALTER GILL, Adelaide.
W. W. WATTS, Sydney.

We are also advised of a shipment from D. McALPINE, Melbourne, sent to our Paris address (107 Boulevard St. Michel) but at the time this pamphlet was written the package had not reached us. Specimens received are acknowledged in detail under the species to which they belong.

CORRECTIONS.

The plates and the first form were printed in the absence from home of the author. Several mistakes have occurred.

- Secotium melanocephalum page 7, correct to melanosporum.
- Phellorina Delastrei page 10, correct to Delestrei.
- Scleroderma verrucosum, Plate 31, correct to verrucosum.
- Castoreum radicans, Plate 38, correct to radicans.
- Scleroderma aurantiacum, Plate 31, correct to aurantium.

C. G. LLOYD,

Paris Address:

107 Boulevard St. Michel, - - - Paris, France.



Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.



Fig. 6.

Explanation of Figures.

Figs. 1, 2, 3 and 4. Plants from Robt. Brown, New Zealand. Figure 5. A section enlarged fourfold. Figure 6. Spores (x 1000).



Fig. 7.



Fig. 8.

Explanation of Figures.
 Fig. 7. Plant from W. H. Long, Jr. Texas, U. S. A. Fig. 8. Spores (x 1000).

SECOTIUM COARCTATUM.



Fig. 9.



Fig. 10.



Fig. 11.



Fig. 12.

Explanation of Figures.
 Fig. 9 and 10. Type specimens at Kew. Fig. 11. Section. Fig. 12. Spores (x 1000).

SECOTIUM MELANOSPORUM.



Fig. 4.



Fig. 5.

Explanation of Figures.

Fig. 4. Plant in Museum at Berlin. Figure 5. Spores (x 1000).

PODAXON MUELLERI.



Fig. 1.



Fig. 2.



Fig. 3.

Explanation of Figures.

Fig. 1 and 2. Plants from R. T. Baker, Sydney, Australia. Figure 3. Spores (x 1000).

.K
00
179



Fig. 1.



Fig. 2.

Explanation of Figures.

Fig. 1. Plant from Algiers sent by A. Acloque, France. Fig. 2. Specimen at Kew from Australia. (Two inches of the stipe of the specimen is cut off from this figure).

PELLORINA DELASTREI.



Fig 3.

Explanation of Figure. Digitized by Google
Fig. 3. Type Specimen in Museum at Berlin.

PHELLORINA STROBILINA.



Fig. 2.



Fig. 3.

Explanation of Figures.

Fig 2. Plant from L. G. Yates, California. Fig. 3. Section of same.

BATTARREA STEVENII.

Q.K
600
.L79

Issued by C. G. LLOYD.

PLATE 28.



Fig. 1.

Explanation of Figure.

Fig. 1. Specimen from L. A. Greata, California.

BATTARREA PHALLOIDES.

-K
300
-79

Issued by C. G. LLOYD.

PLATE 29.



Fig. 1.



Fig. 2.

Explanation of Figures.

Fig. 1. Plant from New Caledonia, (from P. Hariot, Paris). Fig. 2. Plant from Walter Gill, Australia.

POLYSACCUM PISOCARPIUM



Fig. 5.



Fig. 6.

Explanation of Figures.

Fig. 5. Plant from Saxony in Museum at Berlin. Fig. 6. Plant from J. T. Paul, Australia.

POLYSACCUM TUBEROSUM. Digitized by Google



Fig. 3.



Fig. 4.

Explanation of Figures.

Fig. 3. Plant from L. G. Yates, California. Fig. 4. Plant from R. T. Baker, Australia.

POLYSACCUM CRASSIPES.

211
600
.L79



Fig. 1.



Fig 2.



Fig. 3.

Explanation of Figures.

Fig. 1. Mature specimen from A. P. Morgan, Ohio. Fig. 2. Young (unopened) specimen from Simon Davis, Massachusetts. Fig. 3. Section of same.

SCLERODERMA GEASTER.



Fig. 4.



Fig. 5.



Fig 6.

Explanation of Figures.

Fig. 4. Unopened plant from J. B. Ellis, New Jersey. Fig. 5. Same opened. Fig. 6. Specimen from W. R. Guilfoyle, Australia.

SCLERODERMA FLAVIDUM.

K
50
79



Fig. 1.

Explanation of Figure.

Fig. 1. Specimen from Steve C. Stuntz, Wisconsin.
SCLERODERMA CEPA.



Fig. 2.



Fig. 3.



Fig. 4.




Fig. 5.

Explanation of Figures.

Fig. 2 and 3. Plant collected at Cincinnati. Fig. 4. Section of same.
Fig. 5. Section enlarged threefold to show the permanent cells.

SCLERODERMA TEXENSE



Fig. 6.



Fig. 7.

Explanation of Figures

Fig. 6 and 7. Plants from Dr. Wm. Herbst, Pennsylvania.
SCLERODERMA AURANTIACUM.



Fig. 8.



Fig. 9.

Explanation of Figures.

Fig. 8. Plant from Charles Crossland, England. Fig. 9. Plant from
Simon Davis, Massachusetts.

SCLERODERMA VERRUCOSUM.

K
00
-79



Fig. 1.



Fig. 2.



Fig. 3.

Explanation of Figures.

Fig. 1. Specimen at Kew, Fig. 2. Section of same. Fig. 3. Spores (x 1000).

CATASTOMA HYPOGAEUM.



Fig. 4.



Fig. 5.



Fig. 6.

Explanation of Figures.

Fig. 4. Specimen from R. T. Baker, Australia. Fig. 5. Section. Fig. 6. Spore (x 1000).

CATASTOMA ANOMALUM.



Fig. 7.




Fig. 8.

Explanation of Figures.

Fig. 7. Type Specimen at Kew. Fig. 8. Spores (x 1000).

CATASTOMA MUELLERI.



Fig. 9.



Fig. 10.




Fig. 11.

Explanation of Figures.

Fig. 9. Type specimen at Kew. Fig. 10. Section. Fig. 11. Spores (x 1000).

CATASTOMA HYALOTHRIX.

K
00
.79



Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.

Explanation of Figures.

Specimens from J. T. Paul, Australia. Fig. 5. Spores (x 1000).

BOVISTELLA AUSTRALIANA.



Fig. 6.



Fig. 7.



Fig. 8.



Fig. 10.



Fig. 9.

Explanation of Figures.

Fig. 7. Type in Museum at Paris. Fig. 8 & 9. Specimens from W. W. Watts. Sydney, Australia. Fig. 6. Plant enlarged 4 times. Fig. 10. Capillitium (x 100).

BOVISTELLA ASPERA.



Fig. 1.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.



Fig. 6.

Explanation of Figures.

Fig. 1 and 2. Plants from W. R. Guilfoyle, Australia. Fig. 3, 4, 5 and 6. Plants from Miss Jessie Dunn, New Zealand.

LYCOPERDON POLYMORPHUM.

QK
600
.L79



Fig. 7.



Fig. 8.



Fig. 9.



Fig. 10.



Fig. 11.



Fig. 12.

Explanation of Figures.

Fig. 7. Young plant with cortex. Fig. 8, 9, 10, 11 and 12. Mature plants
Fig. 10. Section. All from Robert Brown, New Zealand.

LYCOPERDON PRATENSE.



Fig. 1.

Explanation of Figure.

Mature plant from vicinity of Cincinnati, Ohio.

CALVATIA LILACINA.

2K
600
L79



Fig. 2.



Fig. 3.



Fig. 4.

Explanation of Figures.

Fig. 2. Specimen from J. G. O. Tepper, Australia. Fig. 3. Plant in Museum at Berlin. Collected by Dr. Hennings, near Berlin. Fig. 4. Plant from Dr. Hollos, Hungary.

CALVATIA CANDIDA.



Fig. 5.

Explanation of Figure.

Fig. 5. From type specimen at Kew.

CALVATIA OLIVACEA.



Fig. 1.



Fig. 2.

CALVATIA CAELATA.

QK
600
.179



Fig. 3.




Fig. 4.


Fig. 1. Plant from T. de Aranzadi, Spain. Fig. 2. Plant from E. Bartholomew, Kansas. Fig. 3. Plant from Robert Brown, New Zealand. Fig. 4. Plant from C. V. Piper, (state of) Washington.

CALVATIA CAELATA.

QK
600
.L79

Issued by C. G. LLOYD.

PLATE 3



Explanation of Figure.

A small plant collected near Cincinnati, Ohio.

CALVATIA GIGANTEA.

7.



Q K
600
L79

Issued by C. G. LLOYD.

PLATE 38.



Fig. 1.



Fig. 2.

Explanation of Figures.

Fig. 1. Type specimen at Kew. Fig. 2. Spores (x 1000).

CASTOREUM RADICATUM.

QK
600
L79

Issued by C. G. LLOYD.

PLATE 39.



Fig. 1.



Fig. 2.



Fig. 3.




Fig. 4.



Fig. 5.

Explanation of Figures.

Fig. 1. Plant in exoperidium. Fig. 2 & 3. Section of endoperidium, showing core. Fig. 4. Capillitium (x 100). Fig. 5. Spores (x 1000). All from types at Kew.

MESOPHELLIA ARENARIA.

QK
600
.L79

UNIV. OF MICH:

JUN 10 1910

BULLETIN No. 13. SEPT., 1909. MYCOLOGICAL SERIES, No 4

BULLETIN
of the
LLOYD LIBRARY
of
BOTANY, PHARMACY AND
MATERIA MEDICA

J. U. & C. G. LLOYD
CINCINNATI, OHIO

MYCOLOGICAL SERIES, No. 4

Synopsis
of the
Known Phalloids
With an Illustration of Each Species

By C. G. LLOYD



Ed. Fischer.

In this pamphlet devoted to phalloids I am pleased to present a photograph of Professor Ed. Fischer, Bern, Switzerland, who is the best authority in the world on the phalloid subject.—C. G. L.

INTRODUCTION.

Phalloids are in many respects the most remarkable fungi that grow. Usually they are excessively fetid, and persons who would pass by an ordinary fungus without noticing it have their attention strongly fixed when they chance upon one of these "ill-smelling things." In addition they assume most bizarre shapes and often bright colors. I hope these features, probably intended by nature to attract flies, will attract the attention of those to whom this pamphlet is sent.

From the very nature of phalloids, they should be studied in the countries where they grow. Accurate work can not be done in Europe with such fugitive plants, and a large part of what has been written on the subject is not reliable. More has been added to our knowledge by the observations of Penzig, Moeller, Petch, Long, and Cobb, in very recent years, than from all other sources, and these men observed and studied the phalloids in the countries where they grow.

It was with the view of summarizing what is known of the phalloid subject and making it available to students in all parts of the world that this pamphlet has been written. We hope to interest observers in such unworked fields as India, Japan, Australia, West Indies, and South America (except portions of Brazil). We should be glad if any observer in any country where the phalloids are not well known (cfr. page 6) would publish with good photographs an account of such species as he observes. We believe that all the well-known species can be readily determined from this pamphlet.

We trust, however, that this will not lead to a flood of "new species" by inexperienced observers. The species of phalloids, like all fungi, are widely distributed, and wherever you may be located most of the phalloids you will find are recorded in this work. They may differ in unimportant details and seem new to you, but we strongly advise you before publishing to first submit a good photograph, color notes, and a dried specimen to Professor Ed. Fischer or to myself for an opinion.

C. G. LLOYD,
63 rue Buffon, Paris, France.

WHAT IS A PHALLOID?

It would be out of place in a work of this kind, intended for general distribution, to enter upon any technical, botanical discussion of what constitutes a phalloid. Most persons know them by reputation, and with certainty if they have met them. If not, they will know them as soon as they look through our pictures.

Phalloids are always fleshy fungi, always fetid, and appear as if by magic in our woods and fields. When young, they are enclosed in a gelatinous membrane called a volva, which breaks, and the plant develops so rapidly that I will not go into details for fear that some of my readers will think I am not telling the truth. I have often carried home the eggs, but have never seen them develop, as my specimens have always developed during the night. In a single night the species observed have reached a height of eight inches. One author has a picture showing a plant to have grown 4 cm., or an inch

and a half, in one minute of time. Of course this is not true growth by the accretion of cells, but rather a mechanical process by the expansion of cells.

The "roots," or mycelium, as it is correctly called, of phalloids grow in the earth, or rotten wood, and take the form of long, white cords. The illustration on the opposite page is a cluster of this mycelium, which has developed several "eggs," or young phalloids. If we cut open one of these eggs we will find it to contain an undeveloped plant, as shown in the figures herewith. But it is best not to cut

them open, but to take them home and place them on a dish, and in a few days you will have some perfect plants.

THE COLOR OF PHALLOIDS.

There are only three colors known in the phalloids: red, yellow, and white. Most species are red, or some shade of red, pink, flesh, or orange. A few are yellow, and many are white. The yellow and red phalloids seem quite distinct, and do not run into each other, but the red species are apt to have white forms.





DEFINITION OF TERMS.

In the description of phalloids it is necessary to use a few botanical terms, but they are simple and will be readily understood from the following explanation.

VOLVA.—All phalloids (excepting one genus, *Phallogaster*) when young are enclosed in a subglobose membrane called the volva. In this state a phalloid can well be compared to an egg; in fact, it is customary to speak of young phalloids as “eggs.” The volva or shell, however, is a soft, thick, gelatinous membrane. When the plant develops the volva bursts at the top and remains as a cup at the base of the mature phalloid. All our pictures of phalloids show the volva at the base of the plant, at least all pictures that were made from perfect plants. If there is no volva at the base it is because the illustration was drawn from an imperfect specimen.

RECEPTACLE.—This is a term that is applied to the portion of the plant that bears the greenish, mucilaginous mass (called the gleba). In some phalloids (such as *Clathrus*) the entire plant, exclusive of the volva, forms the receptacle. In others, such as *Simblum*, the receptacle is borne on a stem. Some phalloids are a simple, stem-like structure and bear the gleba directly on the upper portion, then of course the upper portion of the stem is the receptacle.

GLEBA.—This is a greenish, viscid, fetid substance with which all phalloids are supplied. It is in fact the fruiting portion of a phalloid, for it contains innumerable, microscopic spores which are analogous to the seed of flowering plants. It is the gleba of a phalloid that is usually so excessively fetid. This bad odor, as offensive as it may be to us, serves a useful purpose to the plants, as it attracts flies and other insects that are the means of the dispersion of the spores.

STEM.—The stem (or stipe) of a phalloid needs no special explanation. It is used in the ordinary sense of the word. Some phalloids have no stems.

PILEUS.—There are some phalloids (the genus *Phallus*) that have the gleba borne on a special membrane on the top of the stem. This is usually conical or hat-shaped and is called the pileus.

VEIL.—A most striking feature in a few species that have pilei is a thin, net-like membrane that hangs from under the pileus and spreads out as a net around the stem. It is called the veil (or more correctly the indusium) but we prefer to call it the veil.

HISTORY OF PHALLOIDS.

We can not write the history of the phalloids because it is not known. There are only five countries in the world where the phalloids are well known, viz: Europe, the United States, Brazil, Java, and Ceylon. Most of the mycological writers have lived in Europe and the United States, and the easy, conspicuous fungi such as the phalloids are well known. In Java most excellent accounts of the phalloids have been written by Penzig, and in Brazil by Moeller. In the United States a good account of the phalloids of Texas was published by Long, and in Hawaii by Cobb.¹ Very recently—in fact, since this pamphlet was in the printer's hands—we have had an excellent account of the phalloids of Ceylon, by T. Petch. Aside from these five papers, however, most of the work on the subject has been in the line of new species exploitation.

¹ Mr. Cobb marred his paper by discovering some “new species” that were only new to him; otherwise, his paper was most excellent.

If a census were taken of the individual specimens that have reached Europe from foreign countries, probably more than one-half have been discovered to be "new species." Most of these new species finally gravitate where they belong, into the trash pile known as synonyms. Professor Fischer, of Berne, Switzerland, has done good work in disposing of a great many of them. We shall help the subject along to the best of our ability in one of our appendices.

NAMES OF PHALLOIDS.

Like all objects of natural history, phalloids have Latin names, and in addition to each is usually appended a personal name, primarily designed to tickle the vanity of some individual. Under this system they have never acquired any stable names, for each person who writes about them is chiefly interested in getting up new names to which to append his own. By this means the names of phalloids (like all fungi) have been shuffled about like a shuttlecock. There are only forty-nine phalloids that are at all well known, and fifty-eight more or less vague and often inaccurate accounts and forms. These one hundred and seven species have two hundred and ninety-nine different names. One of them alone, *Phallus indusiatus*, has twenty-four different names. It is customary in "scientific" monographs to rake up all these various combinations, tabulate them, usually in chronological order, append with great minutiae the various promoters of these names, and when finished the result is so largely personal it resembles the society notes in a daily newspaper. We present in an appendix (page 77) an alphabetical list of the names which in our opinion have no value, to the number of 192, and in our Index (page 96) the names we have adopted, to the number of 107. Every writer should, of course, use a nomenclature that expresses his views of how the various species are most naturally grouped into genera. And, where changes are advisable in an author's arrangement, it is at best unfortunate, if he is using a system of writing his own name after such changes, as it may give the impression that this is perhaps the strongest reason for the change. We have made but very few changes and have found it necessary to discover but one new genus.

THE STATE OF PHALLOID KNOWLEDGE.

The phalloids of Europe (and there are but six species in Europe) are, with perhaps one exception, well known. The same can be said to-day of those of the United States, though, owing to the vague manner in which several of them were exploited, it is only in recent years that any clear, definite idea has been obtained of them. Taking into account those that occur in both countries, this includes fourteen species and forms. The first foreign paper in which the phalloids were well presented was only ten years ago, an account of the species of Java, by Penzig. In this paper sixteen species and forms were considered, and at least fourteen were well illustrated. Then there appeared a paper on the phalloids of Brazil, by Moeller, in which nine species were well illustrated. Recent writers, and this includes both Penzig and Moeller, have had the benefit of photography, the best method of illustrating a phalloid. Previously the illustrations were mostly made up from dried specimens or copied from sketches, which gives results, sometimes very good, but often more or less doubtful, sometimes very vague and amusing, and in a few instances they seem to be pure fakes.²

There have been a number of compilations similar to this pamphlet, in which the literature has been raked over, and the supposed species arranged with their names more or less shuffled around. This, however, is the first in which all the pictures have been brought together. The first crude attempt was by Ventenat, in presenting one of the first foreign species. Then Fries'

² We know two or three in Europe that in our belief come under this head, and one in America.

Systema (1823) at which time nine foreign species had been figured, and four in Europe, or thirteen in all, and one of the European was a freak. Excluding the freak, all twelve of these species are recognized in this pamphlet, nine of them under the same names as used by Fries. The next general compilation was by Schlechtendal, about fifty years ago. In the meantime, the new species hunters had been quite busy, and Schlechtendal succeeded in finding forty-five species, and he seems to have taken practically all of them at their face value, nor did he indulge in inventing new genera in order to change the names. It is an evidence of the "progress" that since that time, nearly fifty years ago, not an iota of information has been added nor another specimen recorded as to twenty-seven of his forty-five species. Some of them have been discarded as being worthless on their face, but those of the twenty-seven that are retained and known to-day are included in this pamphlet on exactly the same knowledge (?) that Schlechtendal had when he wrote fifty years ago. The next work was by Professor Fischer, in 1886, a compilation of the species described and numbering seventy-six, including the doubtful ones. Practically the same species were included in Saccardo (vol. 7) two years later. After making these compilations, Professor Fischer began his real study of the subject. First, he visited Paris and wrote his first *Untersuchungen* in 1890, then he visited London and Berlin and wrote his second *Untersuchungen* in 1893. A third *Untersuchungen*, principally to include the work of Penzig and Moeller, was issued in 1900. Professor Fischer has studied practically all the specimens in the museums of Europe and the result of his studies has been the rejection of many of the species included in his earliest work, and the reduction of others to forms or varieties. Of the seventy-six species included in his first work, only twenty-three stand as original and good species, and twenty-eight are doubtful. In addition, twenty-eight new species have been added, mostly the work of himself, Hennings, Penzig, and Moeller. This makes a total of fifty-one species, recognized as "good" by Professor Fischer, and twenty-eight doubtful, or a total of seventy-nine.

I have worked over practically the same ground as Professor Fischer, the same museums, and I am in very close accord with him as to the species. As are all who have had the opportunity to see specimens from many localities, Professor Fischer is very liberal in the treatment of species; more so than I, for I maintain a number in this pamphlet that Professor Fischer refers to synonymy. I have not refused to recognize any "new species" that has been exploited in an intelligent manner and that was accompanied by a drawing or photograph showing any material difference. The twenty odd phalloids in this pamphlet, in addition to those recognized in Professor Fischer's latest work, are mostly those that he has referred as forms.

I decline to recognize the alleged "new species" that have been proposed with so much verbosity and so little illustration. No man can give any idea of a phalloid by a mere word description, whether he writes in English, French, German, Chinese, or Pidgin Latin, and it is time this fiction was wiped out of our "literature." In these days of "law-makers" there ought to be a law with a heavy prison penalty for any one who engages in such work. I refer to them in the synonyms as "nomina nuda," although it is a paradox to so call things exploited with so much verbosity.

THE WORK IN THIS PAMPHLET.

We have included in this pamphlet the best illustration known of each phalloid that we recognize. We consider the study of phalloids largely a picture study, and our readers can take these illustrations and form an opinion as to the identity of any phalloid they find with almost as much advantage as if they had access to the types.

In our text we have not entered into minute descriptions, believing that in most cases it is superfluous. We have given the leading facts as to the occurrence of the various species as far as known, the color, and have pointed out the manner in which they differ from each other. We have presented

the best picture possible of each species, and in many cases the copies of the original illustration from which the description was drawn. With these facts before him, the reader can learn just as much about the phalloid as the author who named it and wrote the "description."

CLASSIFICATION.

There are relatively few genera of phalloids, and they are classed by their general form, so that the classification is a very simple matter and will be readily understood by the following table and the illustrations. As a matter of convenience we divide them into five groups:

1st, The simple stem section. Gleba borne directly on the upper portion of a simple stem, or on a pileus borne on top of a simple stem.

Gleba borne on the outer surface of a special pileus.

- Pileus even, rugose, or reticulate.....**Phallus**
- Pileus surface strongly convolute**Clautriavia**
- Pileus of a lamellate structure, the gleba covering the plates.... **Itajahya**
- Gleba borne directly on the upper portion of the stem. No special pileus.
- Smooth, even**Mutinus**
- Rugose, papillate or uneven**Jansia**
- Gleba covering a rudimentary network**Floccomutinus**

2d, The lobed section. Gleba spread over or on the inner surface of free arms or lobes at the apex of the stem.

- Arms free at the apex of a columnar stem**Lysurus**
- Stipe, a flaring tube, the limb lobed**Anthurus**
- Stipe bearing a disk-like expansion, the limb divided into lobes or segments**Aseroe**

3d, The columnar section. Receptacle consisting of simple, vertical columns, united at the top and bearing the gleba on the inner sides.

- Sessile**Laterna**
- Stalked**Pseudocolus**

4th, The clathrate section. Receptacle in the form of a clathrate or latticed structure.

- Sessile, simple**Ciathrus**
- Stalked, receptacle a simple net.

- Borne on a simple stem.....**Simblum**
- Borne on columns that are united into a hollow tube at the base....**Colus**
- Stalked, the net-work having knob-like projections**Kalchbrennera**

5th, Anomalous genus, without volva.....**Phallogaster**

THE GENUS PHALLUS.

This is the original genus of Europe and from whence the name of the order is derived. The genus is very simply characterized by having a pileus, borne on the top of a simple stem. All species of the genus are very much alike as to shape, but differ in color, in size, in smoothness or roughness of the pileus, and in various developments of a veil. This veil, which is only known as rudimentary in the related genus *Mutinus*, varies much in different species of *Phallus*, and even in the same species in degrees of development. Some species have only a rudimentary veil, others a distinct but very

short veil hidden under the pileus, or slightly protruding, others a very conspicuous, long veil. The gleba covers the outer surface of the pileus. In a few species this pileus is even, or relatively smooth; others reticulate, or ridged. Usually the pileus has an apical collar that is entire or perforate, sometimes in the same species. Some species are devoid of this apical collar, and one, *Phallus subtilis*, has been erected into a genus principally on this account. We would divide the species into two sections, as Professor Fischer does, though we would not designate these sections by distinct generic names. We think the old name *Phallus* should cover them both.

Section 1, Veil short or merely rudimentary. Section 2, With distinct veils. Each section is also subdivided on the character, whether the pileus is relatively smooth and even, or is reticulate with ridges.

SECTION I. VEIL SHORT OR MERELY RUDIMENTARY.

PILEUS STRONGLY RETICULATE.

PHALLUS IMPUDICUS (Fig. 1).—It seems to me to be useless to use any space in describing *Phallus impudicus*. It is such a well-known plant, even to every peasant in Europe, and, besides, our photograph is the best description. The stem is white and the pileus has strong reticulations, not shown in our photograph where they are covered with the gleba. *Phallus impudicus* is the original phalloid, and the most common one of Europe. It extends throughout Europe. In the United States we do not have the type form of Europe, but a pinkish variety known as *Phallus imperialis*. In Japan, *Phallus impudicus* (the type form I judge from the drawings I have seen) is common. In Australia it is rare, if it occurs at all. Only one collection is known, now at Kew, which does not accord exactly with the European plant, but is close to it. *Phallus impudicus* probably occurs in other countries, but the above are all that are surely known.

Forms.

PHALLUS IMPERIALIS.—This form differs from the type form only in having a pink volva and in its distribution. I am told that in France it has a different habitat, and a different odor. I can not vouch for that. At any rate it is a rare plant in Europe, widely distributed but infrequent. In the United States it is the only form of *Phallus impudicus* we have. It is common in the West—Colorado, Southern California, and Texas. East of the Mississippi, I know of but one station, Washington, D. C. From its distribution it is evidently a plant that favors a warm climate and a sandy soil.

PHALLUS COSTATUS (Fig. 2).—This species, which was described from Java, is evidently similar to *Phallus impudicus*, and seems to me is better considered as a form. It differs chiefly in having more pronounced, almost winged reticulations to the pileus, and the substance of the pileus is described as yellowish-white.

PHALLUS TENUIS (Fig. 3).—A small yellow-species, native of the Orient. It can easily be known from all others of the section by its yellow color, both of stipe and pileus, and in addition by its



Fig. 1.
PHALLUS IMPUDICUS.



Fig. 7.
PHALLUS RAVENELII.



Fig. 3.
PHALLUS
TENUIS.



Fig. 5.
PHALLUS RUBICUNDUS.



Fig. 9.
PHALLUS RUGULOSUS.



Fig. 2.

PHALLUS COSTATUS.



Fig. 4.

PHALLUS FAVOSUS.



Fig. 8.

PHALLUS RAVENELII.
(reduced)
(With protruding veil.)



Fig. 10.

PHALLUS GLUTINOLENS.

small size and thin substance. The dried specimens appear like a thin skin. *Phallus tenuis* was originally from Java, but must be a rare species there, as Dr. Bernard does not record it. It occurs also in Ceylon (specimens at Kew), and Professor Kusano has found it (very rarely) in Japan. In the latter country it grew on rotten wood. The stipe of the Javanese form is yellow, but in Japan it was represented as white. The original description makes no mention of the plant having a veil, but one of Penzig's figures shows a rudimentary veil hidden under the pileus.

PHALLUS FAVOSUS (Fig. 4).—This species, also known from Java, and rare there, is intermediate between *Phallus impudicus* and *Phallus tenuis*. With the large size of the former, it has a relatively thin pileus and a pale, yellowish stem. The substance of the pileus is also pale, but not so clear yellow as that of *tenuis*. It is only known from the original record.

PILEUS RELATIVELY SMOOTH OR MERELY RUGULOSE.

PHALLUS RUBICUNDUS (Fig. 5).—Stem, red. Pileus, red, smooth, or slightly rugose, covered with the greenish gleba. Apex, perforate, or sometimes imperforate. This is the only red species of the genus *Phallus* that we have, and it is widely distributed. It occurs in abundance in certain localities in our Southern States and many other warm countries. It has been named from India (*Phallus aurantiacus*), Africa (*Phallus sanguineus*), Australia, Hawaii. I have seen a drawing from China, and it is reported from Japan. In Hawaii it has been shown to be the cause of a destructive root disease of the sugar cane. When we get a better knowledge of the distribution of our phalloids, I think that *Phallus rubicundus* will be found in almost all sugar countries. I believe there is only one red *Phallus*. Forms from various countries seem to differ in being slender or obese; the pilei, in being truncate or acute, perforate or imperforate, with an apical collar or without, but the material is not at hand from which to form any opinion as to the systematic value (if any) of these differences.

Forms.

PHALLUS GRACILIS (Fig. 6).—*Phallus rubicundus* varies chiefly in stature. Slender forms have been called *Phallus gracilis*. For a long time the characters of *Phallus rubicundus* were not known other than the fact that we had a red *Phallus* in our Southern States. A recent article of Professor Long has given us a clear idea of its characters and convinced us there is no distinction between it and *Phallus aurantiacus* as it has generally been known in foreign countries.

PHALLUS RAVENELII (Fig. 7).—This is the most common phalloid of the United States, there replacing *Phallus impudicus* of Europe. In general appearance it resembles *Phallus impudicus*, but has a smoother pileus and a veil, usually short and hidden under the pileus. Rarely, however, it occurs with a protruding veil (Fig. 8). Usually *Phallus Ravenelii* grows on logs in the woods, sometimes on




Fig. 6

PHALLUS GRACILIS.



Fig. 12.
PHALLUS INDUSIATUS.



Fig. 15.
PHALLUS DAEMONUM.



Fig. 16.
PHALLUS DUPLICATUS.

the ground, and sometimes it develops in the greatest abundance on old piles of sawdust. The species is only known from the United States and Canada.

PHALLUS RUGULOSUS (Fig. 9).—Pileus, thimble-shaped, almost even or slightly rugulose, with a small, globose, apiculate collar. Color, dark. Veil, none. Stem, pale reddish. This species was described from alcoholic material, and is known only from Japan, where it is reported to be common. I have seen a drawing from Professor Kusano. As I understand it, the substance of the pileus is not red, otherwise the plant seems close to *Phallus rubicundus*. I should not be at all surprised if it develops that it is a slender form (*gracilis*) when the color of the pileus substance is known.

PHALLUS GLUTINOLENS (Fig. 10).—This is a unique species of *Phallus*, known only from Brazil. It has white stipe and no evident veil. The pileus is smooth and differs from all other species in the *globose* shape. It has only been observed by its original author, who gives us a good photograph of it.

PHALLUS SUBTILIS (Fig. 11).—This Brazilian species has only been illustrated by a sectional drawing. A photograph would not show any marked difference from any other small *Phallus*. It was erected into a separate genus because the pileus has no apical collar, and a section shows it to be formed of radiate plates. It is also somewhat gelatinous in its nature. It is only known from Brazil and from the work of the original author.

SECTION 2. VEIL EVIDENT, USUALLY STRONGLY DEVELOPED.

PILEUS STRONGLY RETICULATE.

PHALLUS INDUSIATUS (Fig. 12).—Pileus broadly campanulate, strongly reticulate. Veil strongly developed, of small, slender threads and large meshes. Color of stipe and veil white. This is the most common phalloid of all tropical countries and is found in quantities in all of the museums. We have noted specimens from Australia, India, Andaman Island, Java, Ceylon, East Africa, Mauritius, Mexico, Brazil, British Guiana, French Guiana, South Africa, Surinam, New Caledonia, Cuba, Tonkin, Philippines, Borneo, Jamaica. We have received it from a number of correspondents and have collected it (common) in Samoa.

Forms.

Phallus indusiatus varies in the tropics, chiefly in the shape of the pileus and the veils. Also in color, I think, and I suspect that in time it will not be found practicable to keep distinct *Phallus callichrous* and *Phallus multicolor* as other than color forms. Usually the veil is flaccid, but at other times more rigid. Sometimes it is united above into a distinct membrane. These forms seem to have a geographical significance, but so little is known that at present it is not possible to designate the distribution of the various forms. In Samoa, where I have observed it common, it never takes anything but the type form.




Fig. 13.
PHALLUS MOELLERI.

Digitized by Google

PHALLUS ROSEUS.—A form with a pink veil, which is at Paris, from French Guiana. It is also reported from Java.

PHALLUS MOELLERI (Fig. 13).—A form with a narrow pileus and rigid, spreading veil, as illustrated by Alfred Moeller, from Brazil. Professor Moeller states that in Brazil it runs into the type form so intimately that it is not practicable to keep it distinct.

In the recent article by T. Petch, it is stated that this rigid veil is not a form even, but the *normal* condition of the veil of *Phallus indusiatus* when first expanded and before the sun strikes it. That which I have taken for the type form is a condition after the veil has been wilted by the sun. I have never observed this (in Samoa) nor should I have suspected it, as they seem so different, but Professor Petch undoubtedly knows. In the interest of truth then "*Phallus Moelleri*" must be deleted, even as a form.

PHALLUS ROCHESTERENSIS (Fig. 18).—A form with an elongated thimble-shaped pileus and narrow, cylindrical veil is found at Kew, from Australia. It has been illustrated under the erroneous name, *Phallus merulinus*.

Color Forms.

There are two very showy tropical phalloids that in shape and other characters appear to be the same as *Phallus indusiatus*, but have bright colors. At the present time we can characterize them by their colors, but when the phalloids come to be well known, I think so many intermediate colors will be found that color characters alone will not be held to constitute species. T. Petch finds these color forms abundant in Ceylon, and states that they grade into the white form so intimately that it is not possible to keep them distinct even as forms. I am satisfied, however, that they have a geographical significance. They do not occur in Samoa, and Mr. C. B. Ussher, who has observed the species in tropical Africa, informs me that they are absent there.

PHALLUS MULTICOLOR (Fig. 14).—This was originally from Australia, but has been recently found and photographed from Java. Pileus orange red, veil bright lemon yellow, stipe lemon yellow, volva pink, mycelium purple. The characters, if they are real characters, of the species are the colors as stated above.

PHALLUS CALLICHOUS.—This appears to be different from multicolor only in the coloration. The pileus is orange, the veil and stipe white. It has never been illustrated, but probably could not be distinguished by a photograph alone from either multicolor or *indusiatus*. It was originally named from Brazil, but similarly colored plants have been observed in Java, Africa, and Australia.

PHALLUS DAEMONUM (Fig. 15).—This, which was the original foreign phalloid, illustrated from the island of Amboy, was published one hundred and sixty years ago.³ All that is known of it to this day is the original, crude figure that we present. It seems quite distinct from the usual form in its punctate rather than reticulate pileus, if that proves to be a character of the plant and not of the figure only.

PHALLUS DUPLICATUS (Fig. 16).—Pileus with a strongly developed apical collar and strong reticulations. Veil long, white, of

³ It has therefore strong claims to be taken as the specific name for the species as proposed by Professor McGinty. There are two objections to it, however. First, it may be the "type" in the perverted sense that the word "type" is usually used, but it does not seem to be the typical form as the plant usually occurs. Second, it is not advisable to use so familiarly the name of His Satanic Majesty.



Fig. 14.
PHALLUS MULTICOLOR.

thick threads, which in alcoholic specimens contract and form almost a membrane. This is a common plant in the United States and is so close to the tropical species it may well be considered a temperate region form of it. However, it differs in the nature of the veil and the usual shape of the pileus, and I am convinced that it is as distinct as species generally are. The veil (which is torn in our figure) is a conspicuous feature of the plant.

Forms.

We would be disposed to consider related plants with a similar veil as forms of this species.

PHALLUS SUBUCULATUS, of Algeria, which was inaccurately figured, is, we think, a form of it.

PHALLUS MAURITIANUS (Fig. 17).—This form, which we have received in alcohol from Chas. O'Connor, of Mauritius, we feel is worthy of a separate designation as a form. It differs from the typical plant in the nature of the reticulations of the pileus, and is better shown in our photograph than we can tell it.

Note.—We formerly included in this section, under the name *Phallus irpicinus*, the only known phalloid with a well developed veil and rugulose pileus. It was proposed as a new genus (or a new section) *Clautriavia*, on account of having the pileus minutely convolute. We were not disposed to consider that of generic value, until recently when we saw at Berlin a New Guinea species with such a *strongly convolute* pileus, and such a marked character that we now feel that the genus *Clautriavia* should be maintained. Compare *Clautriavia merulina* on the next page.

IMPERFECTLY KNOWN SPECIES OF THE GENUS PHALLUS.

Many phalloids are known (?) only from old cuts based mostly on dried specimens and, in some instances, fertile imaginations. Naturally they are of not much importance for no one ever finds them again, but there is no way of getting rid of them. The genus *Phallus* has been especially favored (?) in this regard. We give a short synopsis of them here and have relegated the (alleged) pictures to an appendix.

PHALLUS DISCOLOR (Fig. 95).—From Australia, if correctly illustrated (with an emphasis on the "if"), is an intermediate plant connecting the genera *Phallus* and *Mutinus*. It was alleged to have the pileus adnate at the base to the *top* of the stem.

PHALLUS CALYPTRATUS (Fig. 96).—From Australia. Appears to be based chiefly on an accidental mass of gleba dried on top of the pileus.

PHALLUS RETUSUS.—Originally exploited as a new genus, it is reported by Professor Fischer (who has seen the "type") as an obese form of *aurantiacus*. The figure has no resemblance to *aurantiacus*, but it does not follow that the plant has none. It was from Australia.

PHALLUS CAMPANULATUS (Fig. 98).—Known only from the figure (Uruguay). The little cup at the base is not the volva, but the "inner" volva. It seems to have an even pileus and be close to *Ravenelii*, though nothing is known as to its veil. No specimen exists.

PHALLUS CELEBICUS (Fig. 99).—Said to grow in the Celebes and to have a whitish pileus and a yellow stem. It appears from the published account to be very close to *Phallus rubicundus*.

PHALLUS CANARIENSIS (Fig. 100).—If the figure is correct it is a peculiar little species with a slender stipe and large, rugulose pileus. Both




Fig. 18.

PHALLUS ROCHESTERENSIS.



Fig. 17.

PHALLUS MAURITIANUS.



Fig. 11.

PHALLUS SUBTILIS.
(Section.)


pileus and stipe are rose colored. It was from the Canary Islands. I have found no type.

PHALLUS FARLOWII.—This from alcoholic (or dried?) material from Brazil has never been illustrated. It is said to have a membranous veil, otherwise it is very close to *Phallus indusiatus*.

PHALLUS QUADRICOLOR (Fig. 102).—I think is probably based on a specimen of *Phallus multicolor* which has lost its veil. From Australia.

THE GENUS CLAUTRIAVIA.

This genus is characterized by having the surface of the pileus convoluted in folds, the gleba covering the folds and permeating the interspaces between them. Our figure (which is an enlargement six diameters) will give a clear idea of this structure. The original species, *Clautriavia merulina*, which is a frequent plant in Java, Ceylon, and the East Indies, in general (probably) has very minute folds, so that the surface to the eye appears even, but a recently discovered species of New Guinea, *Clautriavia Lauterbachii*, has the folds so strongly convoluted that in the egg the pileus appears to be a crumpled veil covered with gleba.



CLAUTRIAVIA MERULINA (Fig. 19).—This species has the general appearance of being a *Phallus*.

The pileus, however, instead of being a plain or reticulate membrane with the gleba on the outer surface, consists of minutely convoluted folds, the gleba permeating the depressions between the folds. It has long been known as a common species in Java.⁴ Recently T. Petch has published that it is abundant in the grounds of the Botanical Garden at Peradeniya, Ceylon. When the truth of the subject is known it will probably be found to be generally distributed in the East Indies and neighboring countries.

CLAUTRIAVIA LAUTERBACHII (Fig. 20).—This species, which has a most remarkable structure, is unfortunately known only from some undeveloped plants from New Guinea. The pileus in the

⁴ Berkeley named the plant *Phallus merulinus*, many years ago, and while he gave no formal description of it (in pidgin Latin) he characterized it in an unmistakable manner, it appears to me now. Fischer incorrectly referred the name as a synonym for *Phallus indusiatus*, and Cooke illustrated a form of *Phallus indusiatus* of Australia under Berkeley's name. Patouillard discovered it to be a "new species" from Java, and named it *irpincinus*, which name we have previously used, and would continue to use if it had any application to the plant. We adopt Berkeley's name, not on the grounds of "priority," but suitability, believing that when a plant has two names, one very good and one very bad, the better should be chosen.



Fig. 19.

CLAUTRIAVIA MERULINA.

egg is a strongly folded and convolute membrane resembling at first view a crumpled veil. What form it takes in the developed plant is not known, but it is probable that it does not change much, as the form of a pileus is in all known instances well defined in the egg. In addition the volva is covered with wart-like processes, which, while unknown as to any other phalloid, is in my opinion a minor character. The plant is only known from New Guinea, and a photograph of a developed plant is much desired.⁵




Fig. 20.

CLAUTRIAVIA LAUTERBACHII.

Section by Fischer. Photograph of the volva, also of the folds of the inner face of the pileus.

THE GENUS ITAJAHYA.

This genus in general appearance resembles the genus Phallus, but is quite different in the structure of the pileus. This consists of lamellate plates, the gleba covering these plates, permeating the inner structure of the pileus.

⁵ Until I saw the specimens I had a very erroneous idea of the characters of the plant, and I think they have been inaccurately presented in the published accounts. When Dr. Hennings received these phalloid eggs he sent them to Professor Fischer, who made what impresses me as a very accurate drawing of a section that he returned to Dr. Hennings with the suggestion that it be called *Ithyphallus lauterbachii*. We reproduce Professor Fischer's section in our figure (20). Dr. Hennings did not publish Fischer's figure as received, but modified it, showing a "hut" and an "indusium." The plant has but one membrane, which should be called the "hut," as it bears the gleba and is analogous to a pileus. There is no indusium. One of the egg sections at Berlin would at first view seem to have a rudimentary indusium, but on closely examining it I find it is a division of the stem, which in this instance seems to divide above and support the pileus in the manner of a *Helvella*. The pileus in the egg is so convoluted that my first impression (until I noticed that it bore the gleba) was that it was an indusium, and that here we had a type of a new genus of phalloid which had a veil but no pileus. Dr. Hennings first published it under Fischer's name, *Ithyphallus lauterbachii*. Afterwards he republished it as a new genus *Echinophallus*, basing it principally on the protuberances of the volva, a minor character, in my opinion. The main character of the plant, the strongly folded and convoluted pileus, is unique in this species and establishes, for me at least, the validity of Patouillard's genus *Clautriavia*, based on the same character, though in a much less developed form.

ITAJAHYA GALERICULATA (Figs. 21 and 22).—But one species of the genus is known, which is a native of Brazil, from whence it was well described and illustrated by Moeller. It has since been found there by Father Schupp. Robert E. Fries recently records the plant as common in Argentina, and it is probably frequent and widely distributed in South America.⁶ Our photographs and the sectional figure of the pileus are all that are necessary to enable one to recognize the plant.



Fig. 21.

ITAJAHYA GALERICULATA.



Fig. 22.

ITAJAHYA GALERICULATA. (Section.)

THE GENUS MUTINUS.

This genus is distinguished from Phallus, to which it was formerly united by having no distinct pileus, the gleba being borne on the upper portion of a simple stem. Sometimes the gleba-bearing por-

⁶ Mr. Fries suggests, not without reason, that it may be the original of Spegazzini's "new genus" *Albofiella*, which if true is a prior name. In that case I submit, would it not be a rank injustice and a travesty on science to replace the excellent work done by Moeller, or his name, by the inaccurate work of Spegazzini?

tion is distinct from the stem, taking somewhat the nature of a distinct pileus, but in other species it is not clearly marked from the stem. The species of *Mutinus* are all very similar and are distinguished by their general form. All are red, or sometimes have white forms.

MUTINUS CANINUS (Fig. 23).—This, which is the only species of *Mutinus* that grows in Europe, has a short, distinct, spore-bearing portion, which is sharply distinct from the stem. I do not know whether it is a constant character, but I have seen alcoholic specimens where the receptacle was abruptly contracted and of a smaller diameter than the stem. The structure of the receptacle is always different, being of small, thick-walled cells, while those of the stem are large and thin-walled.⁷ *Mutinus caninus* is not rare and is widely spread in Europe. In the United States it is much rarer, and while I think it is well authenticated, it occurs principally in the Eastern States. The stem of *Mutinus caninus* is usually red, though white-stemmed forms have been figured on several occasions.

MUTINUS ELEGANS (Fig. 24).—In this species there is no distinction between the stem and the spore-bearing portion. It is all one uniform, cellular structure, with no sharp line of demarcation. The form is generally tapering from a thickened base to an acute apex. *Mutinus elegans* is the most common *Mutinus* that we have in the United States. It grows in the woods around old logs or soil rich in humus. It is not rare. The color is red or orange.

MUTINUS RAVENELII (Fig. 25).—This species has the same cellular structure as the preceding and has been held to be the same plant. I am satisfied it is distinct in form (usual) and habitat. The shape is club-form, thickened above, and tapering below. The habitat is old fields devoid of woods humus. It is a rare plant in the United States. The color is red.

MUTINUS BAMBUSINUS (Fig. 26).—Receptacle distinct from the stipe, formed of small cells. Color of both stipe and receptacle is red. This, which seems to be the common species of the tropics, is very similar to *Mutinus caninus* of Europe. However, it has a much longer spore-bearing portion and the color is brighter red. It was originally from Java, but occurs in the Celebes, Brazil, and no doubt in many tropical countries. It has been noted, adventitious, in the hot-houses at Kew.

MUTINUS FLEISCHERI (Fig. 27).—The most obese species of *Mutinus* known. It has a thick stem and a very short, contracted spore-bearing portion. Its structure is that of *Mutinus caninus*, of Europe, but it is a much more obese plant. It is known only from Java and is a rare plant there. The color is red.

⁷ This has always been my observation, and my understanding of the essential character of *Mutinus caninus*. I have recently seen at Berlin alcoholic specimens of eggs and sections of eggs of *Mutinus caninus* from Europe, where I can not note any difference in the cells of the stem and gleba-bearing portion.



Fig. 23.

MUTINUS CANINUS.



Fig. 24.

MUTINUS ELEGANS.



Fig. 26.

MUTINUS BAMBUSINUS.

MUTINUS PENTAGONUS (Fig. 28).—All the previous species of *Mutinus* have cylindrical stems, but in this species the stem is pentagonal (or sometimes six-angled). The gleba-bearing portion is also strongly fluted, and the gleba is borne on the channels with free edges. In the genus *Lysurus* the lobes, when young, are connivent, and the young plants of *Lysurus Mokusin* evidently closely resemble this species. In *Mutinus pentagonus* I am convinced from an examination of dried specimens that there are no arms, but that the receptacle consists of a single piece. *Mutinus pentagonus* is known only from Australia, and but scantily there.

MUTINUS XYLOGENUS (Fig. 29).—This is the smallest phalloid known and an idea of its size can be obtained from our photograph, which is an enlargement six diameters. It is only known from a collection made in French Guiana many years ago, and preserved at Paris. It is a question whether it is a *Phallus* or a *Mutinus* (cfr. *Myc. Notes*, p. 336). If a *Mutinus*, it is not only the smallest species known, but differs from all other species in having a globose mass of gleba.



Fig. 25.

MUTINUS RAVENELII.



Fig. 29.

MUTINUS XYLOGENUS.
(Enlarged x6.)

DOUBTFUL AND LITTLE KNOWN SPECIES.

The same remarks apply here as under the same head concerning the genus *Phallus*. *Mutinus minimus*, *Mutinus borneensis*, *Mutinus proximus*, and *Jansia boninensis* may all prove to be the same plant.



Fig. 27.

MUTINUS FLEISCHERI.




Fig. 28.

MUTINUS PENTAGONUS.

MUTINUS MINIMUS (Fig. 103).—Known from a figure reconstructed from a dried or alcoholic specimen. Color, red. Seems to differ from others in its rugulose receptacle. Described from Tonkin.

MUTINUS BORNEENSIS (Fig. 104).—Figured from Borneo, in an Italian journal. Was said to have a white stipe and short red spore-bearing portion. We reproduce a figure from Tonkin.⁸

MUTINUS PROXIMUS.—Based on a dried specimen in the British Museum, from Ceylon. It is a small species, described as having a white stipe, but the plant is accompanied by a sketch showing an orange stipe. It seems to be close to *caninus*. It has not been figured.

MUTINUS CURTUS (Fig. 105).—Only known from one collection made in Australia sixty years ago, which seems to be immature. The figure reconstructed by Corda is no doubt inaccurate, especially as to the lobed volva.

MUTINUS PAPUASIUS (Fig. 106).—Known only from a figure from a dried specimen, from Australia. It is not known whether it is a *Mutinus* or a *Phallus*.

MUTINUS ARGENTINUS (Fig. 107).—This was originally published without illustration and was referred by Professor Fischer, doubtfully, to *Mutinus muelleri*. The latter seems from Fischer's illustration to be *Mutinus bambusinus*, and is so referred by Moeller. Spegazzini has recently published a figure of *Mutinus argentinus* which seems to me quite different from *bambusinus*. It has a short, thick spore-bearing portion. From the figure one could not say it was not *Mutinus caninus* of Europe, though it is rather stocky for that.



Fig. 30.

JANSIA RUGOSA.
(Natural size.)



Fig. 32.

JANSIA ELEGANS.
(Natural size.)



Fig. 35.

JANSIA BONINENSIS.

⁸ In our account, *Myc. Notes*, p. 388, we confused *Mutinus borneensis* of Borneo with *Mutinus boninensis* of Bonin Island. Both are imperfectly known, but the latter seems to be a *Jansia*, and both may in time prove to be the same plant. Since we have seen the types of *Jansia boninensis* we think we have inaccurately referred here (page 402) a species of *Mutinus* from Japan.

THE GENUS JANSIA.

This is a genus of very small phalloids, common in Java and well illustrated by Penzig. The general form is that of a little *Mastigotrochus*, but the spore-bearing portion is strongly differentiated from the stipe, and it is strongly rugulose or papillate. Two species occur in Java and have been well illustrated. They grow on rotten wood. There are two imperfectly known species, one from Bonin Island and one from Australia.



Fig. 31.




Fig. 33.




Fig. 34.

JANSIA RUGOSA.
(Enlarged.)

JANSIA ELEGANS.
(Enlarged.)

JANSIA ANNULATA

JANSIA RUGOSA (Figs. 30 and 31).—This is a very small phalloid, which is common in Java. The short gleba-bearing portion is strongly distinct from the stipe and is strongly rugulose, as shown in our enlargement (Fig. 31). It is the only species of *Jansia* that is common and well known. This little plant is white and grows on rotten wood.

JANSIA ELEGANS (Figs. 32 and 33).—This species is also known only from Java and is rare, at least Dr. Bernard does not report it. It grows on rotten bamboo stems. It is of the same size

nd
u-
om
ur
od.
nd

s the preceding little species, but the gleba-bearing
y covered with little processes, instead of being ru-

LATA (Fig. 34).—This plant is known only from a figure
ralia. No specimen exists. The stipe is white, the gleba-
ochre" and "annulated." The plant is therefore probably
not surely known.

NENSIS (Fig. 35).—This species from Bonin Island, is
with certainty, from one collection in alcohol. The gleba
ightly rugulose, and it seems intermediate between Mutinus
pically neither. The type collection is in alcohol in Berlin.
Mutinus minimus and Mutinus borneensis are not both
cies.



Fig. 36a.

MUTINUS ZENKERI

(g habits of plant.)



Fig. 36.

FLOCCOMUTINUS
ZENKERI.
(Much enlarged.)

GENUS FLOCCOMUTINUS.

very curious and is intermediate between the pileate
halloids. The gleba covers a *loosely attached* net-
the stipe, and while *similar* to the veil of a Phallus,
the *pileus* of a Phallus. The drawing by Professor
e 36) gives a good idea of this structure. The exact

all
on
vn
is
on

so
e-
ze

attachment of this network I could not make out from the type owing to the minuteness of the parts, though that it is attached (loosely) in some manner is evident. It appears to me as a very distinct genus, essentially different in its basic structure from both the genera *Jansia* and *Mutinus*, with which it has been recently united.

FLOCCOMUTINUS ZENKERI (Figs. 36 and 36a).—But one collection of this curious genus is known, which is in alcohol in the museum at Berlin. It is accompanied by a colored sketch of the fresh plant, made by the collector, showing well its habits. We reproduce this drawing (Fig. 36a), though, owing to the difficulty of photographing colors, our figure does not do the drawing justice. In habits *Floccomutinus Zenkeri* is very similar to *Jansia elegans* of Java. The little plants are borne caespitose on a common, mycelial pad. The eggs are elongated in form and open at the apex. The volva is not accurately shown in Figure 36.

THE GENUS *LYSURUS*.

This genus has been very much misunderstood, though of a very simple structure. It consists of free arms borne on a hollow columnar stem. The gleba is borne on the arms. It has been shown that in the original species the gleba is borne on the outer side of the arms, hence species with gleba on the inner surface of the arms have been transferred to *Anthurus*, which genus does not have a columnar stem. I think it is much simpler to define *Lysurus* as originally defined, viz.: a columnar stem bearing free arms at the apex. With respect to the position of the gleba, there are evidently two series, and a new genus will probably be made for those with the gleba on the inner side of the arms. It has recently been shown by Mr. T. Petch, Ceylon, that the arms of *Lysurus Gardneri*⁹ (which was the second species known) are not entirely free, but are united by a delicate membrane. We would therefore modify the definition of the genus to include species with arms free or very slightly united.

⁹ Ever since the species was published there has been a difference of opinion as to whether the arms were united or not, a difference of opinion that was legitimate from the fact that the type specimens at Kew do not bear out the original statement in this respect. Before seeing the specimens Fischer decided they were united, and changed the classification on that account. Massee, who had the type in charge, writes: "The segments are not organically united at the tip, but during the young stage are closely pressed together, and having been dried in that condition appear to be united. When the mucilage is moistened the tips are found to be quite free and are normally so in several out of the twenty-three specimens in the herbarium." Knowing the direct divergence of opinion on the subject, I went very carefully into the question on my previous visit to Kew. Some of the specimens appear to have never been united (see photograph, Figure 38a, from one of the types), and while in many specimens they are convergent and covered with the gleba, I did not believe there was any union between them, and so published. I included them in *Lysurus*, where I think the plant is best classed, though as they are united it becomes necessary to modify the definition of this genus. Mr. Petch, who has observed two specimens fresh, finds the tips of the arms united by a delicate membrane, a fact that could not be told from the dried specimens at Kew. He puts it in the genus *Colus*, although it has no resemblance or analogy to that genus. In order to justify his name he changed the definition of the genus *Colus*, and gives it a definition that excludes from the genus the original and only species known to belong to it. I believe a man has a right to modify a definition of a genus to include species which he thinks should be classed in the genus, but he has no right to draw up his definition so as to exclude the original species and change the original idea entirely.

LYSURUS MOKUSIN (Fig. 37).—This is the original species of *Lysurus* and was one of the first foreign phalloids known. It was figured in 1774 by Father Cibot, a missionary in China. The stem is strongly fluted and bears free arms, which are also fluted. It has been found in several stations in China and Japan, but is unknown from other parts of the world. We have a drawing from Professor Gono, Japan, that shows a white stem and red arms. We do not know, however, that these colors are constant.



Fig. 38a

LYSURUS GARDNERI.
(Photo of a type.)



Fig. 39.

LYSURUS AUSTRALIENSIS.
(From the type.)



Fig. 40.

LYSURUS BOREALIS.
(Stocky form.)

LYSURUS GARDNERI (Figs. 38 and 38a).—This species has been known for many years only from the original collection from Ceylon, at Kew. It has been recently discovered in Ceylon by Mr. Petch, but is of rare occurrence and only recorded from the island. Mr. Petch's observations of the fresh plant show that the arms are united by a very distinct membrane, which would take it out of the genus *Lysurus* as formerly defined. As it was originally classed in this genus, however, and as its relations are evidently with the genus *Lysurus*, I think it better to modify the definition of the genus to in-



Fig. 37.

LYSURUS MOKUSIN.



Fig. 38.

LYSURUS GARDNERI.

clude it.¹⁰ The photograph of *Lysurus Gardneri*, as well as the dried specimens, has a close resemblance to the two following species, and I have heretofore believed that in time they would all three prove to be the same species. We must abandon this idea now that Professor Petch has demonstrated that the arms of *Lysurus Gardneri* are organically united, for they are entirely distinct in both of the following species.



Fig. 42.

LYSURUS
CLARAZIANUS.



Fig. 43.

LYSURUS SANCTAE-
CATHERINAE.



Fig. 45.

LYSURUS WOODII.
(From the co-type.)

LYSURUS AUSTRALIENSIS (Fig. 39).—One collection of a *Lysurus* from Australia is at Kew, published as above. How it differs from *Lysurus borealis* I do not know. Professor McAlpine has advised me of a red *Lysurus* in Australia, but I have not had further details. As I think the published figure of *Lysurus Australiensis* is overdrawn and inaccurate, I present a photograph of the type, which, while not satisfactory, is true as far as it goes.

LYSURUS BOREALIS (Figs. 40 and 41).—This is claimed to be distinct from the preceding, but I know no points of difference.

¹⁰ It has been classed in the genus *Colus*, but for me it has no characters in common with the genus *Colus*, which is a *clathrate* genus. It might be included in *Pseudocolus* according to the definition of that genus, but it is so different from all species of that genus I think it better not to so include it.



Fig. 45a.

LYSURUS (unnamed).
 (The limb and an arm enlarged five times.)



Fig. 41.

LYSURUS BOREALIS.
 (Slender form.)




Fig. 44.

LYSURUS CRUCIATUS.

It has a curious history in the United States and Europe and is supposed to be an introduced plant. It grows in gardens, sod, and other cultivated places. It occurs mostly in our Eastern States. In Europe it has been found in three localities, all in recent years. First by Dr. Hennings in Germany, then by Mr. Carleton Rea in England, and then by Mr. Harold Murray, of Manchester, England. Mr. Murray's plant has a white stem and red arms. Professor Long also advised me of a red *Lysurus* in Texas. We present two photographs, one a stocky plant from England, the other a slender specimen from the United States. We are told, however, that these same "stocky" forms occur in the United States.

LYSURUS CLARAZIANUS (Fig. 42).—This was a small plant, described from Argentina. It is red and small, but otherwise seems about the same as the preceding.

LYSURUS SANCTAE-CATHERINAE (Fig. 43).—This was based on a picture from Brazil. It seems to have the gleba in a globose mass on the center of the apex of the stipe rather than surrounding the arms. The color is red. It may be an *Anthurus*.

LYSURUS CRUCIATUS (Fig. 44).—A very small species with four arms, the gleba forming a ball on the top of the stem. It is only known from the original collection, which was from French Guiana, and is preserved at Paris. We present the original drawing in our illustration.

LYSURUS WOODII (Fig. 45).—This is a small, red species, imperfectly known from South Africa. Our photograph is made from the cotype at Kew. The arms are three or four and are "magnificent scarlet," the stem "waxy yellow." The specimens are from Mr. Wood and are the same as those named and figured by Kalchbrenner as *Anthurus Woodii*. While it is unsafe to draw conclusions from dried specimens, we believe the species is a *Lysurus* entirely distinct from the genus *Anthurus* and that Kalchbrenner misconceived and mis-drew the illustration. We, therefore, present a photograph of the dried specimen, which though a very poor illustration is better than an inaccurate drawing.

UNNAMED SPECIES (Fig. 45a).—We have received from F. M. Reader what is surely an unnamed species from Australia. It is a very small species, as will be seen by reference to our photograph, which is an enlargement four diameters. The limb is four-angled, enlarged above, and bears an arm at each angle. The color is red. The specimen sent us (in formalin) had evidently been cut in two pieces and these arms all broken off, so that we could not make much of a picture of it. We think it will be recognized, if found again by our Australian friends, and we do not name it. We hope some one in Australia will give a good photograph of it from the fresh plant and give it a name. We should be glad to have a perfect specimen in alcohol.

THE GENUS ANTHURUS.

Though largely confused with *Lysurus*, the genus *Anthurus* as originally proposed is very distinct. The stem is a *flaring tube*, the limb divided into segments, and it bears the gleba on the inner side of these segments. But one species is satisfactorily known, and that one is due to the work of Prof. D. McAlpine of Australia.



Fig. 46.

ANTHURUS ASEROEFORMIS.




Fig. 49.

ANTHURUS CALATHISCUS.
(The original drawing.)

ANTHURUS ASEROEFORMIS (Fig. 46).—Professor Mc-Alpine describes the plant as follows:

“Receptacle with hollow stem, expanding above into five arms, directed upwards and outwards. Stem salmon pink, slightly darker at top, fully three inches long, rugose with small depressions running more or less in lines and slight ridges running crosswise, so that it looks as if divided into a series of squares, about $\frac{1}{2}$ inch in diameter towards the tapering base and $\frac{7}{8}$ inch at top. Arms three inches long, merging into stem and tapering to a point, blood-red on inner face, convex and broken up into larger or smaller cavities, on outer face there is a continuation of the color of the upper portion of the stem and gradual darkening until toward the tip it is blood-red like inner face with thickened, slightly raised margins and central furrow broken up into small cavities.



Fig. 47.

ANTHURUS MUELLERIANUS.



Fig. 48.

ANTHURUS ARCHERI.

“Gleba blackish with tinge of bronze green, extending along the inner surface of each arm, but not covering the slender tip.

“Volva somewhat cup-shaped, about as long as broad ($1\frac{1}{2}$ inches) dirty-white, splitting at the apex, tapering towards the base and provided there with turfs of elongated fibrous roots.

“Spores hyaline, cylindrical to elongated ellipsoid, rounded at both ends, sometimes vacuolated but generally homogeneous contents, $6-8 \times 2\frac{1}{2}-3$ mic., occasionally 9 mic. long.

“A solitary specimen growing in a garden among violets, near Melbourne, Victoria, April, 1907. Forwarded by C. French, Jr. It had a very disagreeable smell. Owing to its fragile nature, one of the arms fell away and only the arm to the right in the photograph shows the slender tip.”

This description, taken in connection with the photograph that Professor McAlpine sends, gives a perfect idea of the plant, and it is the only *Anthurus* that is really known.

ANTHURUS MUELLERIANUS (Fig. 47).—This, the original species of the genus, is known only from a drawing supposed to be quite inaccurate. It was from Australia, and the color was described as yellowish-red and shown bright red. I rather suspect that it was based on the same plant as the preceding.

ANTHURUS ARCHERI (Fig. 48).—This is known only from a figure, and that is doubtful. It was from Tasmania. It seems from the figure to be an *Anthurus*, but in the sectional drawing the arms are shown to be bifid, and it seems to incline toward the genus *Aseroe*.

ANTHURUS CALATHISCUS (Fig. 49).—The original of this species, as far as I can learn, is a crude figure found in the herbarium of Montagne from Perrottet, India. I think it was published as *Calathiscus Sepia*, and if so, then a most fantastic and imaginary figure was given of it. Perrottet gives the color as "jaune pale." No similar plant has since been sent from India, but his figure evidently is an *Anthurus*.¹¹




Fig. 51.

ASEROE PENTACTINA.



Fig. 53a.

ASEROE PALLIDA.

THE GENUS *ASEROE*.

Stem tubular, abruptly spreading into a horizontal limb, which is divided into a number of long, slender, usually bifid segments.

¹¹ Although I have hunted diligently for the original of the fantastic picture that for sixty years has embellished our phalloid literature, I have found no other evidence than the cut reproduced (Fig. 49.) It has so little resemblance to the published figure that it does not seem possible to have been the source. It was from "Perrottet, India," and on a sheet with two other sketches taken by Perrottet to be different species, but which appear to me to be forms of the same. Montagne has endorsed this sheet "Perrottet, *Calathiscus* et *Aseroe pentactina* Endl.," and it therefore seems to be the source of his "*Calathiscus*."

These are generally prolonged into long, slender points. The color of most species is bright red, and they are among the most showy phalloids. The genus is at home in Australia, where many forms occur. It also grows in Java and the East. No species is known from America or Europe, and it is vaguely known from Africa.¹² The species



Fig. 50.

ASEROE RUBRA.

all are very similar and have been reduced to two by Professor Fischer. However, the figures that are supposed to represent them seem so different that we would prefer to consider them distinct, at least until more is known about them. We believe, however, that there are three distinct species under which the forms should be arranged: *Aseroe rubra*, which includes the Australian forms and has a *narrow* limb; *Aseroe Zeylandica*, to which all the East Indian forms should be referred, and which has a *broad* limb; *Aseroe arachnoidea*, which is quite distinct from both the others.

¹² At Berlin there is a very imperfect dried specimen of *Aseroe* from Africa!! It is so poor that I would not wish to even venture on its form, but the occurrence of the genus in Africa is not recorded, I think, and is of interest.



Fig. 55.

ASEROE ARACHNOIDEA.



Fig. 52.

ASEROE HOOKERI.




Fig. 56.

ASEROE ARACH-
NOIDEA. (Section.)

ASEROE RUBRA (Fig. 50).—This was the original form known, and was from Australia. It has short, spreading rays. This exact form does not appear to have reached Europe since, but adventitious plants which are exactly the same have appeared in the hothouses at Kew.



Fig. 53.

ASEROE MUELLERIANA.

ASEROE PENTACTINA (Fig. 51).—From the specimens that reach Europe this form seems to be the most common form in Australia. It has a narrow limb and long, slender rays. The name, pentactina, referred to the number (five) of the rays of the original specimen, but the number varies and is of no importance.

ASEROE HOOKERI (Fig. 52).—This was a very small form with a short stem and narrow rays that came from New Zealand. It is the smallest form described and appears to me quite different from the others.

ASEROE MUELLERIANA (Fig. 53).—This form from Australia has a broad limb and a general resemblance to *Aseroe Zeylandica* of Ceylon. How-

ever, the rays are shorter and differently disposed. I think it is known only from the picture. It seems quite different in its broad limb from the other Australian forms, if any reliance can be placed on the picture.

ASEROE PALLIDA (Fig. 53a).—At Berlin I found a dried specimen of an Aseroe from New Caledonia sent by Monsieur Le Rat, with a drawing (Fig. 53a) that seems to be well made. It differs from the Australian form not only in its narrow segments but pale coloration. The stem is "pure white," the limb "pale rose." I think it is worthy of record as a marked form of this variable species.



Fig. 54.

ASEROE ZEYLANDICA.

ASEROE ZEYLANDICA (Fig. 54).—This species is originally from Ceylon and is the largest and most showy of the genus. The broad limb is divided into a number of segments, and the whole plant is bright red. It was collected many years ago in Java (and called

Aseroe Junghuhnii), but is very rare there and was not found by Penzig. It has recently been found again by Dr. Bernard, who has kindly sent us the fine photograph which we publish.

ASEROE LYSUROIDES.—This was figured by Corda from specimens from Australia. It has a long, slender stem and short, broad rays. Corda's figures appear to me to represent two different genera, hence I do not reproduce it as I think there is surely something wrong about it.

ASEROE ARACHNOIDEA (Figs. 55 and 56).—This species differs widely from all that precede. It has simple rays, not bifid, as all others. The color is white: all others are red. It was based on alcoholic material at Paris collected "sur fumier" in Cochin China, by Dr. Harmand. It has since been found abundantly in Java by Penzig and Dr. Bernard, though not on manure. The stem is hollow, and pervious at the top, and the arms crown the limb of the stem.




Fig. 59.

LATERNEA TRISCAPA.



Fig. 60.

LATERNEA PUSILLA.

(From the type.)

THE GENUS LATERNEA.

This genus consists of columns (usually two to five) that are united at the top and bear the gleba clinging to the under side. It is chiefly an American genus, being very common in South America and Southern United States. There is one record from Africa and one species known from Japan.

LATERNEA COLUMNATA (Fig. 57 and 58).—Columns from three to five, usually four. When perfectly developed there is a groove on the outer surface. Color red, or perhaps also white. White plants have been figured from Chile and Africa that are probably the same thing. This is the original species of Laternea, and is the most common one. It is abundant in Southern United States and South America, and is also known from the West Indies and Hawaii.

LATERNEA TRISCAPA (Fig. 59).—This was the second species named, and is known only from the original figure. It is very much the same as Laternea columnata except its small size, and it may be only a small form.




Fig. 62.

LATERNEA ANGOLENSIS.



Fig. 61.

LATERNEA
RHACODES.



Fig. 58.

LATERNEA
COLUMNATA.

It came from the West Indies. The figure shows only three columns, and for a long time that was considered its specific character. It is well known, however, that the number of columns varies in other species and undoubtedly also in this.

LATERNEA PUSILLA (Fig. 60).—This is known from a single specimen from Cuba, preserved at Kew. The character of this specimen is the two columns and its exceedingly small size. As *Laternea pusilla* has never been found since and was never figured, we have used for our illustration a photograph of the type specimen. When these small *Laterneas* are known from more ample collections, it will probably not be possible to draw any line between *pusilla* and *triscapa* and perhaps also *columnata*.



Fig. 57.

LATERNEA COLUMNATA.



Fig. 63.

LATERNEA SPEGAZZINI.

LATERNEA RHACODES (Fig. 61).—In this species the inner cells of the columns are torn and lacerated, and on that account has been made into a new genus (*Blumenavia*). As the same character is afforded by more than one *Clathrus*, which are not separated on this account, we feel it better to include this in *Laternea*, with which it otherwise agrees. *Laternea rhacodes* was originally from Brazil, where it is reported to be common. It is not otherwise known.

LATERNEA ANGOLENSIS (Fig. 62).—This, from the picture, which is all that is known about it, is very similar to *columnata* except that the columns are more slender and reduced at the top, and the color is white. It is probably only a white form of *Laternea columnata*. It is the only record of the genus

Laternea in Africa and was from Angola. The recently described *Blumenavia usambarensis* from Africa is probably the same. The type is in alcohol at Berlin.

LATERNEA SPEGAZZINI (Fig. 63).—This, which we know only from a figure, differs from *Laternea columnata* in having the surface covered with papillate projections. It might well be made the type of a new genus. As far as known, it occurs only in Argentina, South America. The illustration shows only three columns, but the number probably varies.



Fig. 64.

LATERNEA BICOLUMNATA.

LATERNEA BICOLUMNATA (Fig. 64).—Receptacle consists of two columns united at the top and free at the bottom. Columns slightly compressed, cylindrical, tapering above. Gleba attached to the under side of the columns near the apex. Color pale reddish. This species is known only from Japan and is the only *Laternea* recorded from that part of the world. We are under obligations to Professor Kusano for the photograph that we reproduce.

THE GENUS PSEUDOCOLUS.

The genus *Pseudocolus* consists of columns (three, as far as known) which are united at the top and at the bottom are consolidated into a stalk. In other words, it is a stipitate *Laternea*. The best known species are from Java and Brazil. Other and less perfectly known species occur in Australia, Reunion Island (Africa), Java, and Ceylon. All species of *Pseudocolus* appear to be very rare, and most of them are only known from a single record.

PSEUDOCOLUS GARCIAE (Fig. 65).—Receptacle consists of three tapering columns, slightly united at the top and bearing the gleba on the under side. Color, white. This is a rare species, known only from Brazil (Moeller) and rare there, for Father Rick has never found it.



Fig. 66.

PSEUDOCOLUS JAVANICUS.



Fig. 68.

PSEUDOCOLUS FUSIFORMIS.

PSEUDOCOLUS JAVANICUS (Fig. 66).—From the illustration this seems to be very similar to the preceding species from Brazil. However, this is of a pale red color and grows in Java. It is a very rare plant and is only known from one specimen collected by Penzig. We reproduce Penzig's drawing, which is enlarged twofold from the plant.

PSEUDOCOLUS RUGULOSUS (Fig. 67).—Columns three, slender, united at the apex and into a short stipe at the base. The inner side of the columns are strongly rugulose, fluted. The stipe very short and included in the volva. *Color, red.* All that is known of this species is a figure preserved at Kew and made by Kurz in Java. It was referred to *Laternea triscapa*. If it exists at all it must be quite rare, for neither Penzig nor Dr. Bernard has found it.

PSEUDOCOLUS FUSIFORMIS (Fig. 68).—This species is based on a figure in the Museum at Paris, made on the Island of Reunion (near Madagascar). The plant is red; otherwise, our photograph of this figure (Fig. 68) is all that is known about it. If the plant was correctly drawn, as it seems to be, it appears to me to be very distinct from all the other species. Professor Fischer based the name *fusiformis* on this figure, afterwards withdrew it, referring the plant to *Pseudocolus Javanicus* of Java. That does not seem possible to me.



Fig. 67.

PSEUDOCOLUS RUGULOSUS.
(From the original sketch.)



Fig. 69.

PSEUDOCOLUS ROTHAE.
(From the original sketch.)

PSEUDOCOLUS ROTHAE (Fig. 69).—Columns three, slender, united above and below into a short stipe which does not extend beyond the volva. Color, rich orange. This species is represented at Kew by two collections from Australia. It seems very similar to the preceding from Java, but is evidently a much more slender species. As no other illustration of it is known, we give a copy of a crude sketch by Bailey, sent with the plant.



Fig. 65.

PSEUDOCOLUS GARCIAE.

THE GENUS CLATHRUS.

This genus has a receptacle consisting of a simple "sessile" network, bearing the gleba on the inner side. When young, the gleba forms a mass, filling the center of the egg; but as the plant expands, the gleba deliquesces and remains attached to the inner surface of the receptacle. The genus *Clathrus* as comprised in this pamphlet consists of two very distinct genera. *Clathrus* (true), with the receptacle composed of large cells, and *Ileodictyon*, with the receptacle formed of tubes. *Clathrus cancellatus* belongs to the former; *Clathrus cibarius* and *gracilis* to the latter. Where the other species belong we do not surely know, and hence do not attempt to maintain them as two genera.

CLATHRUS CANCELLATUS (Fig. 70).—Color, bright red. Meshes of the network subequal. Receptacle subglobose, composed of large cells, becoming torn and lacerate on the inner surface, the outer surface smooth, even. This well-known species is a native of Southern Europe. It is not rare in Italy and Southern France. It is a plant of warm regions and does not occur in Northern Europe except where the climate is modified by the Gulf Stream. It is found rarely on the channel coast, both of France and England, and even extends up into Holland. It occurs in Northern Africa, and has



Fig. 70.

CLATHRUS CANCELLATUS.



Fig. 71.

CLATHRUS AMERICANUS.



Fig. 74.

CLATHRUS
CAMERUNENSIS.

been collected at a few stations in Florida and Georgia in the United States. In our country it is rare, and only known with certainty from the South.

CLATHRUS AMERICANUS (Fig. 71).—Color, red. Receptacle, elongated. Meshes subequal above, elongated below. Outer surface slightly grooved, smooth. This is a species of Brazil and the West Indies. It reached me first from Father Schupp, of Brazil, who sent a photograph (Fig. 71) and a dried specimen. Then from



Fig. 72.

CLATHRUS TREUBII.

L. J. K. Brace, from the Bahamas, sent in liquid. In general form it appears to be very much like *Clathrus pusillus*, of Australia, but according to the original figure, that has much more slender branches. At Berlin I found a specimen (unnamed) from Paraguay.

CLATHRUS TREUBEI (Fig. 72).—Color, bright red. Receptacle of large meshes above, below columned. The branches of the receptacle are *tubular*, smooth externally and corrugated on the inner surface. They are reduced in diameter above, and when old they



Fig. 73.

CLATHRUS PUSILLUS.

break apart, and the primary columns separate. *Clathrus Treubei* was recently described by Dr. Bernard, from Java. At Upsala there are alcoholic specimens collected in Java by E. Nyman, and an old specimen of the same collection was discovered at Berlin to be a new species of *Laternea* (pentactina).



Fig. 74a.

CLATHRUS CRISPATUS.

(Egg.)

CLATHRUS PUSILLUS (Fig. 73).—Color, bright ruby red. Meshes subequal above, elongated below. Branches of the receptacle wrinkled. This species is only known from the original collection made on the Swan River, Australia, more than sixty years ago. What is apparently a very good figure of it (Fig. 73) was given by Berkeley, though it seems to me the branches of the receptacle are more slender than is borne out by the specimens at Kew.

CLATHRUS CAMERUNENSIS (Fig. 74).—This species was described from Camerun, Africa, and figured. The figure appears to be very much the same as *Clathrus pusillus* from Australia, but the African plant is said to be dark olive and the Australian red. The type is in alcohol at Berlin. It seems to be an *Ileodictyon* with tubular arms. The most marked feature of it to me is the reduced diameter of the upper bars.

CLATHRUS CRISPATUS (Fig. 74a).—This species is only known from the elevated regions of Ceylon and is imperfectly known from there. It was

originally sent to Europe (dried specimens) many years ago, and referred to *Clathrus cancellatus*, to which it seems to have little resemblance. The net is composed of broad, flattened bars which form small meshes. The color is red. No photograph or drawing is known (in Europe) but it must be quite different in appearance from *Clathrus cancellatus*. We reproduce a photograph of an unopened egg. This has a tubercular surface, corresponding to the form of the enclosed net, and is a character not seen at all in the European species. At the British Museum there is a species from *Yucatan* (!) that seems to be this species. Mr. Petch, who has rarely seen it, writes me that the bars are flattened-triangular in section, the broad, flat surface exterior.




Fig. 75.

CLATHRUS GUTTULATUS.

CLATHRUS GUTTULATUS (Fig. 75).—Color, bright red. Branches of the net narrow, thin, smooth. They appear to be tubes. Color, bright red. Nothing is known of this species excepting the original figure in the collection of Fries. It was made by Oersted, from St. Thomas. The guttae appear to me to be spots of white lead on the drawing, intended to show the porous nature of the receptacle.




Fig. 76.

CLATHRUS CRISPUS.

CLATHRUS CRISPUS (Fig. 76).—Color, salmon. Receptacle, subglobose, with subequal meshes. Branches of the receptacle broad,

strongly wrinkled. This seems to be a frequent species in the West Indies and is recorded also from Mexico and South America. Plumier, two hundred years ago gave a crude but evident figure of it. Next it seems to have been very characteristically figured by Turpin (Fig. 76). It was sent to Berkeley from Uruguay. It has been recorded several times, mostly from the West Indies. No photograph is known, but the original drawing seems characteristic.



Fig. 77.

CLATHRUS PSEUDOCRISPUS (reduced one-third).

CLATHRUS PSEUDOCRISPUS (Fig. 77).—A figure (Fig. 77) of what is probably only a form of *Clathrus crispus* is found at Kew from Dr. McCatty, Montego Bay, Jamaica. It differs from *crispus*, as is shown by the figure, in having the meshes below elongated. Whether it is a distinct species, a distinct form, or whether *crispus* really has this character we do not know. The color as shown is dark red.

CLATHRUS PSEUDOCANCELATUS.—This plant was named from Central Africa. It was orange-red and described as having broad, flattened branches. No figure has been given of it from which any idea whatever can be gained of the general appearance of the plant, nor could I form a much more definite idea from the types in alcohol at Berlin. They were probably originally in formalin as they have lost all definite form.



Fig. 79.

CLATHRUS GRACILIS.

CLATHRUS CIBARIUS (Fig. 78).—Color, white. Receptacle with smooth, tubular branches and large, pentagonal meshes. Our figure (78) will give an idea of the general appearance of this plant, but not of the size, for the photograph is evidently much reduced. The plant is four or five inches in diameter. It is a very common species in New Zealand, and it occurs rarely in Australia. It also grows in Chile, and a curious form has been collected in Brazil. It is said that the natives of New Zealand formerly employed the plant for food, hence the name.

CLATHRUS AFFINIS.—At the British Museum there is a specimen collected by G. A. Ramage, Pernambuco, Brazil, which is certainly a distinct form if not specifically distinct. It has the general appearance of Clathrus



Fig. 78.

CLATHRUS CIBARIUS.
(Reduced about one-half.)

cibarius, but the arms of the upper meshes are narrower than those of the lower, and the latter are somewhat columnar, so that the lower meshes are elongated.

AFRICAN FORM(?).—At the British Museum there is a plant from Africa which, if not a form of *Clathrus cibarius*, is very close. There is a sketch with it which is yellowish (and I understand that the New Zealand type form is white), but otherwise it seems to be the same.



Fig. 80.

CLATHRUS CHRYSOMYCELINUS.



Fig. 81.

CLATHRUS PREUSSII.

CLATHRUS GRACILIS (Fig. 79).—Color white or pale. Receptacle large, globose, with large meshes. The branches of the mesh are flattened, very narrow and slender, and vary from 2 to 3 mm. in breadth. *Clathrus gracilis* is the most common phalloid in Australia. There are numerous collections at Kew, and it reaches me from several collectors. It is very much like *Clathrus cibarius* of New Zealand, in fact might be considered as a small form of it. It does not seem to occur in New Zealand. It is reported from South Africa, and at

Paris there is a very poor specimen, which has been called *Clathrus Fischeri*, but which appears to be *Clathrus gracilis*. The specimen is too poor to judge, however. Notwithstanding that *Clathrus gracilis* is the most common phalloid in Australia, we know of no photograph of it and have to resort to one made from alcoholic material, devoid of volva, which gives only a vague idea of the plant.

CLATHRUS CHRYSOMYCELINUS (Fig. 80).—Receptacle white, with large, polygonal meshes; those below somewhat lengthened. The receptacle arms are united at the base. Mycelium described as being bright golden yellow, hence the specific name. This species is only known from Brazil. Father Schupp finds it, and he writes me the mycelium is not always yellow.



Fig. 82.

CLATHRUS DELICATUS.

CLATHRUS PREUSSII (Fig. 81).—This species from Kamerun, Africa, is one of the few white species of *Clathrus* known. The receptacle has broad, flat arms that are more narrow above. The figure which was published by Fischer shows the plant with the volva cut away. It is only known from the original collection in alcohol at Berlin. The bars of the network are cellular (not tubular) and have a somewhat quadrilateral shape, different from all other known species of *Clathrus*.

CLATHRUS DELICATUS (Fig. 82).—This unique little *Clathrus* is the smallest of the genus and disputes with *Mutinus xylogenus* the distinction of being the smallest phalloid known. It occurs only in Ceylon as far as known. The color is white, and the structure of the arms is tubular, hence it should be included in the genus *Ileodictyon* if taken out of *Clathrus*. The gleba is collected in little globose masses at the nodes of the net.

THE GENUS SIMBLUM.

The genus Simblum can be described in a few words as being a Clathrus on a stalk. In most of the species known the meshes are more compact than is usual in Clathrus. The genus Simblum was




Fig. 83.

SIMBLUM PERIPHRAGMOIDES.



Fig. 87.

SIMBLUM MÜLLERI.

originally known from Mauritius, then from South America, Java, and finally from the United States. At Kew we found an unnamed species from Africa, and there is a doubtful one from Australia. It can be divided into two series according to the color, yellow and red, which seem distinct and do not run into each other. However, the red series has pale or white forms.



Fig. 84.

SIMBLUM GRACILE.



Fig. 85.

SIMBLUM TEXENSE.



Fig. 86.

SIMBLUM SPHAEROCEPHALUM.

SIMBLUM PERIPHRAGMOIDES (Fig. 83).—Volva, white. Stipe 3 to 4 inches long by 2 broad, hollow, striate, yellow. Receptacle globose, with small meshes, yellow. This species, which was originally from Mauritius, was sent to Hooker and published in 1831. It is evidently rare in Mauritius, for Dr. O'Connor, who resides there and has collected several other phalloids of this island, but lately found it. The following species, which is common in Java, I at first thought was distinct from its slender form, but at Upsala I have recently seen a series of alcoholic specimens from Java, some so much like the original specimens that I now think them to be one species.




Fig. 88.

SIMBLUM CLATHRATUM.

SIMBLUM GRACILE (Fig. 84).—This has all the characters of the previous excepting the slender form. It is yellow, with a globose head of small meshes. It is a very common species in Java, Ceylon, and India, and has been reported from China. I am convinced, from an examination of a series of alcoholic specimens from Java, at Upsala, that it can not be kept distinct from the preceding species.

SIMBLUM TEXENSE (Fig. 85).—This species, which is only known from Texas, has the same yellow color character as the preceding. It differs in the nature of the network (best shown in our figures) and in the clathrate portion abruptly contracted into the stipe. An excellent account of it has been given by Professor Long.

SIMBLUM SPHAEROCEPHALUM (Fig. 86).—This species differs from those that precede by being red, though pale or white forms occur. It was first noted in South America, where it is an extremely common plant. Then it was published from the United States,¹³ where it is rare, and it reached me from the Bahamas. In shape it is the same as *Simblum Texense*, and the photographs without color notes could not be told apart.

SIMBLUM MÜLLERI (Fig. 87).—This species, which is known from a drawing made from a dried specimen from Australia, is very different from all others in its open network. In its general appearance it is close to *Clathrus pusillus*, excepting that the clathrate portion is borne on a distinct though short stem. When the phalloids of Australia are well known, it may be found that *Clathrus pusillus* varies in this respect and that this is really only a stalked form.

SIMBLUM CLATHRATUM (Fig. 88).—Stem hollow, pale reddish tint, 2½ cm. thick x 7 cm. high. Receptacle a loose, clathrate structure, with large meshes to the net and slender branches. Color, bright red. The clathrate portion is fragile and easily broken. The specimen grew in the botanical garden at Old Calabar, Africa. It is the first red *Simblum* known from Africa, although the original species of the genus came from Mauritius. It was a yellow plant. The only similar plant known is *Simblum sphaerocephalum* from America, which differs widely in having a compact net of small meshes. The specimen and a colored drawing by J. W. Holland are at Kew.

THE GENUS COLUS.

This genus is a *Clathrus* supported on columns which are united at the base into a stipe. Only one species is known, and that only from the Mediterranean regions.

COLUS HIRUDINOSUS (Figs. 89 and 90).—This is a small phalloid, that, as far as is known, grows only in the Mediterranean regions. Originally from Corsica, it was named from Southern France. It has been found in Algeria, and Father Torrend, of Portugal, has recently discovered it abundant in the sand. In Corsica, the original observer stated, it grew only on manure, but the other records are from unnamed places. The color is red; the other characters are all those of the genus and are best shown in our photographs.

In some publications, the genus *Colus* includes plants that in my opinion have very little resemblance or relation to the original species. These we have separated under the name *Pseudocolus*.

¹³ The only stations known are Long Island, N. Y., Gerard; Nebraska, J. M. Bates; Kansas, E. E. Bartholomew; Washington, D. C., W. H. Scudder; Talbot County, Maryland, Chas. McIlvaine. When any one finds this rare plant in the United States I request that it be reported to me so that we can keep a record of its known stations.



Fig. 89.

COLUS HIRUDINOSUS.
(Natural size.)



Fig. 90.

COLUS HIRUDINOSUS.
(Enlarged.)

THE GENUS KALCHBRENNERA.

This is a very peculiar genus, known only from South Africa, and but one species. It has a stipe bearing a clathrate structure similar to the genus *Simblum*, but from the net proceed large, knobbed projections.

KALCHBRENNERA CORALLOCEPHALA (Figs. 91 and 92).—The only species grows in South Africa, and there appears to be rather a frequent plant. It is a very showy plant, of a bright red color in all its parts. The gleba covers the outer portion of the net and to an extent hides the network. It was a number of years before its correct structure was known, and it was Kalchbrenner who made a good picture of it and first showed it.



Fig. 91.

**KALCHBRENNERA CORALLO-
CEPHALA.**




Fig. 92.

KALCHBRENNERA CORALLOCEPHALA.
(Section.)

RELATED PLANTS.

It is a disputed question whether *Phallogaster saccatus* is a phalloid or not. It has no volva as other phalloids have, hence is excluded by some who are theorizing on such things. I do not believe that any one familiar with the fresh plant will ever place it anywhere except with the phalloids. It has the same greenish, fetid gleba that is associated with the phalloids, the same spores and basidia, and it deliquesces in the same way. It seems to me that its relations are entirely with the phalloids, notwithstanding it has no volva.

THE GENUS PHALLOGASTER.

Plants devoid of a volva, the gleba borne in the inner tissue. Peridium white, smooth. In ripening the inner tissue and gleba deliquesce, and the latter adheres to the inner side of the peridium, which breaks irregularly and exposes the adhering gleba.



Fig. 93.

PHALLOGASTER SACCATUS.



Fig. 94.

PHALLOGASTER SACCATUS.

(After dehiscence.)

PHALLOGASTER SACCATUS (Figs. 93 and 94).—This species occurs only in the United States and Canada, as far as known, and it is a rare plant there. It has only been known for a few years. I think there can be no trouble in identifying it from our photographs. Another species has been recently published, which appears to me to be rather a depauperate form.

APPENDIX I.
GEOGRAPHICAL DISTRIBUTION.

The real study of the phalloids, as I view it, is the correct characters of the species, the simplest grouping of them into genera, and their distribution. We present a synopsis of the number of phalloids known to occur in various countries, and where the same species occurs in different countries it is included in each. We include as different phalloids all the various forms named in this pamphlet, and all the alleged species so named, whether doubtful or well known.

General Distribution.

When the subject is well known, I think, it will be found that several species are of very wide distribution, but at present we only know two.

Phallus indusiatus occurs without doubt in every tropical country of the world. We give on page 18 the countries from which we have seen specimens, and the list does not embrace perhaps half of the countries where it occurs.

Phallus rubicundus (under the names *aurantiacus*, *gracilis*, etc.) also seems to occur in most warm countries.

	Europe.	North America.	West Indies.	South America.	Australia and New Zealand.	Africa.	Ceylon and India.	Japan and China.	East Indies.
<i>Anthurus</i> ,	3	..	1
<i>Aseroe</i> ,	6	1	2	2	2
<i>Clathrus</i> ,	1	1	4	5	3	4	2	..	1
<i>Clautriavia</i> ,	1	..	1	..	1
<i>Colus</i> ,	1	1
<i>Floccomutinus</i> ,	1
<i>Itajahya</i> ,	1
<i>Jansia</i> ,	1	..	1	1	2
<i>Kalchbrennera</i> ,	1
<i>Laternea</i> ,	1	3	3	1(?)	1	..	1	..
<i>Lysurus</i> ,	1	1	..	3	2	1	1	1	..
<i>Mutinus</i> ,	1	3	..	3	3	..	1	2	3
<i>Phallogaster</i> ,	1
<i>Phallus</i> ,	2	5	2	8	10	6	3	5	9
<i>Pseudocolus</i> ,	1	1	1	2
<i>Simblum</i> ,	2	1	1	1	2	1
Total ,	6	14	10	25	31	19	12	12	21

Europe.

There are but six phalloids in Europe (including one form). *Phallus impudicus* is the most common and widespread. The form *Phallus imperialis* is rare and local. *Mutinus caninus* is not rare. *Clathrus cancellatus* is of a southern range. It occurs mostly in southern France, Italy, etc. *Colus hirudinosis* is confined to the Mediterranean region. It occurs in Corsica, Southern France, Portugal. *Lysurus borealis* is probably an introduced species. It is known from one collection in Germany and two in England.

United States and Canada.

We have fourteen phalloids in our country. *Phallus Ravenelii* and *Phallus duplicatus* are the most frequent. The form *Phallus imperialis*, which with us replaces *Phallus impudicus* of Europe, is of a Western range, found in California, Colorado, and Texas. But one Eastern station is known, Washington, D. C. *Mutinus elegans* is our most common *Mutinus* in rich woods. *Mutinus caninus* is an Eastern species, and *Mutinus Ravenelii* is local and rare. *Laterna columnata* is common in the South, and *Clathrus cancellatus* is very rare and only known with certainty from the South. *Phallus rubicundus* seems to be fairly common in the South. *Simblum sphaerocephalum* is very rare. A list of known stations is given on page 67. *Simblum Texense* is known only from Texas. *Lysurus borealis* seems to be an introduced plant. Of late years it has been found a number of times, chiefly in the East and in cultivated stations. *Phallogaster saccatus* is of rare occurrence. In addition I have a specimen in alcohol from Florida, species not sure, but probably *Phallus gracilis*.

West Indies.

The phalloids of the West Indies are not well known. Undoubtedly when well observed, several of the Brazilian species will be found in the West Indies. *Clathrus crispus* (and a doubtful form, *pseudocrispus*); *Clathrus Americanus*, recently found in the Bahamas by Mr. Brace; *Clathrus guttulatus*, known only from an old drawing; *Phallus indusiatus*, common; *Phallus rubicundus*, probably common; *Laterna columnata*, common; *Laterna pusilla*, known from one collection; *Laterna triscapa*, known only from an old drawing, and *Simblum sphaerocephalum*, recently collected in the Bahamas by Mr. Brace.

South America.

Most excellent work has recently been done on the phalloids of Brazil by Moeller, and to this work is due most of our knowledge of South American phalloids. He has published in a superb manner *Clathrus chrysomycelinus*, *Pseudocolus Garciae*, *Laterna columnata*, *Laterna rhacodes*, *Mutinus bambusinus*, *Phallus subtilis*, *Phallus glutinolens*, *Phallus indusiatus* (and a form, Moelleri), *Phallus callichrous* (which is probably only a color form of *indusiatus*), and *Itajahya galericulata*, a genus only known from South America.

Simblum sphaerocephalum is a most common phalloid in South America, but does not seem to have been found by Professor Moeller. *Clathrus Americanus* is a recent species from Rev. F. A. Schupp, Brazil.

Rev. J. Rick finds in his locality (Sao Leopolda) the following: *Simblum sphaerocephalum*, *Phallus indusiatus*, *Pseudocolus Garciae*, *Laterna columnata*, *Laterna rhacodes*, and *Clathrus Americanus*.

There have been several imperfectly known phalloids from South America. We would list *Phallus Farlowii*, *Mutinus australis*, *Lysurus Sanctae-Catherinae*, *Phallus roseus* (a form of *indusiatus*), *Mutinus xylogenus*, *Lysurus cruciatum*, *Phallus campanulatus*, *Lysurus Clarazianus*, *Clathrus affinis* (a form of *cibarius*, known only from a specimen in the British museum), *Laterna Spegazzini*, and *Laterna crispus*. In addition, several have been proposed by Spegazzini, but they are mostly only word-descriptions, and nothing can be told about them. For me an unillustrated phalloid has no place excepting in the rejected columns. There has also been a "new genus," *Alboffiella*, illustrated by Spegazzini. Professor Fischer has suggested, not without reason it seems to me, that it was based on a *Phallus* with an accidental volva cap. Robert E. Fries suggests it was based on *Itajahya galericulata*. If true, in either case, the work was very poorly done.

Australia and New Zealand.

I consider the phalloids of Australia and New Zealand for the most part very imperfectly and inaccurately known. The new species were mostly proposed forty or fifty years ago and illustrated by figures reconstructed from dried specimens, often inaccurate it seems to me, and nothing since has been learned of them. The subject has gotten into such a condition that the local workers in these countries seem to be able to make but very little of their species, and the result is there have been very few original papers by the mycologists of these countries. It is time our friends there observed their phalloids and gave us good accounts and photographs of them. If Australian mycologists will take as a model the photographs and account given on page 42 by Professor D. McAlpine, of *Anthurus aseroeformis*, and supply similar photographs and accounts, it will only be a few years until we have a much better knowledge of the subject.

The two most frequent phalloids are *Clathrus cibarius* and *Clathrus gracilis*, the former in New Zealand, the latter in Australia. Neither has been satisfactorily illustrated. *Anthurus aseroeformis*, a rare species but well known, due to Professor McAlpine. *Phallus indusiatus* is a frequent plant, but the forms and color forms are not worked out. The genus *Aseroe* is at home in Australia. It seems to take very different forms, but their value in classification is not known. With the exception of the above, I consider all the other Australian species more or less doubtful and little known, viz: *Phallus impudicus*, *Phallus rubicundus*, *Phallus multicolor*, *Phallus callichrous*, *Phallus Rochesterensis*, *Phallus discolor*, *Phallus calypttratus*, *Phallus retusus*, *Phallus quadricolor*, *Mutinus pentagonus*, *Mutinus curtus*, *Mutinus papuasius*, *Jansia annulata*, *Lysurus Australiensis*, *Lysurus* (unnamed), *Anthurus Muellerianus*, *Anthurus Archeri*, *Aseroe* (all the five recorded forms, *rubra*, *pentactina*, *Hookeri*, *Muelleriana*, *lysuroides*), *Laternea columnata* (very ??), *Pseudocolus Rothae*, *Clathrus pusillus*, *Simblum Mülleri*. In addition, there is a curious species, *Clautriavia Lauterbachii*, only known from an egg from the neighboring island of New Guinea, and a pale *Aseroe* (*pallida*) is recorded from New Caledonia.

Samoa.

I have spent two winters in Samoa and have hunted the fungi thoroughly. I am satisfied that *Phallus indusiatus* is the only common phalloid that grows on the island, and it is not at all rare. In the museum at Berlin is a specimen labeled *Clathrus gracilis* (and it seems to be correct), also a *Mutinus* (unnamable). Both genera must be very rare in Samoa, as I found neither.

Africa.

Many years ago *Simblum periphragmoides* was well illustrated from Mauritius, and was only recently found again. A slender form is very frequent in the East Indies. *Kalchbrennera corallocephala*, a most striking species, was well illustrated by Kalchbrenner thirty years ago. *Phallus indusiatus* is a common species and has reached me several times from Africa. *Colus hirudinosus* occurs in North Africa. *Lysurus Woodii*, *Laternea Angolensis*, *Phallus subacutus*, and *Phallus canariensis* were imperfectly published years ago, and nothing has been added to them since.

In recent years Africa has been a fertile field for "new species," but the work has not been done as it should have been. Such work would have passed forty years ago, but it is out of date now. The following have been added, mostly in this manner, in comparatively recent years: *Floccomutinus Zenkeri*, *Phallus rubicundus* (?), *Clathrus camerunensis*, *Clathrus pseudocancellatus*, *Clathrus Preussii*, *Clathrus gracilis* (?), *Simblum clathratum*, *Pseudocolus fusiformis*, *Phallus callichrous*, *Clathrus cibarius*(?). Fine specimens of many of these are in alcohol in the museum at Berlin.

Mr. Chas. O'Connor has been observing the phalloids of Mauritius. He finds the only common one to be *Phallus gracilis*. More rarely he has observed *Phallus indusiatus* and *Phallus Mauritanus*, a related plant. He has only recently rediscovered *Simblum periphragmoides* which was originally from Mauritius, but is very rare there.

Ceylon.

For many years we have had a very imperfect knowledge of the phalloids of Ceylon, but a very recent paper by T. Petch has set the matter right. The following species occur in Ceylon: *Jansia rugosa* (rare, and considered by Petch to be *Mutinus proximus*), *Mutinus proximus* (known only from dried specimens and sketch), *Phallus tenuis* (only previously known from Ceylon from dried specimen at Kew, but recently again reported from Ceylon), *Phallus indusiatus*, the most common phalloid and takes many color forms, viz: callichrous and multicolor), *Clautriavia merulina* (common in the Botanic Gardens at Peradeniya), *Simblum gracile*, common, *Lysurus Gardneri* (rare in Ceylon, but most abundantly represented in the museums at Kew, there being 25 specimens), *Aseroe Zeylandica*, rare in the elevated regions, *Aseroe arachnoidea*, very rare. In addition the unique little *Clathrus delicatus* is only known from Ceylon.

India.

Seventy years ago Perrottet sent Montagne a few phalloid sketches and dried specimens on which were based *Phallus rubicundus* (published as *aurantiacus*), *Anthurus Calathiscus* (supposed to have been very inaccurately published). In addition a few specimens of *Phallus indusiatus* have reached Europe from India, and these are all, I think, that are known from India. At the British Museum there are ten times as many specimens of extinct elephant remains from India as there are of the *live phalloids* that every naturalist in India must observe.

Mr. Hutchins writes me from North Bengal that *Phallus indusiatus* is common, but is the only phalloid he finds. Mr. G. H. Krumbiegel sent me from North Bengal a dried phalloid which, while I would not attempt to reconstruct it, I recognize as a genus unknown.

Java.

From no country in the world have we had a better account of the phalloids than from Java, which was published by Penzig. The following were well illustrated and described by him: *Mutinus bambusinus*, *Mutinus Fleischeri*, *Jansia elegans*, *Jansia rugosa*, *Phallus tenuis*, *Phallus costatus* (form?), *Phallus favosus* (form?), *Phallus indusiatus*, *Clautriavia merulina*, *Phallus multicolor*, *Simblum gracile* (form), *Pseudocolus Javanicus*, *Aseroe arachnoidea*.

Dr. Chas. Bernard has given us a good photograph and account of *Clathrus Treubei*, and has sent me a collection of the Javanese species in alcohol, from which some good photographs have been made.

Aseroe Zeylandica (under the name *Junghuhnii*) was published from Java many years ago, but is very rare and only rediscovered by Dr. Bernard recently. *Pseudocolus rugulosus* is based on an old drawing from Java, and no specimen is known. From the neighboring islands, *Mutinus borneensis* is vaguely described from Borneo and *Phallus celebicus* from the Celebes.

Dr. Chas. Bernard gives the following synopsis of the relative frequency of the phalloids he has observed in Java: *Mutinus bambusinus*, *Clautriavia merulina*, *Phallus indusiatus*, and *Simblum gracile* are common throughout the season, though more abundant, of course, during the rainy season. *Aseroe arachnoidea*, *Jansia elegans*, *Jansia rugosa*, *Phallus multicolor*, and *Clathrus Treubei* are rarer species and will probably only be found during the rainy season. *Aseroe Zeylandica* is a very rare phalloid and only recently rediscovered.

Japan.

An account of the phalloids of Japan was published in *Mycological Notes*, page 400. It was based on notes, drawings, and specimens from Professors Kusano, Gono, and Yasuda. The following were included: *Phallus indusiatus*, *Phallus impudicus*, *Phallus rugulosus*, *Phallus tenuis* (rare), *Jansia boninensis* (as *Mutinus*), *Lysurus Mokusin*, and *Laternea bicolonnata*. In addition, *Phallus rubicundus* under the name *aurantiacus* has been said to grow in Japan.

China.

Little is known as to the phalloids of China, although *Lysurus Mokusin* from China was among the first foreign phalloids figured.

Some alcoholic specimens were sent to Patouillard at Paris a few years ago from Tonkin, and the following species recorded: *Aseroe Zeylandica*, *Phallus indusiatus*, *Phallus gracilis*, *Mutinus bambusinus*, *Mutinus minimus*, *Mutinus borneensis*.

APPENDIX II.

LOST, STRAYED, OR STOLEN.

The following phalloids have not been heard from since they were originally exploited and grave fears are entertained as to their survival. Vague rumors have been circulated of one or two of them having been seen, but when traced to the source have usually resulted from a mistaken identification. Any one noticing a stray phalloid in their neighborhood is requested to seize it and send it in with such notes and marks as may lead to its identification.

	Whence Exploited	Has not been heard from for
<i>Anthurus Müllerianus</i>	Australia	Thirty years.
<i>Anthurus Archeri</i>	Australia	Fifty years.
<i>Anthurus Calathiscus</i>	India	Sixty-eight years.
<i>Clathrus pusillus</i>	Australia	Sixty-five years.
<i>Laternea pusilla</i>	Cuba	Forty years.
<i>Laternea triscapa</i>	West Indies	Eighty-seven years.
<i>Laternea angolensis</i>	Africa	Forty years.
<i>Lysurus cruciatus</i>	French Guiana	Sixty-five years.
<i>Lysurus Clarazianus</i>	South America	Thirty-six years.
<i>Lysurus Sanctae Catherinae</i>	South America	Twenty years.
<i>Lysurus Woodii</i>	South Africa	Thirty years.
<i>Mutinus curtus</i>	Australia	Sixty-five years.
<i>Mutinus papuasius</i>	Australia	Thirty years.
<i>Mutinus discolor</i>	Australia	Thirty years.
<i>Mutinus xylogenus</i>	French Guiana	Fifty-five years.
<i>Phallus Daemonum</i>	Amboy	One hundred and sixty-six years.
<i>Phallus quadricolor</i>	Australia	Twenty-six years.
<i>Phallus calyptratus</i>	Australia	Twenty-six years.
<i>Phallus retusus</i>	Australia	Twenty-five years.
<i>Phallus subuculatus</i>	North Africa	Sixty years.
<i>Pseudocolus fusiformis</i>	Reunion	Thirty years.
<i>Simblum Mülleri</i>	Australia	Twenty years.

APPENDIX III.

SYNONYMS.

There have been nearly three hundred names proposed for phalloids and only about one hundred have been retained in this pamphlet. The other two-thirds are, in our opinion, superfluous. It is an easy matter to propose a new name, but when once proposed it is impossible to ever get rid of it. Writers can refer it to "synonymy" all they please, but the next man that comes along has to dig it up and go all over it again, for no two men ever agree as to all the details, and each man is entitled to his own opinion.

Many of the following names are the discoveries made by those who discover "new species," which seem to me to have been "new" chiefly to the discoverer. A large part of the synonyms are from changing plants from one genus to another or making new genera out of sections of old genera. Personally we do not maintain many of these innovations, for the old established genera seem better to us. Of the new genera proposed in the last twenty years we only maintain *Itajahya*, *Jansia*, *Phallogaster*, *Clautriavia*, *Floccomutinus*, and *Pseudocolus*. (The latter we had the assurance to propose ourselves.) Professor Fischer has worked over this same ground and reduced many of these same names to synonymy, and while we agree with him in many instances we have copied him in none, for in every case we have looked up the evidence and formed our own opinion. We have not been as free as he in reducing species, for perhaps twenty names recognized as good in this pamphlet Professor Fischer puts in synonymy. While we suspect many of these have little value, we give them, in all instances, the benefit of the doubt.

There is one class of "new species" exploiters that I have not bothered much with—those who propose new species without illustrating them. In a subject such as the phalloids, where a good illustration tells most of the story, there is no excuse for any one to try to describe a phalloid in words. It ought to be a recognized crime, with a heavy penalty. Such species are listed here as "Nomina nuda." The phalloid fakers who fake up pictures are perhaps worse. The following names are those which in our opinion should be placed in synonymy and the reasons.

<i>Anthurus australiensis</i>	See <i>Lysurus</i> .
<i>Anthurus borealis</i>	See <i>Lysurus</i> .
<i>Anthurus Clarazianus</i>	See <i>Lysurus</i> .
<i>Anthurus cruciatus</i>	See <i>Lysurus</i> .
<i>Anthurus Sanctae Catherinae</i>	See <i>Lysurus</i> .
<i>Anthurus trifidus</i>	Nomen nudum.
<i>Anthurus Woodii</i>	See <i>Lysurus</i> .
<i>Aporophallus subtilis</i>	See <i>Phallus</i> .
<i>Albofiella argentina</i>	Supposed to be a break.
<i>Aseroe actinobolus</i>	= <i>Aseroe pentactina</i> .
<i>Aseroe Ceylanica</i>	See <i>Aseroe Zeylandica</i> .
<i>Aseroe Calathiscus</i>	See <i>Anthurus</i> .
<i>Aseroe corrugata</i>	Nomen nudum.
<i>Aseroe Junguhnii</i>	= <i>Aseroe Zeylandica</i> .
<i>Aseroe multiradiata</i>	= <i>Aseroe Zeylandica</i> probably.
<i>Aseroe viridis</i>	= <i>Aseroe Hookeri</i> .
<i>Aserophallus cruciatus</i>	See <i>Lysurus</i> .
<i>Blumenavia rhacodes</i>	See <i>Laternea</i> .
<i>Blumenavia usambarensis</i>	= <i>Laternea angolensis</i> (?).
<i>Calathiscus Sepia</i>	See <i>Anthurus Calathiscus</i> .
<i>Calathiscus Puiggarii</i>	Nomen nudum.
<i>Caromyxa elegans</i>	See <i>Mutinus</i> .
<i>Clathrella camerunensis</i>	See <i>Clathrus</i> .
<i>Clathrella crispa</i>	See <i>Clathrus</i> .
<i>Clathrella chrysomycelina</i>	See <i>Clathrus</i> .
<i>Clathrella delicata</i>	See <i>Clathrus</i> .

<i>Clathrella</i> <i>Muelleri</i>	See <i>Simblum</i> .
<i>Clathrella pseudocancellata</i>	Nomen nudum.
<i>Clathrella</i> <i>Preussii</i>	See <i>Clathrus</i> .
<i>Clathrella pusilla</i>	See <i>Clathrus</i> .
<i>Clathrella</i> <i>Treubei</i>	See <i>Clathrus</i> .
<i>Clathrus</i> <i>angolensis</i>	See <i>Laternea</i> .
<i>Clathrus</i> <i>albidus</i>	= <i>Clathrus gracilis</i> .
<i>Clathrus</i> <i>australis</i>	Nomen nudum.
<i>Clathrus</i> <i>Baumii</i>	Nomen nudum.
<i>Clathrus</i> <i>Berkeleyi</i>	= <i>Laternea pusilla</i> .
<i>Clathrus</i> <i>Brasiliensis</i>	= <i>Laternea columnata</i> .
<i>Clathrus</i> <i>columnatus</i>	See <i>Laternea</i> .
<i>Clathrus</i> <i>colonnarius</i>	= <i>Laternea columnata</i> .
<i>Clathrus</i> <i>Fischeri</i>	= <i>Clathrus gracilis</i> (?)
<i>Clathrus</i> <i>hirudinosus</i>	See <i>Colus</i> .
<i>Clathrus</i> <i>intermedius</i>	Nomen nudum.
<i>Clathrus</i> <i>parvulus</i>	Too poorly illustrated.
<i>Clathrus pseudocancellatus</i>	Nomen nudum.
<i>Clathrus</i> <i>Tepperianus</i>	= <i>Clathrus gracilis</i> .
<i>Clathrus</i> <i>triscapus</i>	See <i>Laternea</i> .
<i>Clathrus</i> <i>trilobatus</i>	= <i>Laternea columnata</i> .
<i>Colus</i> <i>fusiformis</i>	See <i>Pseudocolus</i> .
<i>Colus</i> <i>Garciae</i>	See <i>Pseudocolus</i> .
<i>Colus</i> <i>Gardneri</i>	See <i>Lysurus</i> .
<i>Colus</i> <i>Javanicus</i>	See <i>Pseudocolus</i> .
<i>Colus</i> <i>Muelleri</i>	See <i>Simblum</i> .
<i>Colus</i> <i>Rothae</i>	See <i>Pseudocolus</i> .
<i>Colonnaria</i> <i>truncata</i>	Rafinesque's ravings.
<i>Colonnaria</i> <i>urceolata</i>	Rafinesque's ravings.
<i>Corynites</i> <i>brevis</i>	= <i>Mutinus Ravenelii</i> .
<i>Corynites</i> <i>Curtisii</i>	= <i>Mutinus elegans</i> .
<i>Corynites</i> <i>elegans</i>	See <i>Mutinus</i> .
<i>Corynites</i> <i>Ravenelii</i>	See <i>Mutinus</i> .
<i>Cryptophallus</i> <i>albiceps</i>	= <i>Phallus imperialis</i> .
<i>Cynophallus</i> <i>bambusianus</i>	See <i>Mutinus</i> .
<i>Cynophallus</i> <i>caninus</i>	See <i>Mutinus</i> .
<i>Cynophallus</i> <i>papuasius</i>	See <i>Mutinus</i> .
<i>Dictybole</i> <i>texensis</i>	A phalloid fake.
<i>Dictyophallus</i> <i>aurantiacus</i>	= <i>Phallus rubicundus</i> .
<i>Dictyophallus</i> <i>discolor</i>	See <i>Phallus</i> .
<i>Dictyophora</i> <i>bicampanulata</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>brasiliensis</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>Braunii</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>callichrous</i>	See <i>Phallus</i> .
<i>Dictyophora</i> <i>campanulata</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>chlorocephala</i>	= <i>Phallus callichrous</i> .
<i>Dictyophora</i> <i>collaris</i>	= <i>Phallus duplicatus</i> .
<i>Dictyophora</i> <i>Daemonum</i>	See <i>Phallus</i> .
<i>Dictyophora</i> <i>duplicata</i>	See <i>Phallus</i> .
<i>Dictyophora</i> <i>echinata</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>Farlowii</i>	See <i>Phallus</i> .
<i>Dictyophora</i> <i>irpicina</i>	= <i>Clautriavia merulina</i> .
<i>Dictyophora</i> <i>Lilloi</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>merulina</i>	See <i>Clautriavia</i> .
<i>Dictyophora</i> <i>multicolor</i>	See <i>Phallus</i> .
<i>Dictyophora</i> <i>nana</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>phalloidea</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>radicata</i>	= <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>rosea</i>	= <i>Phallus indusiatus</i> (form).
<i>Dictyophora</i> <i>speciosa</i>	See <i>Phallus indusiatus</i> .
<i>Dictyophora</i> <i>subuculata</i>	See <i>Phallus</i> .

<i>Dictyophora tahitensis</i>	= <i>Phallus indusiatus</i> .
<i>Echinophallus Lauterbachii</i>	See <i>Clautriavia</i> .
<i>Floccomutinus Nymanianus</i>	= <i>Jansia rugosa</i> .
<i>Foetidaria coccinea</i>	= <i>Simblum sphaerocephalum</i> .
<i>Hymenophallus alboindusiatus</i>	= <i>Phallus indusiatus</i> .
<i>Hymenophallus brasiliensis</i>	= <i>Phallus indusiatus</i> .
<i>Hymenophallus duplicatus</i>	See <i>Phallus</i> .
<i>Hymenophallus Hadriani</i>	= <i>Phallus impudicus</i> .
<i>Hymenophallus indusiatus</i>	See <i>Phallus</i> .
<i>Hymenophallus radicans</i>	= <i>Phallus indusiatus</i> .
<i>Hymenophallus roseus</i>	= <i>Phallus indusiatus</i> (form).
<i>Hymenophallus speciosus</i>	= <i>Phallus indusiatus</i> .
<i>Hymenophallus subuculatus</i>	See <i>Phallus</i> .
<i>Hymenophallus tahitensis</i>	= <i>Phallus indusiatus</i> .
<i>Hymenophallus togatus</i>	= <i>Phallus duplicatus</i> .
<i>Hymenophallus tunicatus</i>	= <i>Phallus indusiatus</i> .
<i>Neodictyon cibarium</i>	See <i>Clathrus</i> .
<i>Neodictyon gracile</i>	See <i>Clathrus</i> .
<i>Ithyphallus aurantiacus</i>	= <i>Phallus rubicundus</i> .
<i>Ithyphallus Balansoe</i>	= <i>Phallus rubicundus</i> .
<i>Ithyphallus calyptratus</i>	See <i>Phallus</i> .
<i>Ithyphallus campanulata</i>	See <i>Phallus</i> .
<i>Ithyphallus canariensis</i>	See <i>Phallus</i> .
<i>Ithyphallus celebicus</i>	See <i>Phallus</i> .
<i>Ithyphallus coralloides</i>	= <i>Phallus rubicundus</i> .
<i>Ithyphallus costatus</i>	See <i>Phallus</i> .
<i>Ithyphallus cucullatus</i>	= <i>Phallus Ravenelii</i> .
<i>Ithyphallus favosus</i>	See <i>Phallus</i> .
<i>Ithyphallus glutinolens</i>	See <i>Phallus</i> .
<i>Ithyphallus impudicus</i>	See <i>Phallus</i> .
<i>Ithyphallus Lauterbachii</i>	See <i>Clautriavia</i> .
<i>Ithyphallus Muellerianus</i>	= <i>Phallus retusus</i> .
<i>Ithyphallus Novae Hollandiae</i>	= <i>Phallus gracilis</i> .
<i>Ithyphallus purpuratus</i>	= <i>Phallus imperialis</i> .
<i>Ithyphallus quadricolor</i>	See <i>Phallus</i> .
<i>Ithyphallus Ravenelii</i>	See <i>Phallus</i> .
<i>Ithyphallus retusus</i>	See <i>Phallus</i> .
<i>Ithyphallus rubicundus</i>	See <i>Phallus</i> .
<i>Ithyphallus rugulosus</i>	See <i>Phallus</i> .
<i>Ithyphallus sanguineus</i>	= <i>Phallus rubicundus</i> ?
<i>Ithyphallus tenuis</i>	See <i>Phallus</i> .
<i>Jansia Nymaniana</i>	= <i>Jansia rugosa</i> .
<i>Jansia Zenkeri</i>	See <i>Floccomutinus</i> .
<i>Kalchbrennera Tuckii</i>	= <i>Kalchbrennera corallocephala</i> .
<i>Kirchbaumia imperialis</i>	See <i>Phallus</i> .
<i>Laterna australis</i>	Nomen nudum.
<i>Laterna pentactina</i>	= <i>effete Clathrus Treubei</i> .
<i>Lysurus Archeri</i>	See <i>Anthurus</i> .
<i>Lysurus argentinus</i>	Nomen nudum.
<i>Lysurus aseroeformis</i>	See <i>Anthurus</i> .
<i>Lysurus Beauvaisi</i>	= <i>Lysurus Mokusin</i> .
<i>Lysurus corallocephalus</i>	See <i>Kalchbrennera</i> .
<i>Lysurus pentactinus</i>	= <i>Anthurus Archeri</i> .
<i>Lysurus Texensis</i>	Nomen nudum.
<i>Mutinus annulatus</i>	See <i>Jansia</i> .
<i>Mutinus boninensis</i>	See <i>Jansia</i> .
<i>Mutinus bovinus</i>	= <i>Mutinus elegans</i> .
<i>Mutinus brevis</i>	= <i>Mutinus Ravenelii</i> .
<i>Mutinus Curtisii</i>	= <i>Mutinus elegans</i> .
<i>Mutinus discolor</i>	See <i>Phallus</i> .
<i>Mutinus elegans</i> (of Java only)	See <i>Jansia</i> .

Mutinus Muelleri	=Mutinus bambusinus.
Mutinus Nymanianus	=Jansia rugosa.
Mutinus proximus	Nomen nudum.
Mutinus proximus (in sense of Petch)	=Jansia rugosa.
Mutinus Watsoni	Nomen nudum.
Mutinus Zenkeri	See Floccomutinus.
Omphalophallus Muellerianus	=Phallus retusus.
Omphalophallus retusus	See Phallus.
Phallogaster whitei	=depauperate Phallogaster saccatus.
Phallus aurantiacus	=Phallus rubicundus.
Phallus bambusinus	See Mutinus.
Phallus brasiliensis	=Phallus indusiatus.
Phallus caninus	See Mutinus.
Phallus collaris	=Phallus duplicatus.
Phallus curtus	See Mutinus.
Phallus foetidus	=Phallus impudicus.
Phallus Hadriani	Based on a freak.
Phallus irpicinus	=Clautriavia merulina.
Phallus inodoratus	=Phallus impudicus.
Phallus iosmos	=Phallus impudicus.
Phallus merulinus	See Clautriavia.
Phallus Mokusin	See Lysurus.
Phallus Muellerianus	=Phallus retusus.
Phallus Novae Hollandiae	=Phallus gracilis.
Phallus purpuratus	=Phallus imperialis.
Phallus radicans	=Phallus indusiatus.
Phallus sanguiners	=Phallus rubicundus (?).
Phallus senegalensis	=Imagination chiefly.
Phallus speciosus	=Phallus indusiatus.
Phallus tahitensis	=Phallus indusiatus.
Phallus truncatus	=Mutinus unknown.
Phallus tunicatus	=Phallus indusiatus.
Phallus Watsoni	Nomen nudum.
Phallus xylogenus	See Mutinus.
Protubera Maracuja	(Not for me a phalloid.)
Satyrum rubicundus	See Phallus.
Simblum australe	=Simblum sphaerocephalum.
Simblum flavescens	=Simblum gracile.
Simblum Lorentzii	=Simblum sphaerocephalum.
Simblum pilidiatum	=Simblum sphaerocephalum.
Simblum rubescens	=Simblum sphaerocephalum.
Sophronia brasiliensis	=Phallus indusiatus.
Staurophallus senegalensis	=Something unknown.
Xylophallus xylogenus	See Mutinus.

APPENDIX IV.

LIST OF PHALLOIDS IN THE MUSEUMS.

All specimens are not listed, for some are so uncertain that I feel they should not be recorded. In case a plant has been named from these specimens, I sometimes record it under this name, even if I do not maintain it as a valid species.

KEW, ENGLAND.

Phallus multicolor, 3 collections, Australia—*Phallus namus*, type, Andaman Island—*Phallus indusiatus*, Australia, Africa, British Guiana, Uganda, India, Ceylon, Java, Brazil, Mexico, Surinam Cape, several specimens from each country, also Cuba (? depauperate), Australia (var. *Rochesterensis*)—*Phallus duplicatus*, Carolina—*Phallus Ravenelii*, Connecticut—*Phallus truncatus*, poor specimens, and a drawing from which it appears to me to be rather a *Mutinus*—*Phallus rubicundus*, Southern United States, several—*Phallus rubicundus* (as *aurantiacus*), several from Australia and the form *gracilis*—*Phallus impudicus*, Australia, one only from Bailey and doubtful, England several (one the type of *iosmos*), East Indies but very ??, France, Germany—*Phallus tenuis*, Ceylon, named *Phallus pallidus* by Berkeley but never published—*Phallus aurantiacus*, co-type, ex Montagne—*Phallus gracilis*, South Africa—*Mutinus curtus*, Australia, type—*Mutinus elegans* (type of *Corynites Curtisii*)—*Mutinus Ravenelii*, type, one with a short apex called *brevis*—*Mutinus bambusinus*, nice drawing from Kurz, I think; Java, also dried from Java, also adventitious in hothouses, England—*Mutinus caninus*, a number from England—*Mutinus proximus*, type, Ceylon, poor—*Mutinus pentagonus* (labeled *Australiensis*) also labeled *pentagonus* from Bailey, Australia—*Kalchbrennera corallocephala*, South Africa, collection and also *Kalchbrenner's* fine drawing¹⁴—*Simblum gracile*, Ceylon, also several specimens from Kurz, Java, and a nice drawing originally named "*Thyridocephalus flavescens, Mihi*"—*Simblum sphaerocephalum*, Brazil, Glazion (labeled *Simblum Brasiliense*, also drawing of the type of *S. pilidiatum*)—*Simblum periphragmoides*, type in good condition, Mauritius in Hooker's herbarium—*Clathrus pusillus*, type, Australia—*Pseudocolus rugulosus*, type drawing ex Kurz, Java, labeled *Clathrus triscapus*—*Pseudocolus Rothae*, two collections and sketch from Bailey—*Laterna columnata*, Brazil, also (very ???) from Australia, also Cuba, the latter more slender, also several from the United States—*Laterna pusilla*, type, Cooke—*Clathrus cancellatus*, England, France—*Clathrus pseudocrispus* type drawing ex McCarty, Jamaica, also poor specimens—*Clathrus crispatus* ex Thwaite, Ceylon (published as *cancellatus*)—*Clathrus cibarius*, about a dozen collections, all from New Zealand excepting one from Chiloe, an island off the coast of Chile—*Clathrus gracilis*, several, all from Australia—*Clathrus delicatus*, type, Ceylon—*Clathrus crispus*, Cuba, San Domingo, Uruguay—*Simblum clathratum*, type drawing and specimens, Old Calabar, Africa ex J. H. Holland—*Lysurus Gardneri*, abundant types, Ceylon—*Colus hirudinosus*, Corsica, Alpes Maritimes—*Lysurus Australiensis*, type, Australia—*Lysurus Woodii*, South Africa ex Wood, and same as co-type of *Anthurus Woodii*—*Aseroe Zeylandica*, type, Ceylon—*Aseroe rubra*, several from Australia—*Aseroe Hookeri*, type, New Zealand.

BRITISH MUSEUM, LONDON.

Phallus indusiatus from Philippines, Angola (Africa), China, Ceylon, India, Borneo, and St. Vincent (the latter a small form = *nana*)—*Phallus multicolor*, type with the original colored sketch and also a colored sketch by Broom—*Phallus impudicus*, several exsiccated from Europe, also specimens from Britain—*Phallus tenuis*, Ceylon—*Phallus quadricolor*, type—*Phallus calyptratus*, type—*Phallus Ravenelii* from Ravenel with his original notes—*Phallus auran-*

¹⁴ "The *Kalchbrennera* is very rare. Only twelve specimens have been found in five years."—Extract from letter from MacOwan.

tiacus, Australia—*Mutinus caninus*, photo ex. Krieger, also several specimens from England and the continent—*Mutinus elegans*, sketch from Morgan, labeled *Ravenelii*—*Mutinus Ravenelii* with a letter from Ravenel stating that the plant has been mis-cited in *Grevillea* (which is true)—*Mutinus proximus*, drawing by Broom—*Kalchbrennera corallocephala*, original description but no specimen from Welwitsch, drawing as reproduced in *Trans. Linn. Soc.*—*Simblum gracile*, Ceylon—*Laterna columnata*, abundant specimen from Ravenel—*Laterna Angolensis*, original description and drawing (no specimen) from Welwitsch. It was described as “*splendida albida*”—*Simblum Muelleri* (?) poor, Australia—*Clathrus cancellatus*, several specimens from Europe—*Clathrus delicatus*, co-type, Ceylon—*Clathrus gracile*, co-type, Australia—*Clathrus cibarius*, New Zealand, also Chiloe, Chile, also a form (see page 60) Pernambuco, also a specimen and sketch of a very similar species from Mombasa, East Africa. There are no color notes, but the sketch is dull yellow—*Clathrus crispus*, Vera Cruz, also specimen, poor, so labeled, from Australia, but not the species, I think—*Clathrus crispatus*, *Yucatan*, dried specimen, but seems the same as the original at Kew from Ceylon—*Lysurus Gardneri*, co-type, Ceylon—*Colus hirudinosus* (alcohol) from Meadow Valley, Asia Minor—*Aseroe rubra*, Australia.

CRYPTOGAMIC MUSEUM, PARIS, FRANCE.

Aseroe arachnoidea, type specimens in alcohol from Harmand, Cochinchina—*Anthurus trifidus*, in alcohol, type from Japan, Dr. Harmand, specimen is broken, but I think is a *Pseudocolus*—*Aseroe rubra*, New Zealand, Raoul—*Phallus impudicus*, several, France—*Phallus Ravenelii*. There is an historical specimen more than two hundred years old in the herbarium of Vaillant. It was “*Boletus phalloides*” and was “ex Canada, 1702”—*Phallus aurantiacus*, type, India—*Mutinus caninus*, several, France—*Mutinus elegans*, type, Ohio, good condition—*Mutinus Ravenelii* ex Ravenel—*Mutinus curtus*, fragment from Berkeley—*Mutinus xylogenus*, types ex French Guiana and drawing (good) from Leprieur—*Pseudocolus fusiformis*, type drawing from Reunion, all that is known—*Lysurus Gardneri* ex Berkeley—*Laterna triscapa* from Chile, the specimen is very small, but probably a small *columnata*—*Laterna columnata* ex Ravenel, also Chile—*Clathrus gracilis* ex Berkeley, also a specimen so referred from New Caledonia, the latter also published as *Colus hirudinosus*, but so poor it should not have been named—*Clathrus cancellatus*, a number all from Southern France and one Algeria—*Clathrus gracilis*, ex Berkeley, Australia, also a very poor specimen ex Africa, similar to *gracilis*, but too poor to judge. It was named *Clathrus Fischeri*—*Clathrus cibarius*, type ex New Zealand, in alcohol, also from Chile!—*Lysurus cruciatus*, a number of type specimens, but all much broken, French Guiana—*Colus hirudinosus*, specimen from Pyrenees, Algeria, and Corsica, the latter received by Montagne in 1820, thirteen years before it was published, and is labeled “*Clathrus hirudinosus nobis*”—*Phallus subuculatus*, type from Algeria, “very common and less fetid than *impudicus*,” says the collector—*Phallus duplicatus* ex Ravenel—*Phallus indusiatus*, in alcohol, from French Guiana, also dried, the type of “*Sophronia brasiliensis*” from Brazil, also type of “*Phallus radiatus*” from French Guiana, also from Tonkin and New Caledonia—*Anthurus Calathiscus*, no specimen was received, but the drawing is there from Perrottet, India, on which I think Montagne based his imaginary “*Calathiscus Sepia*.”

UPSALA, SWEDEN.

In alcohol.—There is a very abundant collection made by E. Nyman in Java a number of years ago.

Phallus indusiatus, twelve collections, ten of the usual form with broad pilei and two with slender pilei.

Clautriavia merulina, two collections. This is a frequent species in Java. *Simblum periphragmoides* (and the form *gracilis*), five collections, which convince me that *gracilis* is at the best a form of *periphragmoides*.

Mutinus bambusinus, one collection.

Clathrus Treubei, three collections, two old, with the arms broken apart, as shown in Myc. Notes, p. 382, fig. 212.

Jansia rugosa, one collection.

There is also at Upsala, in alcohol, a specimen of *Aseroe rubra* from New Zealand, collected by G. von Scheele; *Clathrus cancellatus* from Montpellier, France, and ten collections of *Phallus impudicus* by various collectors in Sweden.

Dried specimens.—*Aseroe rubra* from New Zealand, Berggren, and a drawing from the fresh specimens—*Clathrus cancellatus*, Tirol, Bresadola—*Clathrus pusillus*, "New Holland, ex. Berk."—*Mutinus elegans* from Curtis, and labeled "Corynites brevis," which was a manuscript name for it—*Clathrus cibarius*, New Zealand, Berggren—*Lysurus Gardneri*, co-types, ex. Berkeley—*Macowanites agaricinus*, co-type from Kalchbrenner. (Not usually classed in the phalloids, but to my mind closely related)—*Mutinus caninus*, ex. Quélet, France—*Mutinus* (unnamed), Guadeloupe, L'Herminier. (Something curious but unnamed, and I think this specimen unnamable)—*Phallus impudicus* Fautrey, France—*Clathrus guttulatus*, no specimen but the type drawing from Örsted on which the species was based.

BERLIN, GERMANY.

Dried Specimens: *Clathrus cancellatus*, from a hothouse at Berlin. It probably does not occur in the open as far north as Berlin. Three collections from southern Europe—*Clathrus gracilis*, three from Australia, also so-labeled from Samoa (!) and it seems correct.—*Clathrus Baumeri*, the types, dried specimens but better unnamed from such material.—*Clathrus crispus*, Guadeloupe—*Simblum sphaerocephalum*, three from Brazil and Uruguay—"Anthurus" Woodii, co-types and the type drawing. I think it is a *Lysurus*—*Lysurus borealis* var. *Klitzingii*, same exactly I think as our American form—*Aseroe* (sp.?) from Africa!—*Aseroe pallida*, type and drawing from New Caledonia—*Phallus indusiatus*, specimens from Samoa, Usambara, New Guinea, Brazil and Australia—*Phallus rubicundus*, specimens from South Africa and Australia (labeled *aurantiacus*)—"Omphalophallus" *calvescens* and *Muellerianus*, both co-types (Australia) and both same, but specimens too poor for comment, much less to be named—*Phallus* (unnamed) from Brazil, on the order of *Ravenelii* but much too large—*Mutinus caninus* seven collections from Germany—*Mutinus elegans* from Rau, Penn.—*Mutinus* (sp.?) from Dr. Reinecke, Samoa. I found no *Mutinus* in Samoa.—*Phallus impudicus*, many specimens, mostly from Germany—*Kalchbrennera corallocephala* from MacOwan, South Africa, also a drawing (labeled *Aseroe Tuckii* and the type of this "new species")—*Floccomutinus Zenkeri*, the original drawing from Zenker.

In alcohol: *Clautriavia merulina*, Java—*Simblum periphragmoides*, Java—*Phallus indusiatus*, nine from Java, two from Africa, three from New Guinea. One of the African forms has unusually large meshes to the veil—*Clathrus camerunensis*, type, Africa—*Clathrus Americanus* (unnamed) from Paraguay—*Clathrus chrysomycelinus*, type Brazil—*Laternea columnata*, Brazil—"Laternea pentactina", type, Java. It is an old condition of *Clathrus Treubei*, the arms broken apart as shown in figure 212, page 382, Myc. Notes—*Simblum sphaerocephalum*, Brazil—*Aseroe rubra*, New Guinea. I can not say as to the exact form, but it seems to have a broad limb and to tend towards the East Indian species.—*Lysurus borealis*, var. *Klitzingii*, Berlin, same I think as our American form—*Clautriavia Lauterbachii* (type of *Echinophallus Lauterbachii*) unfortunately only known from eggs as it is a most peculiar genus—Eggs of a *Clathrus* determined as *Preussii*? but I think the species is?—*Mutinus boninensis*, type, all known, intermediate between *Mutinus* and *Jansia*—*Jansia elegans*, abundant, from Java, type of "*Floccomutinus Nymanianus*" but quite different from the genus *Floccomutinus*, I think.—*Floccomutinus Zenkeri*, type Africa, a very distinct genus, in my opinion—*Mutinus bambusianus*, several collections from Java. Some have rather short heads and in size approach *Mutinus caninus* of Europe. The gleba-bearing portions are more pointed and not so even as *caninus*—*Mutinus caninus*, Germany—*Blumenavia rhacodes*, type, Brazil—*Itajahya galericulata*, type Brazil.—*Phallus glutinoides*, type, Brazil. All are in egg state.—

Clathrus pseudocancellatus, type, Africa. It was probably originally in formalin as it is now flabby and shapeless.

Phalloids in alcohol in the show department of the Museum at Berlin: At Berlin there is the finest collection of phalloids, both as to numbers and condition, that exists anywhere. It was the work of the late Dr. Hennings, and the specimens are most beautifully prepared and displayed. The following is the list: *Laterna columnata*, Brazil—*Clathrus cancellatus*, Europe—*Colus hirudinosus*, Sardinia—*Clathrus gracilis*, Australia—*Mutinus Moelleri*, type, Brazil (=for me, *bambusinus*)—*Floccomutinus Zenkeri*, type, Africa—*Jansia elegans* (type of *Floccomutinus Nymaniensis*)—*Phallus rubicundus*, Africa (type of *Phallus sanguineus*)—*Phallus tenuis*, Java—*Clautriavia Lauterbachii* (type of *Echinophallus*)—*Itajahya galericulata*, type, Brazil—*Phallus impudicus*, two from Java, one Africa, also a slender form from Java (type of *echinata*)—*Clautriavia merulina*, two, Java—*Lysurus borealis* (var. *Klitzingii*)—*Aseroe rubra*, New Caledonia, also New Guinea—*Blumenavia usambariensis*, type, Africa—*Clathrus Preussii*, type, Africa—*Simblum sphaerocephalum*, Argentina—*Simblum periphragmoides*, Java—*Mutinus caninus*, Berlin—*Mutinus Fleischeri*, type, Java—*Mutinus bambusinus*, two, Java—*Phallus celebicus*, type, Celebes—*Phallus impudicus*, three, Berlin—*Kalchbrennera corallocephala*, South Africa (labeled *Kalchbrenneri Tuckii* var. *clathroides*, Henn.)—*Clathrus camerunensis*, type, Africa—*Blumenavia rhacodes*, type, Brazil.

THE LLOYD MUSEUM, CINCINNATI, OHIO.

Note.—As this list is made in England from the published records without having access to the specimens, some may have been overlooked. All listed are dried specimens unless otherwise noted.

Europe.

Clathrus cancellatus, Portugal, Rev. Torrend.
Clathrus cancellatus, Italy, M. Bezzi.
Clathrus cancellatus, France, L. Rolland.
Clathrus cancellatus, Spain, T. de Aranzadi.
Clathrus cancellatus, France, Auguste Bernin (fresh!) (alcohol).
Mutinus caninus, Ireland, Greenwood Pim.
Mutinus caninus, Germany, C. Engelke.
Mutinus caninus, Germany, Otto Jaap.
Mutinus caninus, France, C. G. Lloyd.
Mutinus caninus, Germany, W. Krieger, photograph.
Phallus imperialis, Italy, M. Bezzi.
Phallus impudicus, France, C. G. Lloyd.
Phallus impudicus, France, L. Rolland.
Phallus impudicus, Italy, M. Bezzi.
Lysurus borealis (red arms), England, Harold Murray, photograph.
Colus hirudinosus, Portugal, Rev. Torrend (alcohol).

United States.

Laterna columnata, Florida, L. N. Fowler (alcohol).
Laterna columnata, Florida, Dr. J. F. Maddox (alcohol).
Laterna columnata, Florida, C. E. Pleas.
Laterna columnata, Florida, C. G. Lloyd (alcohol).
Laterna columnata, Florida, C. E. Pleas (photograph).
Mutinus caninus, Canada, Jas. Fletcher.
Mutinus caninus, Maryland, W. T. Lakin.
Mutinus caninus, New Jersey, E. B. Sterling (alcohol).
Mutinus elegans, Ohio, Pennsylvania, and Kentucky, C. G. Lloyd (alcohol).
Mutinus elegans, Pennsylvania, Dr. Herbst (alcohol).
Mutinus elegans, Ohio, M. E. Hard (photograph).
Mutinus elegans, Connecticut, C. C. Hammer (eggs).
Mutinus Ravenelii (?), New Jersey, E. B. Sterling.
Mutinus Ravenelii, Ohio, Chas. Dury (alcohol).

Mutinus Ravenelii (?), Florida, G. C. Fisher.
Mutinus Ravenelii, Ohio, A. P. Morgan.
Lysurus borealis, Ohio, H. C. Beardslee.
Lysurus borealis, Massachusetts, Geo. B. Fessenden.
Lysurus borealis, Connecticut, C. C. Hanmer (photograph).
Lysurus borealis, Massachusetts, Miss L. C. Allen.
Lysurus borealis (? red), Texas, W. H. Long, Jr.
Lysurus borealis, Massachusetts, G. E. Stone (alcohol).
Phallogaster saccatus, Ohio, C. G. Lloyd (alcohol).
Phallogaster saccatus, West Virginia, C. G. Lloyd.
Phallus duplicatus, Ohio, H. C. Beardslee (alcohol).
Phallus duplicatus, California, L. A. Greata.
Phallus duplicatus, Iowa, L. R. Waldron.
Phallus duplicatus, Ohio, C. G. Lloyd (alcohol).
Phallus duplicatus, Florida, G. C. Fisher.
Phallus duplicatus, Ohio, A. P. Morgan (alcohol).
Phallus duplicatus, Ohio, Prof. W. H. Aiken (alcohol).
Phallus imperialis, Colorado, E. B. Sterling.
Phallus imperialis, Texas, W. H. Long, Jr.
Phallus imperialis, California, L. G. Yates.
Phallus imperialis, Washington, D. C., F. J. Braendle.
Phallus imperialis, California, L. A. Greata.
Phallus imperialis, California, W. H. Henderson.
Phallus imperialis, Colorado, E. Bethel.
Phallus gracilis, Florida, L. N. Fowler (alcohol).
Phallus Ravenelii, Pennsylvania, Wm. Herbst (alcohol).
Phallus Ravenelii, Iowa, F. J. Fitzpatrick.
Phallus Ravenelii, Ohio, C. G. Lloyd.
Phallus Ravenelii, Ohio, Mrs. Carl Langenbeck (alcohol).
Phallus Ravenelii, Ohio, M. E. Hard (photograph).
Phallus Ravenelii, New Jersey, E. B. Sterling (fresh) (alcohol).
Phallus Ravenelii, Florida, G. C. Fisher.
Phallus rubicundus, Texas, W. H. Long, Jr.
Simblum sphaerocephalum, Nebraska, Rev. J. M. Bates.
Simblum sphaerocephalum, Texas, W. H. Long, Jr.
Simblum Texense, Texas, W. H. Long, Jr.

Brazil.

Itajahya galericulata, Rev. A. Schupp (photograph).
Clathrus chrysomycelinus, Rev. A. Schupp.
Clathrus Americanus, Rev. A. Schupp (dried).
Clathrus Americanus, Rev. A. Schupp (photograph).
Laterna rhacodes, Rev. A. Schupp.
Laterna rhacodes, Rev. A. Schupp (photograph).
Laterna (cfr. *columnatus*), Rev. J. Rick.
Laterna (unnamed?), Rev. J. Rick.
Phallus (labeled *rugulosus*), Rev. J. Rick.
Simblum sphaerocephalum, Rev. J. Rick.

West Indies.

Clathrus crispus, Jamaica, Miss Barrett.
Clathrus crispus, Jamaica, Wm. Chadwick.
Clathrus (sp.?), Bahamas, L. J. K. Brace (egg).
Clathrus Americanus, L. J. K. Brace (formalin).
Laterna pusilla (?), Jamaica, W. Jekyll.
Phallus indusiatus, Jamaica, H. E. Cox.
Phallus indusiatus, Jamaica, Miss Barrett.
Simblum sphaerocephalum, Bahamas, L. J. K. Brace.

Samoa.

Phallus indusiatus, C. G. Lloyd (photograph and dried).

Hawaii.

Phallus rubicundus, D. D. Baldwin.
Phallus rubicundus (form gracilis), N. A. Cobb.

Australia and New Zealand.

Aseroe Hookeri, Miss Jessie Dunn.
Aseroe Muelleriana, A. G. Hamilton.
Anthurus aseroeformis, Prof. McAlpine (photograph and description).
Clathrus cibarius, Robert Brown.
Clathrus cibarius, Miss Jessie Dunn.
Clathrus cibarius, S. Duncan.
Clathrus cibarius, W. H. Laing.
Clathrus cibarius, Rev. J. Wilson.
Clathrus gracilis, Prof. D. McAlpine.
Clathrus gracilis, J. T. Paul.
Clathrus gracilis, F. M. Reader.
Clathrus gracilis, J. H. Spencer (alcohol).
Clathrus gracilis, Margaret Flockton (alcohol).
Clathrus gracilis, Edmund Jarvis.
Clathrus gracilis, J. G. O. Tepper.

Africa.

Kalchbrennera corallocephala, Cape, J. M. Wood.
Clathrus (undetermined), Dr. Labesse (alcohol).
Clathrus (unnamed), Congo, Edouard Luja.
Phallus indusiatus, Congo, Edouard Luja (dried).
Phallus indusiatus, Congo, Edouard Luja (photograph).
Phallus indusiatus, C. B. Ussher.
Unnamed genus, C. B. Ussher.

Mauritius.

Phallus gracilis, Chas. A. O'Connor (alcohol).
Phallus indusiatus, Chas. A. O'Connor (alcohol).
Phallus Mauritianus, Chas. A. O'Connor (alcohol).

India.

Genus unnamed, G. H. Krumbiegel.

Japan.

Phallus indusiatus, Professor Gono (drawing).
Phallus impudicus, Professor Kusano (drawing).
Phallus rugulosus, Professor Kusano (drawing).
Phallus rugulosus, Professor Kusano (alcohol).
Phallus rugulosus, T. Nishida (alcohol).
Phallus tenuis, Professor Kusano (drawing).
Phallus tenuis, Professor Kusano (alcohol).
Mutinus boninensis (?), Professor Kusano (alcohol).
Mutinus boninensis (?), Professor Kusano (alcohol).
Lysurus Mokusin, Professor Gono (drawing).
Lysurus Mokusin, Professor Kusano (drawing).
Laternea bicolonnata, Professor Kusano (photograph).

Java.

Clathrus Treubei, Dr. Ch. Bernard (photograph).
Clathrus Treubei, Dr. Ch. Bernard (alcohol).
Phallus indusiatus, Dr. Ch. Bernard (alcohol).
Clautriavia merulina, Professor Patouillard (photograph).
Clautriavia merulina, Dr. Ch. Bernard (alcohol).

Simblum gracile, Dr. Ch. Bernard (photograph).
Simblum gracile, Dr. Ch. Bernard (alcohol).
Jansia rugosa, Dr. Ch. Bernard (alcohol).
Aseroe arachnoidea, Dr. Ch. Bernard (alcohol).
Aseroe arachnoidea, Dr. J. P. Lotsy (alcohol).
Mutinus bambusinus, Dr. Ch. Bernard (alcohol).

APPENDIX V.

SOURCE OF ILLUSTRATIONS.

Photographs.

The best illustration of a phalloid is a good photograph, and we confidently look to photography to dispel much of the doubt that surrounds many of the species of foreign phalloids. We present herewith a list of those who have published or supplied photographs of phalloids or furnished material to illustrate phalloids by photography, and have indicated our figures that are taken from these sources. America leads the world in the use of photography to illustrate fungi. Well illustrated books have appeared by Atkinson, McIlvaine, Hard, and Marshall, all containing illustrations of phalloids. We have not cited them in detail, however, as they all cover the same restricted field of a few species. We think the following is otherwise a complete list of those who have aided in the work. We hope this pamphlet will awaken interest in the subject in other countries and that the next résumé of the subject will have a much larger list. If you find a phalloid that is not illustrated in this work by a good photograph, we hope you will not fail to secure a good photograph of it, if possible.

Dr. Chas. Bernard, Java.

Aseroe Zeylandica (Fig. 54).
Clautriavia merulina
Jansia rugosa (Figs. 30 and 31).
Clathrus Treubei (Fig. 72).
Simblum gracile (Fig. 84).

Auguste Bernin, Monaco.

Clathrus cancellatus (Fig. 70).

N. A. Cobb, Hawaii.

Phallus gracilis (Fig. 6).

Robt. E. Fries.

Itajahya galericulata.

C. C. Hanmer, Connecticut.

Lysurus borealis (Fig. 41).

M. E. Hard, Ohio.

Phallus Ravenelii, (Fig. 8).

W. Krieger, Germany.

Mutinus caninus (Fig. 23).

Professor Kusano, Japan.

Laternea bicolonnata (Fig. 64).

W. H. Long, Jr., Texas.

Phallus rubicundus (Fig. 5).
Simblum Texense (Fig. 85).
Simblum sphaerocephalum (Fig. 86).

D. McAlpine, Australia.

Anthurus aseroeformis (Fig. 46).

Alfred Moeller (from Brazil).

Phallus indusiatus.
Phallus Moelleri (Fig. 13).
Clathrus chrysomycelinus (Fig. 80).
Laternea columnata.
Laternea rhacodes.
Phallus glutinolens (Fig. 10).
Pseudocolus Garciae (Fig. 65).
Mutinus bambusinus (Fig. 26).
Itajahya galericulata (Fig. 22).

Harold Murray, England.

Lysurus borealis.

Chas. O'Connor, Mauritius.

Phallus Mauritanus (Fig. 17).

N. Patouillard, Paris

Clautriavia merulina.

Otto Penzig (from Java).

Aseroe arachnoidea (Figs. 55 and 56).
Phallus indusiatus.
Phallus favosus (Fig. 4).
Clautriavia merulina.
Phallus multicolor (Fig. 14).
Simblum gracile.
Jansia elegans (Figs. 32 and 33).
Jansia rugosa.
Mutinus Fleischeri (Fig. 27).
Pseudocolus Javanicus (Fig. 66).

T. Petch, Ceylon.

Jansia rugosa.
Phallus indusiatus.
Clautriavia merulina (Fig. 19).
Clathrus crispatus (Fig. 74a).
Clathrus delicatus (Fig. 82).
Simblum gracile.
Lysurus Gardneri (Fig. 38).

C. E. Pleas, Florida.

Phallus duplicatus (Fig. 16).
Laternea columnata (Figs. 57 and 58).

Carleton Rea, England.

Lysurus borealis (Fig. 40).

Rev. J. Rick, Brazil.

Laternea rhacodes (Fig. 61).
Simblum sphaerocephalum.

Rev. A. Schupp, Brazil.

Itajahya galericulata (Fig. 21).
Clathrus Americanus (Fig. 71).

Rev. J. Torrend, Portugal.

Colus hirudinosus (Figs. 89 and 90).
Simblum sphaerocephalum.

Photographed by the writer.

Phallus impudicus (France) (Fig. 1).
Phallus Ravenelii, Ohio (Fig. 7).
Phallus indusiatus, Samoa (Fig. 12).
Mutinus elegans, West Virginia (Fig. 24).
Mutinus Ravenelii, Ohio (Fig. 25).
Phallogaster saccatus, Ohio (Figs. 93 and 94).

Figures.

Next to a photograph, an accurate drawing is the best illustration. In a few instances we have used Penzig's figures in preference to his photographs. I think all figures are not good, especially the old ones reconstructed from dried specimens. However, as to many species they are all we have, and the following are the sources from which they have been reproduced. There are phalloids unillustrated by even a crude drawing. It is a standing reproach that authors are found to engage in such work. In a few such cases we have photographed the type as a makeshift illustration, but the most of such work we think is better considered as "nomina nuda" and relegated to "synonymy."

- | | | | |
|----------|--------------------------------|----------|---------------------------------|
| Fig. 2 | Drawing by Otto Penzig. | Fig. 62 | Drawing by Welwitsch. |
| Fig. 3 | Drawing by Ed Fischer. | Fig. 63 | Drawing by Spegazzini. |
| Fig. 9 | Drawing by Ed. Fischer. | Fig. 67 | Drawing by Kurz. |
| Fig. 11 | Drawing by Alfred Moeller. | Fig. 68 | Drawing at Paris. |
| Fig. 15 | Drawing by Rumphius. | Fig. 69 | Drawing by F. M. Bailey. |
| Fig. 18 | Drawing by M. C. Cooke. | Fig. 73 | Drawing by Berkeley. |
| Fig. 20 | Drawing by Ed. Fischer. | Fig. 74 | Drawing by Ed Fischer. |
| Fig. 28 | Drawing by F. M. Bailey. | Fig. 75 | Drawing by Oersted. |
| Fig. 29 | Photographed from the type. | Fig. 76 | Drawing by Turpin. |
| Fig. 34 | Drawing by F. M. Bailey. | Fig. 77 | Drawing by Dr. McCatty. |
| Fig. 35 | Drawing by Ed Fischer. | Fig. 78 | Photograph, Museum at Kew. |
| Fig. 36 | Drawing by Ed Fischer. | Fig. 79 | Photograph from alc. specimens. |
| Fig. 36a | Drawing by Zenker. | Fig. 81 | Drawing by Ed Fischer. |
| Fig. 37 | Photograph from alc. material. | Fig. 83 | Drawing by Hooker. |
| Fig. 38a | Photograph from dried type. | Fig. 87 | Drawing by Ed Fischer. |
| Fig. 39 | Photograph from dried type. | Fig. 88 | Drawing by J. W. Holland. |
| Fig. 42 | Drawing by Mueller. | Fig. 91 | Drawing by Kalchbrenner. |
| Fig. 43 | Drawing by Ed Fischer. | Fig. 92 | Drawing by Kalchbrenner. |
| Fig. 44 | Drawing by Montagne. | Fig. 95 | Drawing by Kalchbrenner. |
| Fig. 45 | Photographed from the type. | Fig. 96 | Drawing by Berkeley. |
| Fig. 47 | Drawing by Kalchbrenner. | Fig. 98 | Drawing by Berkeley. |
| Fig. 48 | Drawing by Berkeley. | Fig. 99 | Drawing by Hennings. |
| Fig. 49 | Sketch by Perrottet. | Fig. 100 | Drawing by Montagne. |
| Fig. 50 | Drawing by La Billardiére. | Fig. 102 | Drawing by Berkeley. |
| Fig. 51 | Drawing by Berkeley. | Fig. 103 | Drawing by Patouillard. |
| Fig. 52 | Drawing by Berkeley. | Fig. 104 | Drawing by Patouillard. |
| Fig. 53 | Drawing by Kalchbrenner. | Fig. 105 | Drawing by Corda. |
| Fig. 53a | Drawing by Le Rat. | Fig. 106 | Drawing by Kalchbrenner. |
| Fig. 59 | Drawing by Turpin. | Fig. 107 | Drawing by Spegazzini. |
| Fig. 60 | Photograph of the type. | | |

APPENDIX VI.

Reproduction of the original figures of doubtful and little known species of the genera *Phallus* and *Mutinus*. Most of them I think have no value whatever, but there is no way of getting rid of them.



Fig. 95.

PHALLUS DISCOLOR.



Fig. 99.

PHALLUS CELEBICUS.



Fig. 102.

PHALLUS QUADRICOLOR.



Fig. 98.

PHALLUS CAMPANULATUS.



Fig. 96.

PHALLUS CALYPT-
TRATUS.



Fig. 106

MUTINUS PA-
PUISIUS.



Fig. 107.

MUTINUS ARGENTINUS.



Fig. 105.
MUTINUS CURTUS.



Fig. 100.
PHALLUS CA-
NARIENSIS.




Fig. 103.
MUTINUS MINIMUS.



Fig. 104.
MUTINUS BORNE-
ENSIS.

APPENDIX VII.

ASEROË RUBRA LA BILL. VAR. JUNGHUHNII SCHLECHT.

PAR DR. CH. BERNARD.

Avec deux photographies.

Il y a peu de temps, j'ai publié dans les Annales du Jardin botanique de Buitenzorg (Vol. XXII, 1908. 2^{ème} partie, pp. 224-238), une petite note sur cette Phalloïdée très curieuse, assez rare, et jusqu'alors assez mal connue. Je déplorais à cette occasion de n'avoir pas pu faire une photographie convenable de ce type, et je signalais certains points dont l'étude demandait à être reprise ultérieurement. Depuis lors, j'obtins, toujours grâce à l'extrême amabilité de M. le Dr. *J. Bosscha*, plusieurs magnifiques exemplaires de cet organisme, et entre autres les deux individus dont M. *Huysmans* a bien voulu faire les deux photographies qui accompagnent cette note. Tous ces échantillons provenaient de la plantation de Taloen, sur le plateau de Pengalengan, au Sud de Bandoeng, c'est à dire de la même station où avaient été récoltés les



exemplaires décrits dans ma précédente note. Ces nouveaux individus n'ont permis de faire certaines observations venant jeter quelque lumière sur des détails laissés jusqu' à présent dans l'ombre, et je ne crois pas inutile de publier ici ces quelques lignes qu' illustreront les deux photographies en question, et qui viendront fixer ou rectifier certains points d' importance secondaire, car je dois dire dès le début que, dans leurs caractères importants, ces individus coïncidaient très exactement avec ceux déjà observés.

Je disais entre autres: "*Aseroë rubra* est une espèce extrêmement polymorphe; . . . il est impossible de trouver dans la série de ses formes de "passage une solution de continuité permettant de séparer des espèces . . .

“et la forme qui nous occupe ici vient diminuer encore la valeur des variétés nettement délimitées. . . . Il importe donc de ne séparer les types qu’ avec la plus grande prudence, car il est probable que pendant bien longtemps, chaque fois qu’ on découvrira un exemplaire de ces champignons éminemment variables, ce nouvel individu constituera un anneau de cette longue chaîne de types voisins, atténuant les différences et supprimant telle ou telle variété.”

Les échantillons dont il sera question dans la présente note apporteront un argument de plus en faveur de cette opinion, et si, par certains de leurs caractères, ils se rapprochent des individus que j’ai décrits antérieurement, la disposition de leurs bras, qui est fort typique, établit un passage vers d’autres formes. Je me contenterai de donner une description de ces exemplaires, les points sur lesquels je veux fixer l’attention ressortiront d’eux mêmes de la description et de l’examen des photographies.



Les deux échantillons que j’ai pu faire photographier n’étaient pas de dimensions particulièrement considérables, et le nombre des bras était de 18 chez l’un et de 22 chez l’autre, ce qui correspond aux indications que j’ai données antérieurement. De même l’extrémité plus ou moins régulièrement enroulée des bras est caractéristique. La disposition de la volve, du pied assez court, de la glèbe, les couleurs, l’odeur, etc., ne distinguaient en rien ces formes de celles déjà décrites. Pour tous ces détails je renvoie donc à ma précédente

publication. Mais le point important et sur lequel je tiens à insister est le suivant: tandis que les exemplaires que j'ai observés jusqu'ici établissaient un passage entre les var, *zeylanica* et *Junghuhnii* de *Aseroë rubra*,—certains de leurs caractères, comme je l'ai démontré p. 235, rappelant ceux de l'une ou de l'autre de ces deux variétés,—les types dont nous nous occupons aujourd'hui et surtout l'un d'entre eux sont beaucoup plus voisins de la variété *Junghuhnii*; et même par la plupart de leurs détails, ils coïncident presque exactement avec cette variété comme nous la trouvons décrite d'ordinaire; cependant en comparant tous mes échantillons, j'ai pu me convaincre qu'il n'y a nulle raison de séparer les unes des autres ces différentes formes qui toutes du reste proviennent d'une seule et unique localité; elles appartiennent non seulement à la même espèce, mais aussi à la même variété, cela ne fait aucun doute et il me semble que la question que se posait *E. Fischer* dans le "Sylloge Fungorum," Vol. VII, p. 25, quand il se demandait à propos de *A. Junghuhnii*: "An ab *A. zeylanica* diversa?" doit être certainement résolue par le négative. Non seulement il ne saurait s'agir de deux espèces différentes, mais encore il me paraît que les deux types doivent être rangés sous un même nom de variété.

Le caractère important auquel je fais allusion est la disposition des bras; je disais à ce propos: "Les bras étant séparés les uns des autres et le disque, dans certains échantillons étant très peu développé, cela parle en faveur d'une identité avec *A. zeylanica*. Cependant il faut remarquer que le disque peut être remarquablement développé (caractère de *A. Junghuhnii*) puis que, si les bras sont le plus souvent nettement séparés les uns des autres jusqu'à leur base, il existe cependant des cas, assez rares, où l'on pourrait croire à de vagues indications de rapprochements par paires." Dans les formes qui nous occupent, il ne s'agit plus de vagues indications. Un des deux individus photographiés n'est pas encore très convaincant à cet égard, il est cependant facile de reconnaître que ses 18 bras sont rapprochés par paires les uns des autres. Mais l'autre individu est des plus typiques, et ses 22 bras sont très nettement et très régulièrement rapprochés deux par deux les uns des autres. Dans les deux échantillons le disque est assez fortement développé.

Pour terminer je crois qu'il m'est permis de maintenir, en la renforçant, la conclusion que j'énonçais à la fin de mon précédent travail, mais que je n'osais encore affirmer, à savoir que les variétés *zeylanica* et *Junghuhnii* d'*Aseroë rubra* devront être réunies dans la suite sous un même nom.

Cette petite note n'a pas d'autre prétention que de présenter deux individus de cet intéressant champignon qui, s'il a été souvent décrit, et plusieurs fois dessiné, n'avait pas encore, que je sache, été photographié jusqu'ici.

Les deux photographies reproduisent le champignon à peu près en grandeur naturelle.

EDITOR'S NOTE.

We publish the above article by Dr. Chas. Bernard, and we take no editorial liberties with it, but publish it just as received. We are particularly glad to get the photograph, which is the first published of this species of the East Indies. As we have stated in detail on page 44 our views as to the species of *Aseroe*, we shall not discuss the matter here. We believe, however, that the Ceylonese plant, *Aseroe Zeylandica*, and the Javanese plant, *Aseroe Junghuhnii*, are one and the same but quite distinct from the New Zealand and Australian forms which go to make up *Aseroe rubra*.—C. G. L.



THE LLOYD LIBRARY, CINCINNATI, O.

Devoted exclusively to a library of Botany and Pharmacy. It contains at the present time about twenty-five thousand volumes on the above named subjects.

INDEX.

In making up the pages it was not practicable to keep the figures in serial order nor always in close relationship to the corresponding text. In this index we have given the page on which will be found the text and also the figure for each species.

	Text on Page	Figure on Page		Text on Page	Figure on Page
Anthurus	40		Lysurus canctae Catherinae	40	38
" Archeri	43	42	" (unnamed)	40	39
" aseroeformis	42	41	" Woodii	40	38
" calathiscus	43	41	Mutinus	27	
" Muellerianus	43	42	" argentinus	32	90
Aseroe	43		" bambusinus	28	29
" arachnoidea	48	45	" borneensis	32	91
" Hookeri	46	45	" caninus	28	29
" lysuroides	48		" curtus	32	91
" Muelleriana	46	46	" elegans	28	29
" pallida	47	43	" Fleischeri	28	31
" pentactina	46	43	" minimus	32	91
" rubra	46	44	" papasius	32	90
" Zeylandica	47	47	" pentagonus	30	31
Clathrus	54		" proximus	32	
" affinis	60		" Ravenelii	28	30
" (African form)	62		" xylogenus	30	30
" Americanus	56	55	Phallogaster	71	
" camerunensis	57	55	" saccatus	71	71
" cancellatus	54	55	Phallus	9	
" Chrysomycelinus	63	62	" callichrous	20	
" cibarius	60	61	" calypttratus	22	90
" crispatus	57	57	" campanulatus	22	90
" crispus	58	58	" canariensis	22	91
" delicatus	63	63	" celebicus	22	89
" gracilis	62	60	" costatus	10	13
" guttulatus	58	58	" Daemontum	20	16
" Preussii	63	62	" discolor	22	89
" pseudocancellatus	60		" duplicatus	20	17
" pseudocrispus	59	59	" Farlowii	24	
" pusillus	57	56	" favulus	14	13
" Treubei	56	56	" gracilis	14	15
Clautriavia	24		" glutinolens	18	13
" Lauterbachii	24	26	" imperialis	10	
" merulina	24	25	" impudicus	10	11
Colus	67		" indusiatus	18	16
" hirudinosus	67	68	" Mauritianus	22	23
Floccomutinus	34		" Moelleri	20	19
" Zenkeri	35	34	" multicolor	20	21
Itajahya	26		" quadricolor	24	89
" galericulata	27	27	" Ravenelii	14	11
Jansia	33		" retusus	22	
" annulata	34	33	" Rochesterensis	20	23
" boninensis	34	32	" roseus	20	
" elegans	33	32	" rubicundus	14	12
" rugosa	33	32	" rugulosus	18	12
Kalchbrennera	68		" subuculatus	22	
" corallocephala	68	69	" subtilis	18	23
Laternea	48		" tenuis	10	12
" angolensis	50	49	Pseudocolus	51	
" bicolumnata	51	51	" fusiiformis	53	52
" columnnata	48	49	" Garciae	52	54
" pusilla	50	48	" javanicus	52	52
" rhacodes	50	49	" Rothae	53	53
" Spegazzini	51	50	" rugulosus	52	53
" triscapa	48	48	Simblum	64	
Lysurus	35		" clathratum	67	66
" australiensis	38	36	" gracile	66	65
" borcalis	38	36	" Mülleri	67	64
" Clarazianus	40	38	" periphragmoides	66	64
" cruciatus	40	39	" sphaerocephalum	67	65
" Gardneri	36	37	" Texense	67	65
" Mokusin	36	37			

ak
600
.L79

JUN 2 1920

BULLETIN No. 14. JUNE, 1910. MYCOLOGICAL SERIES, No. 5

BULLETIN
of the
LLOYD LIBRARY
of
BOTANY, PHARMACY AND
MATERIA MEDICA

J. U. & C. G. LLOYD
CINCINNATI, OHIO

MYCOLOGICAL SERIES, No. 5

Synopsis
of the
Genus Hexagona

By C. G. LLOYD



P. HARIOT.

To Monsieur P. Hariot, Curator of the Cryptogamic Museum at Paris, France,
I beg to dedicate this pamphlet, in recognition of the many courtesies
extended to me while working in the Museum.—C. G. LLOYD.

THE GENUS HEXAGONA.

Definition.—The genus *Hexagona* can be described in a few words as being fungi with large, round or hexagonal pores, or *Polyporus* with large, round or hexagonal pores. It is a tropical or sub-tropical genus, and does not occur in temperate regions. Theoretically the genus is purely artificial, based on a single character, but it is a very convenient genus, and to attempt to break it up would only make a needless complication.¹

In context *Hexagonas* are usually corky-woody, and as Fries knew only such species, he specified this as a character of the genus. There are a few species with hexagonal pores of which the context is rather fleshy-cartilaginous. We should include them also in *Hexagona*, believing it to be better to take the genus literally on its pore character.

Size of the pores.—While the basic idea of the genus is *large*, round or hexagonal pores, the word *large* is, of course, only relative. There has been associated (unintentionally perhaps) with the genus *Hexagona* the idea of regularity of pores. Thus there is a series of forms related to *Hexagona tenuis* which have by common consent always been classed in *Hexagona*. The pores are rather small, but they are *regular* and *shallow*. Many other polyporoids with larger pores are classed with *Polyporus* or *Trametes* when the pores are deep.

There is also a group of species, such as *Polystictus pinsitus*, with equally as large and as shallow pores, but they are irregular, with thin, angular, uneven pore-walls. This section is generally included in *Polystictus*, and I think that is preferable, but sometimes species of this section have been called *Hexagona*.

Spores.—The color of the spores was not taken into account by the old authors in the genus *Hexagona*, but as they are practically all supposed to have white spores, we should make that a character also of the genus. We have never seen a single spore of any species of what we include in *Hexagona*, as spores of white-spored polyporoids are *rarely found* in herbarium specimens. But the fact that we do not find the spores is a strong evidence that they are white.²

¹Any one who is so disposed, however, can discover as many and as useless "new genera" in it as have several times been discovered in *Polyporus*.

²It is well known that species of polyporoids with colored spores have the spores usually (if not always) in abundance. I have no doubt that all *Hexagonas* have white spores, though I have not positive evidence of the fact in a single instance.

There are but two cases where plants have been classed as *Hexagona* which I have found to have colored spores, viz.: *Hexagona decipiens*, of Australia, and *Hexagona gracilis*, of Brazil. I shall exclude them both, however, from *Hexagona* and include them in the section of *Polyporus*, where they belong, not only on their spore color, but their general natures, which are closer to other *Polyporus* than to *Hexagona*.



Fig. 276 X 6




Fig. 277.

Surface markings.—The next character that we use in grouping the species is the nature of the surface.

Setosus.—There is a very marked section of *Hexagona* with dense, coarse, rigid, black hairs or setae on the surface. This is the same character that is so familiar in the common *Trametes hydnoides* of the tropics. In fact, there is a series of species beginning with *Trametes hydnoides* with minute pores and ending with *Hexagona apiaria* with the large pores. This series has the same form, context, color, peculiar surface hairs, and differs chiefly in the size of the pores. It is a very natural group and might well be discovered to be a new genus. The hairs (Fig. 276 X6) are peculiar, being more or less wedge-

shaped and incised. They are detersive and fall away from old specimens.³ Our figure (277) shows the same species (of Australia) in three different conditions. But there are species that we include in this section in which these hairs are not so strongly developed. *Hexagona Deschampsii* (Fig. 282) has the hairs or fibrils strongly agglutinate with only a few free fibrils. It resembles old, worn conditions of the previous species. And finally we include in this section species strongly marked with entirely agglutinate fibrils as *Hexagona elegans* (Fig. 284).

Velutinus.—This section includes species with fine, soft, velutinate hairs (Fig. 289). It embraces but a few species.

Glaber.—Surface smooth, but often zoned or uneven, but not hirsute, velutinate or with strongly agglutinate fibrils.

General shape.—There are of course intermediate specimens, but most species can be arranged in one of three sections according to their general shape. All *Hexagonas* are sessile without stems,⁴ or (as in *albida*) sometimes a rudimentary-lateral stem.

Ungulaformis.—This section, which comprises but a very few species, has the pileus thick with deep pores. It is usually hard and sub-woody and corresponds to the genus *Fomes*, though the pores are never stratified.

Applanatus.—The texture is softer and general shape is flatter than in the previous section. The shape corresponds to the usual shape of *Fomes applanatus*. Usually it is a centimeter or more thick. We include in this section also some species such as *Kurzii* that are thinner and tend toward the next section but have deeper pores, 5 mm. or more.

Tenuis.—The character of the section *tenuis* is the very *thin* pileus, rarely over two or three mm. thick, and the small (for *Hexagona*) shallow, regular pores.

The setae in the pores.—There are some species of *Hexagona* that have conspicuous, colored setae in the pores. Sometimes these setae are large enough to be seen with the naked eye, and can easily be



Fig. 278 (X 6).

noted on the accompanying figure (278), which is enlarged six diameters. These setae are always in connection with ferruginous or cinnamon context. They are much larger than the microscopic setae found on many polyporoids, (Cfr. Pol. Issue, page 2), and often called cystidia. Klotzsch noticed these setae and proposed a new tribe,

Scenidium, but the idea never found favor. The inner surface of the

³ Mr. Murrill states that the plants are "nearly glabrous when young" and are finally clothed with these hairs. I think his explanation should be taken backwards.

⁴ This statement excludes *Hexagona gracilis*, which, however, is not a *Hexagona* for me.

pores of *Hexagona leprosa* appears finely pubescent, but under the lens the hairs are sub-hyaline and are of quite different nature from the colored setae of other species.

Color of context.—Most species have a colored context. It is hard for me to designate the exact color, though it is customary to describe it as cinnamon, ferruginous, gilvous, etc. In this pamphlet I designate these as having “colored context.” A few number have a context color white or pale ochraceous, much paler than the former. We indicate these as “pale context.” The difference between these two context colors is so marked that the character can be used to advantage in classification, and we base on it one of our groups.

The glaucescence of pores.—Many collections of *Hexagona* have pores strongly silvery glaucous, and it is a puzzling question how much stress to place on it in classification. In itself, I think it is not of much value, for many collections show some specimens glaucous and others not or only partially so. It seems to me to be a sort of deposit on the pores, with age perhaps. At Kew there is a specimen of *Hexagona apiaria* where most of the pores are strongly glaucous and the outer (younger) ones not at all. It is one of the species that has setae on the pores. The setae are quite noticeable on the non-glaucous pores, but in the glaucous ones they are not visible and have been covered up (apparently) with this deposit.

History of the genus.—In the very old days all fungi that had pores were called *Boletus*, and under this name are included in Linnaeus' *Species Plantarum*. Palisot-de-Beauvois, who was a collector of African plants, included a few fungi in his plates, and the Polypores he divided into two genera, those with large, round, hexagonal, or elongated pores, that he called *Favolus*, and those with small, round pores that he called *Microporus*. Fries divided Palisot's first genus into two, those with hexagonal pores which he called *Hexagona*, and those with elongated pores that he called *Favolus*. He took the name *Hexagona* from a probably inaccurate illustration of Pollini, which however showed hexagonal pores, and the genus was based on this one character. It has been taken in this sense by mycologists for about eighty years and about a hundred alleged new species have been named in accordance. The early mycologists, Persoon, Klotzsch, and Berkeley, were at first not disposed to consider a “large pored” genus of much value, but after the appearance of Fries' *Epicrisis* (1838), where he collated all the known species, no one has presumed to deny the genus.⁵ The first species to reach Europe was undoubtedly sent to Linnaeus, who named it *Boletus favus*. It was from China, but is not in the Linnaean herbarium, but was stated by Klotzsch, Berkeley, and Hariot to be the same as *Hexagona Wightii*, which it probably is.⁶ The next were two species from Africa, beautifully illustrated by Palisot-de-Beauvois, and the specimens (one at least) are

⁵ Recently a little cheap juggling was attempted to change all *Hexagonas* to *Favolus* and all *Favolus* to *Hexagona*, thus making new combinations for them all, and a muss in general. It was based on such a flimsy pretext that it is not worth discussing in detail. Monsieur Hariot, who first showed how the trick could be turned (*Bull. Soc. Myc. de France*, 1891. p. 203), dismissed it with the very sensible remark: “Mais la tradition Friesienne s'est imposée et il serait difficile dans l'état actuel de la science d'intervir les désignations générales de plus d'une centaine d'espèces. Le remède deviendrait pire que le mal et force est de s'en tenir aux idées admises.”

⁶ How this was overlooked in the priority hunt I do not know. 1753 is such a “prior” date to arrange alleged synonyms that a good thing like that ought not to be overlooked.

now in the Delessert Herbarium at Geneva.⁷ Next, *Hexagona tenuis*, the most common species of the tropics was named by Hooker from South America. The specimen is in fair condition in Hooker's herbarium at Kew. Then Persoon published *Hexagona apiaria* and *Hexagona vespacea* from the island of Rawak, and both are in good condition at Paris, the latter rather scanty however. When Fries issued his *Epicrisis* (1838) twelve species were supposed to be known and Fries had overlooked one of Palisot's.⁸ Since the *Epicrisis* there has been a steady output of "new species." There are in my index 125 names of supposed species, but this includes a few that were not called *Hexagona* but which in my opinion should have been, and a larger number that were called *Hexagona* and should not have been.

The work in this pamphlet.—In my work with the polyporoids I have visited and studied and photographed the species of *Hexagona* in the following museums: Kew London, British Museum London, Leiden, Berlin, Upsala, and Paris, and the private herbarium of Monsieur Patouillard at Paris. Most of the historic material is preserved in these museums and of the 125 species named in my index I have seen the type (or in a few cases the co-type) of all that exist but eight.

Classification.—For convenience we arrange the species in sections or groups. We have tried to make these groups as natural as possible for the benefit of the future "new genus" discoverers.

First (*Setosus*).—Typically the surface is clothed with dense, coarse, branched, rigid hairs, usually detersive and then the surface is fibrillose. We include also species which are covered with agglutinate fibrils generally with free ends.

Second (*Velutinus*).—Surface covered with fine, soft pubescence, or velutinate hairs. Plants that we include here are all thin and as to shape belong in section 5 (*Tenuis*). A few species that are pubescent we arrange on general relations in other groups, viz.: *Pobeguini* in *Applanatus*; *macrotrema* in *Pallidus*; *bipindiensis* in *Pseudofavolus*.

Third (*Ungulaformis*).—Thick, hard, with long pores and generally hoof-shaped. This section corresponds to the woody (*Fomes*) section of *Trametes* and might be called *Hexagona-Fomes*. We include here also *Hexagona resinosis* on its general nature, although it has a pale context and might be placed in the group *Pallidus*.

Fourth (*Applanatus*).—General shape applanate or flattened as distinguished from *Ungulaform* or hoof-shaped. Plants of this section are of a softer nature than the previous section. We include here for this reason *Hexagona amplexens*, that from its shape alone should be included in the preceding, and for a similar reason we include in the preceding *resinosa* which from its shape alone belongs in this section.

Fifth (*Tenuis*).—Pileus very thin, rarely two or three mm. thick with small (for *Hexagona*) regular, shallow pores. Surface smooth. We include in *Velutinus* species with pubescent surface.

Sixth (*Pallidus*).—Context white or pale ochraceous. (All the preceding except *resinosis* have context that is more deeply colored, ferruginous, cinnamon, etc.)

Seventh (*Pseudofavolus*).—All the previous have context suberose or subligneous. In this section we include the species with fleshy, tough nature. Usually they have been classified with the genus *Favolus* from which they differ essentially in their pore shapes. Patouillard bases on them a genus (*Pseudofavolus*). In our opinion, they should be included in *Hexagona*.

⁷I have never seen these, but Palisot gave such fine illustrations that it is not necessary to see them. One can be as sure from such illustrations as from the specimen itself.

⁸Viz: *Hexagona glabra*. As Saccardo seems to have started his compilation of the polyporoids with Fries's *Epicrisis*, this species does not occur and has been lost to all modern books.

Eighth (Resupinatus).—I think no resupinate species occur, but the section is convenient for a resupinate "species" so claimed.

GROUP 1, SETOSUS.

HEXAGONA APIARIA (Fig. 279).—Color dark. Surface densely covered with coarse, branched, dark hairs which are deterrentive,




Fig. 279

Hexagona apiaria. Type at Paris.

and old specimens have the surface coarsely fibrillose. Pores large (3-4 to cm.) from 5-10 mm. deep, ferruginous, often glaucous *with prominent setae* (cfr. page 3, fig. 278 x6.) Context thin, ferruginous.

History.—A frequent plant in the Philippines, India, Ceylon, and Australia, and as found in the museums usually called *Hexagona Wightii*. At Berlin there is a specimen from New Guinea and one from Guadeloupe.⁹ Linnaeus named something (none knew what) *Boletus favus* which came from China, and was evidently a *Hexagona* and probably this plant.¹⁰ The first specimen

⁹ This is the only specimen from an American station, but the plant is not included in N. A. F., which professes to include all West Indian species. Owing to the superficial work done by the author in the museums of Europe, he probably never saw the specimen.

¹⁰ The specimen does not exist in the Linnaean herbarium, though there is in the herbarium a specimen of *Hexagona tenuis* named "*Boletus favus*, Linn." by Dickson, many years after Linnaeus died.

known to reach Europe was from Rawak and was named *Polyporus*¹¹ *apiarius* by Persoon and a good figure given. The specimen (Fig. 279) is in good condition at Paris. Next Klotzsch got a specimen from Wight, India, which he called *Polyporus Wightii*, and also gave a good figure of it.¹² He noticed the setae (See fig. 278, page 3) in the tubes, which are evident even to the naked eye, and gave an exaggerated figure of them and based on them a "new tribe," *Scenidium*.¹³ A number of specimens have since reached Europe and are usually referred to *Wightii*. At the time he described the plant Klotzsch published that it was the same as *Boletus favus* of Linnaeus, and that was also Berkeley's opinion, and I think was probably true.¹⁴




Fig. 280
Hexagona hirta.

HEXAGONA HIRTA (Fig. 280).—Color dark. Surface covered with a dense coat of rigid, branched, dark hairs. These are often detersive. Context dark, ferruginous. Pores medium (about 8 to 10 to cm.) about 5 cm. deep. Owing to the depth and relatively small size of the pores it is often put in *Trametes*, and it belongs there about as well as in *Hexagona*.

History.—It seems to be a common plant in Africa, but only in Africa as far as I know. It was most beautifully and accurately illustrated by Palisot-de-Beauvois more than a hundred years ago (1805) and his specimen is at Geneva. Notwithstanding it frequently reaches Europe, but one single specimen has ever been referred to Palisot's name. Fries discovered it was a new species

¹¹ Persoon at the time was aware of the genus *Hexagona*, but declined to consider it a genus, stating that the size of pores is only a relative character.

¹² I think the type does not exist. The only specimen I have seen from India is at the British Museum, but was not collected by Wight.

¹³ Under these conditions it seems to me very careless, to say the least, for Mr. Murrill to describe the pores of *Hexagona Wightii* as "glabrous within."

¹⁴ Klotzsch does not seem to have been consistent in his views of "*Boletus favus*, Linn." He gives this plant as being the same, and then he refers another plant to *Hexagona sinensis*, which was only a name-change of "*Boletus favus*."

and called it *Hexagona crinigera*. Klotzsch got it from Mauritius and referred it to Linnaeus' (alleged) species under a Friesian name-change, *Hexagona sinensis*. Berkeley decided it was not the Linnaean species and changed it to *Trametes Klotzschii*. (He was only guessing, but probably guessed right.) I think this name has been most generally used for it. Then L veill  got a specimen from Madagascar, and found it to be another new species, *Trametes crassa*. Then Cooke got the *same collection* (Perville, Madagascar) and described it as *Trametes adelphica*, but he does not seem to have taken himself very seriously, for he never changed his manuscript name on his specimens and they are found to-day in his collection as *Hexagona strigosa*.




Fig. 281
Hexagona capillacea.

HEXAGONA CAPILLACEA (Fig. 281.) — Color light, ferruginous or cinnamon, covered with a dense coat of concolorous hairs. Pores large, 3-4 mm. deep, with thin, flaccid walls. Bright ferruginous in color, devoid of setae.

History.—This is known from a single specimen (Fig. 281) from Venezuela, South America, now in the herbarium of Patouillard. From the figure it is evidently close to *apiaria*, but is lighter color, has finer hairs and thinner, more flaccid pore-walls.



Fig. 282
Hexagona Deschampsii.

HEXAGONA DESCHAMPSII (Fig. 282.)—Pileus dark reddish brown, with adpressed fibrils, a few with free ends. Context thin, ferruginous. Pores large, 3 to cm., rather shallow (3-4 mm. deep) bright, ferruginous (never glaucous) and with prominent setae.

This species is quite similar to *apiaria* but is smaller, thinner, and never has the dense coat of rigid hairs characteristic of *apiaria* in its prime. It is only known from Ceylon. Abundant specimens reached Berkeley and were by him referred to *crinigera* of Africa (from which it is quite different). Then a single specimen, having strayed into Paris, was named *Hexagona Deschampsii*.




Fig. 283

Hexagona aculeata. Type at Paris.

HEXAGONA ACULEATA (Fig. 283).—Color reddish brown, with appressed, fibrillose, zonate surface. Pileus thin. Pores medium, 5-6 to cm., regular. Color ferruginous.

This is known only from one collection made in French Guiana by Leprieur. It is in Montagne's herbarium, and there is also a co-type at Upsala. It has about the same sized pores as *Hexagona hirta*, but is a lighter colored plant, is thinner, and does not have the same dense coat of hairs.

HEXAGONA ELEGANS (Fig. 284).—Color dark, reddish brown. Surface with appressed, rigid fibrils and zonate. Pores medium, 5-6 to cm., 6-8 mm. deep, glaucous.

A single specimen of this is in the museum at Paris and its origin is not known. It is not as close to *Hexagona aculeata* as might appear from the

photograph. It is a thicker plant and the pores are glaucous, also the surface is not so strongly zoned.




Fig 284

Hexagona elegans. Type at Paris.



Fig. 285

Hexagona Dybowski. Type at Paris.

HEXAGONA DYBOWSKI (Fig. 285).—Pileus thin, flexible, of a pale color. Surface rugulose, zoned with a dense coat of pale, slender hairs, which are detersive, and old specimens evidently become almost smooth. Pores medium, pale, with angular walls disposed to become somewhat irpicoid.

There are three collections of this plant from the Congo, Africa, in the Museum at Paris, but it has never reached any other museum. It is a unique species, very different from all others of this section in its pale context color and the general color of the plant. Its affinities are rather with *Trametes* or *Polystictus* than with *Hexagona*.



Fig. 286

Hexagona Henschalli. Type at Kew.

HEXAGONA HENSCHALLI (Fig. 286).—Color reddish brown. Pileus thin, strongly zoned. Most of the zones are smooth or appressed fibrillose, a few with free fibrils. Pores large, rather shallow, glaucous, with thin walls and disposed to become a little irpicoid.

A single specimen is at Kew from Java, and named *Hexagona Henschalli* by Berkeley. It was never published, but was placed in the apiaria cover from which species it seems to me to be quite different.

GROUP 2, VELUTINUS.

We include in this group only the thin, velutinate plants that correspond to the group *Tenuis* in form and thickness. There are three other pubescent or velutinate plants (mentioned on page 14) which are included in other groups. As included here, the entire group might be considered a single species. All are very similar plants, thin with zonate, velutinate surface, and small, regular, shallow pores. It is chiefly an American group and abundant specimens are in the museums from the West Indies, Mexico, and South America. Of other than American specimens there are only three collections known, viz.: one each from New Caledonia, Africa, and Ceylon.



Fig. 287.

Hexagona variegata.

HEXAGONA VARIEGATA (Fig. 287).—Pileus thin, with ferruginous context. Surface velutinate with fine hairs and strongly marked with variegated, colored zones. Pores small, regular, shallow, smooth, usually ferruginous color, but sometimes glaucous.

This is a strongly marked species, the upper surface resembling bright forms of *Polystictus versicolor*. The contrast of zones usually alternate seal brown and blood brown. Sometimes plants are more evenly colored and then it runs into the next "species." It is a common plant in Mexico, Central America, West Indies, and South America, and many specimens are in the museums. Most of them are called *variegata*, and Berkeley so labeled most of his specimens. There is no question, however, that it is the same plant that Berkeley at a much earlier date named *Hexagona papyracea* and as he himself virtually so

stated. The name *variegata* is a much better name and has been generally employed.¹⁵

The two following may be called varieties or species, as you may prefer.



Fig. 288
Hexagona scutigera.

HEXAGONA SCUTIGERA (Fig. 288).— In Balansa's *exsiccateae* are found specimens so labeled on Spegazzini's authority. It is practically the same as the preceding except the surface is of a uniformly brownish color and is perhaps more rugulosely zoned. In any large collection of *Hexagona variegata*, however, all connecting forms occur, and it is at best a form. No type exists, but I judge it is correctly named, in which event it is a case for the date dictionary experts.




Fig. 289
Hexagona velutina. - Type at Paris.

¹⁵ Even Mr. Murrill uses it, though in order to excuse his disregard of dates he puts a question mark after *Hexagona papyracea*. If any doubtful mark should be used, it should be after *variegata*, for the type specimen of *Hexagona papyracea* is in good condition and unquestioned, and there is no type of *Hexagona variegata* so labeled. As a matter of fact I believe there is a "type" from which *variegata* was named, but it is labeled *Hexagona papyracea*.

HEXAGONA VELUTINA (Fig. 289).—The only specimen of this section known from Africa has a uniform brown color with narrow zones. It also has smaller pores. Otherwise it is the same as variegata. It is known from a single specimen and was called velutina. Nearly the same plant (one collection) reached Berkeley from Ceylon. He referred it to variegata.

Note.—The following species with pubescent pilei are placed in other sections: Pobeguini in Applanatus; macrotrema in Pallidus; bipindiensis in Pseudofavolus.

GROUP 3, UNGULAFORMIS.

(Hexagona of the "Fomes" type are few in number, but very marked species.)



Fig. 290.

Hexagona nitida. Type at Paris.

HEXAGONA NITIDA (Fig. 290).—Pileus with a hard, smooth, sulcate, polished crust. Context ferruginous, hard. Pores medium (5 to cm.) deep, $1\frac{1}{2}$ to $2\frac{1}{2}$ cm., reaching the crust.

This is the only *Hexagona* that occurs in Europe, and it is known only from two stations in the extreme south. It was first found in 1829 in the Pyrenees, and in Algeria in 1844. Both were on the live oak (*Quercus Ilex*) and came to Montagne, who named the species. He gave a good figure in Flora of Algeria. Then Dr. Marcucci seems to have collected it abundantly in Sardinia and his specimens (Exsiccatæ No. 69) are in most of the museums under the name "*Hexagona* (*Favolus*) Mori Poll.," Dr. Marcucci having referred it in error to an old figure of Pollini, to which it has little resemblance. Afterwards Baglietto, noting the mistake, naturally discovered it must be a new

species and called it *Hexagona Marcucciana*.¹⁶ At Paris, where it can be compared with Montagne's specimen, I doubt if one can be told from the other if they were transposed. I think Professor Maire has collected it in Greece, but I have seen none of his specimens. *Hexagona nitida* from what is known seems to occur only on the live oak (*Quercus Ilex*) and only in the Mediterranean countries. There are several of Marcucci's collections in the museums, mostly now badly eaten. The only good specimen I have seen is the Algerian collection in Montagne's herbarium, from which our figure (290) has been made.




Fig. 291

Hexagona Gunnii. Co-type at Paris.

HEXAGONA GUNNII (Fig. 291).—Pileus unguiform, with a thin, fragile, smooth, reddish brown crust which appears to me slightly lacate. Context thick, ferruginous.¹⁷ Pores large, concolorous, with thick walls.

This species is represented at Kew by several collections from Tasmania and Australia. I think it grows on Eucalyptus trees. Berkeley named it in 1839 as *Polyporus vesparius*, and then changed it (unfortunately without Otto Kuntze's consent) to *Hexagona Gunnii*.¹⁸

HEXAGONA SULCATA (Fig. 292).—Pileus subligneous, with a hard crust and deep, sulcate ridges. Context ferruginous. Pores medium (4-5 to cm.) deep, rigid, pale wood color.

¹⁶ This was in "Erbario Crittogamico Italiano," where Marcucci's collection was again distributed. The advertisement is given in Saccardo as "Bagl. & de Not," but they were both lichen men. I suspect Cesati was really responsible for it, as he seems to have been the chief fungus man of these exsiccatae.

¹⁷ I think this is the only species known with a strong development of the context. Usually the pores almost reach the crust.

¹⁸ I do not know why Berkeley changed the specific name when he put it in *Hexagona*, but he no doubt had good reasons for it, and he thought he had the right. At any rate it was before our "lawmakers" had legislated on the subject as to what a man has a right to do in his own private affairs.

This strongly marked species is only known from Ceylon. Berkeley published it in 1847 with a good figure and sent specimens to both Fries and Montagne. That sent to Montagne (Fig. 292) was typically sulcate, but the specimen to Fries (Fig. 293) was more even. Berkeley did not retain a specimen in his own herbarium, and when some twenty years later he received the smooth form also from Ceylon he described it as *Hexagona durissima*. It is the same as the specimen of *sulcata* he sent Fries. Whether or not it is the same species as *sulcata* I do not know, but I think probably only a smooth form.



Fig. 292.
Hexagona sulcata.




Fig. 293.
Hexagona durissima.

HEXAGONA DURISSIMA (Fig. 293).—This seems to be practically the same plant as *sulcata* except it has a more even crust. It is known only from Ceylon, but recently I have seen a specimen so referred, and probably correctly, from Java.

HEXAGONA RESINOSA (Fig. 294).—Pileus applanate, with a dark resinous crust. Context hard, sub-woody, pale alutaceous or pale ferruginous. Tubes medium, 5 to cm., 1 to 1½ cm. deep, pale color, with rigid walls. Spores (teste Murrill) hyaline, smooth, 4 x 6.

This species was recently well named by Murrill, from the Philippines, and is very different from all others. It is known only from one collection or record.¹⁹ Its natural relations I think are with *Fomes pinicola*, the same

¹⁹ *Polystictus Copelandii*, as distributed by the Philippine Bureau of Science, No. 1214 (specimen sent to Kew), is evidently *Hexagona resinosa*, through some transposition of specimens. Mr. Murrill has introduced enough confusion with his jargon of names among the Philippine polyporoids, without having the subject further confused by transposition of specimens in distributed sets.

resinous crust, same context, and the coloration both of context and crust is similar. There is no similar plant in the genus *Hexagona*, and it might well be made the type of a new genus. We place it in this section on account of its evident "Fomes" relationship, though as to form it belongs in the next, and as to context color it approaches the section *Pallidus*.



Fig. 294.

Hexagona resinosa. Co-type in museum at Berlin.

Note.—*Hexagona laevis* was based on nondescript material from Andaman Islands. There is one poor specimen at Kew and another at the British Museum. I judge it belongs in this section.

GROUP 4, APPLANATUS.

This is an artificial group to include species that have no one prominent character to throw them into other groups, and which are flat but not too thin.

HEXAGONA POBEGUINI (Fig. 295).—Pileus applanate, with concentric, sulcate zones, and minutely pubescent. I think the pubescence wears off to a certain extent on old specimens. Context subligneous, harder than others of this group, ferruginous. Pores large 2-3 to cm., ferruginous, with rigid walls and setae.

This seems to be a frequent plant in Africa, and several collections are at Paris and Berlin. One at the British Museum was named *Hexagona Welwitschii*. In fact, the plant was discovered to be a "new species" in each of

the three museums where it is to be found.²⁰ The zones of the pileus are sometimes (in the type specimen) colored with different shades of brown. The pores of the type were crenate, but that was only an accidental character of this particular specimen. The pores vary in size, as shown in our figures.



Fig 295
Hexagona Pobeguini.

HEXAGONA NIAM-NIAMENSIS (Fig. 296).—Pileus *smooth*, unicolorous, with narrow zones. As to surface it much resembles *Hexagona tenuis*. Context ferruginous. Pores medium, about 4 to cm., regular, with thin hexagonal walls, $\frac{1}{2}$ cm. deep. They have no evident setae and some of them (not all) are glaucous.

This is known from a single specimen (Fig. 296) from Africa at Berlin. The specific name, while alleged to be Latin, is more probably from an Ethiopian dialect. It is a terrible misfortune for a plant to have to bear such a name as that.

²⁰ In a case of this kind we are very much disposed to take the best name, as we believe plants should be given decent names. Where a poor plant has had the misfortune to be named Pobeguini, Stuhlmanni, and Welwitschii, there is not much choice.




Fig. 296.

Hexagona niam-niamensis. Type at Berlin.




Fig. 297

Hexagona chartacea. Type at Paris.

HEXAGONA CHARTACEA (Fig. 297).—Pileus rather thin, smooth, with narrow zones. Pores large, about 3 to cm., 5 mm. deep, with thin walls. Color ferruginous and setae evident.

This is rather a thin species for this group and is known from two collections from Africa. One of the collections was named *Hexagona obversa*, but seems to be too close to be kept distinct. I can see very little application of the name *chartacea* to this plant.




Fig. 298.

Hexagona leprosa. Type at Upsala.

HEXAGONA LEPROSA (Fig. 298).—Pileus with a thin, dull, slightly pubescent crust, not zoned. Context soft, spongy, ferruginous. Pores medium, 4-5 to cm., 1½-2 cm. long, with thin walls. The inner surface of the pores is pubescent under the lens, with short, pale hairs. The color of the entire plant is almost uniform.

This is known only from a single specimen, collected in the West Indies,²¹ about sixty years ago and preserved in a jar at Upsala. There is a small co-type fragment also at Kew.

²¹ It has been stated "also in Brazil." I know not the source of this statement, but am sure there is no specimen from Brazil in any museum of Europe that I have visited.

HEXAGONA SPECIOSA (Fig. 299).—Pileus with a thin, zoned, smooth crust. Context and pores ferruginous. Pores medium large, about 4-5 to cm., 1-1½ cm. deep, with thin walls becoming lacerate.

In size, shape, and color this plant is much like the preceding. It is known from a single specimen in a jar at Upsala and a co-type at Kew. It came from South Africa sixty years ago.




Fig. 299

Hexagona speciosa. Type at Upsala.



Fig. 300.

Hexagona Kurzii.

HEXAGONA KURZII (Fig. 300).—Surface dark reddish brown, rugulose, zoned. Pores medium 5 to cm., 5 mm. deep, strongly glaucous.

This came from India and has a general resemblance to Hexagona polygramma.²² The pores are too deep, however, to be entered in the section with polygramma.

HEXAGONA ERUBESCENS (Fig. 301).—Pileus *rigid*, about 2 cm. thick, with a smooth zoned crust. Pores irregular, angular, about 5 to cm., 1-1½ cm. deep, with *rigid*, rather thick walls. No setae.

²² The co-type collection is at Kew on a sheet of polygramma.

This is based on a collection by Spruce, Brazil, and is at Kew. The collector states "Hymenium vinosum," hence Berkeley named it *erubescens*. The pores have lost all vinous color now. The strong character of the species for me is its *rigidity*, both of pileus and pores. The species is found in Saccardo in section "Hirtae." The type has not a sign of a hair of any kind. There is a collection at Kew, however, that was referred to *erubescens*, and which has appressed fibrils, but I think it quite another (and a "new") species. I should prefer that some one else name it.




Fig. 301.

Hexagona erubescens. Type at Kew.




Fig. 302.

Hexagona amplexens.

HEXAGONA AM-
PLEXENS (Fig. 302).—
Pileus small, ungulaform,
gibbose, smooth, with sulcate zones. Context brown,
suberose. Pores 5-6 to cm.,
5-8 mm. deep, concolorous,
with thin walls, no setae.

This little species is unique in size and shape. It is known from one collection in the herbarium of Patouillard, and came from New Caledonia. It evidently grew on small branches which it partially encircled.

SECTION 5, TENUIS.

This group is the most important of all for it embraces the only common species that grows in many warm countries, viz.: *Hexagona tenuis*. They are *thin* plants, rarely over two or three mm. thick, with smooth, concentric-zoned pilei. The pores are small and shallow. Sometimes the plants are called membranaceous, but I think are not thin enough to be called membranes.



Fig 303

Hexagona tenuis. (Type form.)

HEXAGONA TENUIS (Fig. 303).—Pileus rigid, with a smooth, concentric-zoned surface. Context thin, about 2 mm., ferruginous. Pores small, regular, round, 8-12 to cm., shallow.



Fig. 304

Rugulose form classed as *Hexagona tenuis*. (Kew.)

This is a widely distributed plant and occurs in most warm countries of the world. Like all widely distributed plants it varies, and it is not practicable to maintain all the specific names that have been given to it. As to pore size those with the smallest pores were named *Hexagona pulchella* (Fig. 305), then the type size (Fig. 306) and the largest size (Fig. 307) were called *Hexagona polygramma*. With hardly two collections with exactly the same sized pores, it is difficult to maintain these "species." And yet the pore sizes have some value for they are usually uniform in specimens of the same collection. They also vary as to color, and particularly in the development of a glaucous pore covering. Some collections have no sign of it, others are partially glaucous, and others strongly glaucous. *Hexagona cervino-plumbea* is only a glaucous form.




Fig. 305.

Fig. 306

Fig. 307

Comparative pore sizes. Fig. 305, *pulchella*. Fig. 306, *tenuis* (type). Fig. 307, *polygramma*.

History.—The first specimen recorded was brought by Humboldt from South America and is still preserved in Hooker's herbarium. It was published by Hooker as *Boletus tenuis* in Kunth Synopsis (1822) and in the preceding paragraph an anomaly of the same species as *Boletus reticulatus*.²³ However, this was not the first specimen to reach Europe, for it is found in the Linnæan herbarium with no clue to its source. It is labeled "*Boletus favus*, Linn.," an obvious error as pointed out first by Klotzsch, then by Berkeley, and very recently by Mr. Murrill.²⁴ *Hexagona tenuis* is a very common species in many

²³ As this was published at a "previous date," according to Kuntze's method of reckoning dates, it was necessary to find another species called *reticulatus* to put forth as a reason for not taking the name. This was not a *Hexagona*, but that was a minor matter compared to the importance of Hooker having published *reticulatus* in a previous paragraph to *tenuis* in the same book. It was Klotzsch who first recognized that *reticulatus* was only an altered condition of *tenuis*, and he so indorsed it on the label, from whence was obtained the information that was dilated upon at length recently, forgetting to mention that it had all been published in full by both Klotzsch and Berkeley many years ago.

²⁴ "Im Linne'schen Herbarium, *Boletus favus* ist *Polyporus tenuis*, Hooker."—Klotzsch, 1832.

"*Hexagona tenuis* is marked in the Linnæan herbarium *Boletus favus*, but not by Linnæus, with whose description it does not correspond. The name is evidently not authoritative."—Berkeley, 1842.

"This species is found in the Linnæan herbarium marked *Boletus favus*, but not by Linnæus, nor with his sanction."—Murrill, 1905.

Had Mr. Murrill, instead of copying Berkeley, done a little investigating in the Linnæan herbarium he would have found that "this species" was named by Dickson many years after Linnæus died, and under the circumstances he would have had considerable trouble in obtaining Linnæus' "sanction."

countries and like all such species has been discovered to be "new" on numerous occasions.²⁵ A number of these seem to me to be absolutely the same plant, and I can see no difference whatever on which to base "new species."²⁶ Others do differ slightly from the type form, but whether this is of specific importance or not it is difficult to say.

Forms of *Hexagona tenuis* or related plants.

HEXAGONA PULCHELLA (Fig. 305).—This plant from Java seems exactly the same as the type form except smaller pores.




Fig. 308.

Hexagona polygramma.

HEXAGONA POLYGRAMMA (Fig. 308).—Originally from Cuba, the type is practically the same as that of *tenuis* with pores slightly larger. In most museums, however, all these similar plants are arranged in two covers, one "*Hexagona tenuis*, Hooker," the other "*Hexagona polygramma*, Mont." I can not believe that the namers have any distinct idea of a difference, for in both covers I have found indiscriminately collections varying as follows:

Size of spores.—From very small, as shown in Fig. 305, to size medium, as shown in Fig. 307.

Surface.—Relatively smooth and evenly zoned, as Fig. 303, to strongly rugulose, as shown in Fig. 304.

Color of pileus.—Very pale, almost white to brown, and many deep reddish brown.

²⁵ Not necessarily all, however, that are raked up and tabulated. Thus "*Polyporus bivalvis*, Pers.," given as a synonym, has little resemblance to it and is not a *Hexagona*. A good specimen is in Persoon's herbarium. "*Hexagona cingulata*, Lév.," and "*Hexagona unicolor*, Fries," are also said to be synonyms, but that is only a vague guess, as no specimens of either exist, and the compiler knew nothing about them.

²⁶ For further details see list of synonyms, pages 43 to 45.

Color of pores.—Pure cinnamon or ferruginous to dark (fuscus), sometimes bright silvery, glaucous.

At Kew there are ninety-two collections in these two covers and scarcely any two of them exactly the same. Under these conditions it is only practical to do as has been done and refer all to one or two species.

The following we should consider as forms of *Hexagona tenuis*, and we could manufacture as many more if we were so disposed.



Fig. 309

Hexagona umbrinella. Small one is the type at Upsala.

HEXAGONA UMBRINELLA (Fig. 309).—This has a dark reddish brown, rugulose surface.²⁷ The same thing is also called *Hexagona Dregreana*. *Hexagona Boneana* is also too close.

HEXAGONA CONCINNA is a very thin plant with small, dark pores and dark reddish pileus.

HEXAGONA DISCOPODA is a plant with the reddish stain only *partially developed* over the base of the pileus so that the plant is decidedly *two-colored*. It seems to be a frequent form in Africa and abundant specimens reached Hennings and were referred by him to *Hexagona polygramma*. It is probably the same as *tricolor* named by Fries from Africa many years ago (because of its color contrasts), but no specimen is known now.

HEXAGONA SUBTENUIS was named by Berkeley from India, but I think not published. It has ferruginous colored pores that to me under a lens appear slightly pubescent.

HEXAGONA PHAEOPHORA is a form with pale pileus and *dark* pores.

All the preceding are *thin* plants, very similar to *Hexagona tenuis*, and differing principally in color, pore color, and pore size. The four following, *rigida*, *similis*, *Muelleri*, and *nigrocincta*, are thicker plants (relatively), but with the same general characters and small regular pores. They differ among themselves as do the forms of *tenuis*. All come from the same geographical region, Australia, New Caledonia, and the Pacific Islands, and all, I think, are better considered as forms of one species.

HEXAGONA RIGIDA (Fig. 310).—Pileus about 3 mm. thick, with smooth or slightly rugulose zoned surface and small regular pores.

²⁷ The "co-type" that Fries sent Berkeley of *Hexagona umbrinella* is a much thicker specimen than is to be found in his own herbarium. I think it is not the same species.

The plant is close to *Hexagona tenuis* as to color and general appearance, with slightly deeper pores. It came from Australia. *Hexagona Muelleri*, also from Australia, and based on a single specimen, is practically the same with slightly more rugulose surface. *Hexagona nigrocincta* is for me a pale form of *rigida*, paler color and smoother than type of *rigida*. It came from New Caledonia, and the ordinary form of *rigida* also occurs there.



Fig. 310.

Hexagona rigida. Type at Kew.

HEXAGONA SIMILIS.—The type specimen is of a very dark color with agglutinate, fibrillose zoned surface, so that it has relations to the section *Setosus*. None of the fibrils are free, however, and I think it is closer to *rigida*. The pores are quite small and dark colored. The “type” is the only one in the cover that has the agglutinate fibrils strongly marked. Others so referred seem to me much closer to *rigida*.

HEXAGONA ATROSANGUINEA (Fig. 311).—Plant growing on under side of stick, and largely resupinate with narrow, pileate margin. Pileus thin, smooth, deep *blood brown* color. Pores small, rugulose, shallow, many *colored similar to the pileus*, and also with a glaucous deposit.

This species is very marked and the only one I have noted where the pores are strongly colored red-brown. Its habit of growth is also different from usual, and abundant specimens at Berlin all seem to have the same habits. Whether it ever takes a truly pileate form I do not know, but I judge not, as I think all the abundant types at Berlin are of the same nature. It came from Africa and is only found in the museum at Berlin.

HEXAGONA SACLEUXII (Fig. 312).—Pileus rigid, with a smooth, pale, strongly concentrically ridged surface. Pores vary in the

same collection (as shown in our figures) as to size. The large pores are about 5 mm. deep and strongly glaucous.




Fig. 311

Hexagona atrosanguinea. Type at Berlin.



Fig. 312

Hexagona Sacleuxii. Type at Paris.

This is known from three collections at Paris, all from Africa. The pores are deeper than others of this group, otherwise it is close to *rigida*. The variation of pore sizes in the same collection is unusual, and I am not sure but that it is due to different ages. The large pores are strongly glaucous, the small pores not at all.

SECTION 6, PALLIDUS.

This section is quite distinct from all that precede in the *pale color* of its context. One species (*albida*) when fresh is *pure white* but discolours some in drying. The usual color of the museum specimens of this section may be called pale ochraceous or isabelline. I doubt if there are any in this section that are true *Hexagonas*. Probably all have the character of the *variation of the hymenium* strongly developed. In *Hexagona albida* (cfr. Figs. 313 and 314) and particularly in *Hexagona ochroleuca* the hymenium takes *hexagonal, daedaloid, and lenzitoid* forms and this tendency to *variation* is as much a *specific character* as any character a species can have.




Fig 313

Fig 314

Hexagona albida. Photographed in Samoa.

HEXAGONA ALBIDA (Figs. 313 and 314).—Pileus pure white with soft, smooth, faintly zonate surface. Context soft, almost fleshy when fresh, in drying it becomes discolored in time and more tough. Pores large, irregular, about 5 mm. deep with thin walls. Sometimes lenzitoid forms (Fig. 313) are found growing with the hexagonal forms and our figures (313 and 314) are specimens that grew from the *same mycelium*.

History.—This plant was described under this name by Berkeley from the Philippines, and seems to occur mostly in Australia and the Pacific Islands. I found it in Samoa, but it is not common there. In Samoa it usually took the hexagonal form, rarely the lenzitoid form, but in other localities it may run more often to lenzitoid forms. It is the same plant, I believe, as *Daedalea inconcinna*, also from the Philippines, and *Daedalea intermedia* from Australia. I think *Hexagona Cesatii* from Borneo is exactly the same thing, with a tendency to become a little cyclomyroid. When the history of these polymorphic plants is worked out it will probably be found to have names in other genera such as *Lenzites*.




Fig. 315

Hexagona macrotrema. Type at Leiden.

HEXAGONA MACROTREMA (Fig. 315).—The description of *Hexagona albida* covers this species also, for it is the same thing excepting that the surface is distinctly pubescent. However, they are undoubtedly forms of the same species and they occur over the same region (Pacific Islands).

Hexagona macrotrema was first collected by Junghuhn in Java and so named by him on the label. The specimen is to-day found in Leiden in good condition and bears only Junghuhn's original label. Before he published it, however, Lévillé visited the museum, saw the specimen, changed its name, and published it as *Hexagona Molkenboeri*. This did not please Junghuhn (naturally) and he wrote to Fries, who, when he published it used Junghuhn's name.²⁸ The name *Molkenboeri* is therefore "prior" from an Otto Kuntze point of view, and while I have great regard for priority it does not appeal to me when served with so much rascality. Therefore I use the name *macrotrema*.

Hexagona macrotrema is rather rare in the museums and has been mostly named *albida*. I have seen only the following three specimens: Java (Leiden),

²⁸ The plant is found in Saccardo, vol. 6, p. 369, under both names, pretending to be two different species, although both names were based on exactly the same specimen.

New Guinea (Berlin), Tropical Africa (Kew). As in the case of *Hexagona albida*, it is probable that it takes other hymenial forms.

HEXAGONA OCHROLEUCA (Figs. 316 to 319).—We shall not enter here into any detailed account of *Hexagona ochroleuca*, for it is usually not a *Hexagona*. In fact the name *glabra* is the only specific name that was given to it as a *Hexagona*, although it has a dozen other names, as *Trametes*, *Daedalea*, *Lenzites*, *Sistrotrema*, etc. Generally it is a *Lenzites*, and if we ever consider it in detail it will be as a *Lenzites*, its usual form.

Hexagona ochroleuca is the most polymorphic species known, I think, and takes hexagonal, lenzitoid, irpicoid, and daedaloid forms, often *in the same specimen*. Our figure (318) shows three distinct hymenial forms. The hexagonal forms are rare and the type of *glabra* (Fig. 317) is the only one so named as a *Hexagona*, though several "species" of *Trametes* are based on the same thing. I have seen many lenzitoid forms. Lévillé named this specimen *Hexagona glabra*, and another specimen of the same collection (Roux, India) he called in the same paper *Sistrotrema ochroleucum*. These plants are in the same cover at Paris, and they are surely the same species notwithstanding the hymenium is so different.

Hexagona ochroleuca has but few constant characters, none of a hymenial nature, and can only be learned by experience. Its consistency, *color of context* (alutaceous, not white when fresh), surface, and *distant* plates are the main characters by which it can be known from its equally abundant and equally polymorphic neighbor, *Lenzites repanda*. *Hexagona ochroleuca*, in its various forms as *Trametes*, *Lenzites*, etc., is a very abundant plant in India, Java, Philippines, and the East in general, and also in Australia.

We have not thoroughly investigated its synonymy, though we believe the following should be included: *Polystictus lenziteus* (Zollinger Col.), *Sistrotrema ochroleucum*, *Hexagona glabra*, *Daedalea lurida*, *Daedalea pruinosa*, all by Lévillé, who seems to have discovered it was a "new species" every time he saw a specimen.

Trametes Beyrichii (as to Berkeley's Philippine determination, Cummings 2202),²⁹ *Trametes colliculosa* from Ceylon, *Trametes lobata* from India, *Trametes laeticolor* from Ceylon, *Daedalea Hobsoni* from India (or Australia?),³⁰ and numerous recent determinations from the Philippines.³¹ *Daedalea Schomburgkii*

²⁹ As to Fries, from Brazil, it is doubtful, as *Hexagona ochroleuca* is not known from America. No specimen of *Trametes Beyrichii* exists, and what it was is unknown.

³⁰ *Daedalea Hobsoni* was published in a paper on Australian fungi and was based on a collection cited, made by Schomburg in Australia. Hobson collected in India, and Berkeley refers to his specimen incidentally as "the original specimen," though never formally published. Under these conditions our lawmakers ought to specifically tell us which is the "type locality" and which the "type specimen," as they put so much stress on those things. In this instance I think it makes no vital difference, because both are the same plant. Cooke afterwards discovered the "type specimen" of *Daedalea Hobsoni*, which Berkeley had labeled *Daedalea Schomburgkii* and sent Saccardo (cf. Vol. 6, p. 376) a description of this interesting "new species," though based on exactly the same specimen that Berkeley had described sixteen years before. As evidence of the value of our literature Saccardo puts it in a different section of the genus from the one in which he places *Daedalea Hobsoni*, although both were based on the same specimen.

³¹ It appears to me that Mr. Murrill's priority investigations were very superficial as to these plants (as with most others). He uses the name *Hobsoni* (1865), and it has a dozen names "prior" to that. Bresadola habitually calls the plant "*Daedalea lenzitea* (Lév.), Bres.," which was 1854, and Lévillé had four names prior to lenzitea, to say nothing of Berkeley's discoveries. In the whole list it does not have a suitable name, or I should use it without regard to the date. Lévillé's name *ochroleuca* is probably the best. I presume, however, some enterprising individual could take the synonyms I have cited, look up their dates, arrange them chronologically, and produce weighty evidence why *ochroleuca* can not be used.



Fig. 316



Fig 317.



Fig. 318



Fig 319.

Hexagona ochroleuca. Figs. 316 and 317, type forms (as *Hexagona*). Fig. 318, a specimen reduced, showing three variations of the hymenium. Fig. 319, an irpicoid form named *Sistrotrema ochroleucum*.

from Australia, *Daedalea tenuis* from Philippines, *Daedalea aulacophylla* from Australia, *Daedalea flabellum* from Andaman Islands, *Daedalea ochracea* from India, *Daedalea sub-confragosa* from the Philippines, *Lenzites Guilfoylei* from Australia, *Lenzites ochrophyllus* from India.

The old mycologists took the "genera" *Trametes*, *Daedalea*, *Lenzites*, etc., literally, and based a new species on almost every specimen of this plant in every hymenial form that reached them. When the science of mycology gets beyond its "new species" babyhood, and workers take a broader view of species than single "type" collections, then I believe all the species I have mentioned will be held to be the same plant. I call it in this paper *Hexagona ochroleuca*, but were I writing on *Trametes* it would be *Trametes ochroleucum*. The same applies to *Daedalea*, *Lenzites*, or *Irpex*. I think its better name is as *Lenzites* to correspond with its usual hymenial form, and the equally polymorphic *Lenzites repanda*.



Fig. 320.

HEXAGONA VESPACEA (Fig. 320).—This may be exactly the same plant as *macrotrema*. It was one of the early Persoonian names (1826) and came from the island of Rawak. No other collection has ever been referred to Persoon's name, and the original collection is only known from two little specimens, one at Paris (Fig. 320), another in Persoon's herbarium at Leiden. These are *thinner* than specimens of the preceding species, and *darker*, though the dark color may be due to age. Persoon described them as smooth, but that they are somewhat pubescent can be seen from our photograph. (Fig. 320.)

HEXAGONA SEURATI (Fig. 321).—Context surface and pores unicolorous, pale alutaceous or isabelline. Context soft, homogeneous



Fig. 321

Hexagona Seurati. Type at Paris.

with the pores. Surface smooth, no distinct crust. Pores large, shallow, many superficial.

This species is known only from one collection (Fig. 321) in the herbarium of Professor Patouillard. It came from Raiatea, one of the Society Islands.

HEXAGONA AEQUALIS (Fig. 322).—No better description of this can be given than to say that it is a hexagonal, tropical form of *Daedalea quercina*. The color, context, surface, everything, is exactly the same as the common



Fig. 322.

Hexagona aequalis. Type at Paris.

plant of Europe excepting the hymenial configuration. The pores, as will be seen from our photograph, are not truly hexagonal, but tend to daedaloid. *Daedalea quercina* is presumed not to occur in the tropics. If it does, this plant must be referred to it. It is known from but one collection, South America.³²



Fig. 323.
Hexagona rhombipora.

HEXAGONA RHOMBI-PORA (Fig. 323).—Color pale alutaceous or isabelline, concolorous. Context thin. Surface smooth, no distinct crust. Pores large, flaccid, concolorous, tending to favoloid.

This is known from a single half specimen (Fig. 323) in the herbarium of Montagne. It came from Brazil. No similar specimen has ever reached Europe from South America. I have an impression that it is an aberrant, hexagonal form of some *Lenzites* perhaps.

³² I think it was a Mr. Smith who some years ago distributed some specimens from Central America "determined" by Ellis. Among others was a specimen labeled "*Irpex maximus*." It has no resemblance whatever to "*Irpex maximus*," which is only an irpicoid condition of the common *Polystictus occidentalis* of the tropics. Ellis, of course, had no way of knowing that, and his determinations of tropical species were but little more than a vague guess. The plant that he called "*Irpex maximus*" I have always considered as a tropical, irpicoid form of *Daedalea quercina*, the same as I consider *Hexagona aequalis* to be a tropical, hexagonoid form of the same species.

GROUP 6, PSEUDOFAVOLUS.

This section differs from all that precede it in its fleshy, tough nature, rather than corky-woody. It is not usually classed in *Hexagona*.³³ Several have been placed in *Favolus*, from which it differs in the basic idea of the genus *Favolus*, viz.: the shape of the pores. In my opinion, the species should be included in *Hexagona*, or if not, should be made into a separate genus. There are but few species known, including, however, the only *Hexagona* known from the United States.




Fig. 324

Hexagona cucullata, natural size and pores x6.

HEXAGONA CUCULLATA (Fig. 324).—Pileus orbicular, reniform, attached by a short disk-like stem. Surface smooth, even, when fresh Mars yellow, when old deep, reddish brown. Pores concolorous, orbicular, a scant mm. wide, shallow.

This seems to be a rather rare plant, occurring in Southern United States, West Indies, and South America. At Kew there are but ten collections, including one from Ceylon (but probably the same) named by Cesati, *Favolus chartaceus*. In addition it has two other synonyms from the United States, *Favolus curtipes* and *Favolus Taxodii*, and I think a third, the recently described *Pseudofavolus auriculatus*³⁴ from Louisiana.

³³ But one of the species, *Hexagona Miquelii*, is placed in this genus in Saccardo.

³⁴ I have seen no specimen of this, as when I called it was not to be found. I feel so well convinced, however, that *Hexagona cucullata* is the only one we have in the United States that I have very little doubt as to its being the same thing.

Related plant.—POLYPORUS ORINOCENSIS (Fig. 325).—With the same color and other characters except its small pores, we mention Polyporus Orinocensis here, for we feel it is a very closely related plant notwithstanding its small pores. Professor Patouillard now places it in the same section with cucullata. It has small pores and they are paler than those of cucullata. When fresh they were probably white. It would not do to classify such a small-pored plant as a Hexagona and we wish Nature would be more consistent and make her species so they

would fit into the man-made genera. It would be so much simpler. Polyporus Orinocensis is known from but one collection (Fig. 325) in the herbarium of Professor Patouillard.



Fig. 325.




Fig. 326

Hexagona Miquelii. Type at Paris.

HEXAGONA MIQUELII (Fig. 326).—Pileus orbicular, reniform. Color deep reddish brown. Surface glabrous, but *strongly tessellate*. Pores orbicular, shallow, colored.

This, as to coloration, texture, and all characters, is exactly the same as Hexagona cucullata, except that it is strongly marked with a tessellate pileus. It is a very rare plant and but three specimens have ever reached Europe, all of which were discovered to be "new species." First, from Surinam, named Polyporus Miquelii by Montagne, a nice specimen (Fig. 326) in the herbarium of Montagne. Then from Java, by Zollinger, named Polyporus pustulosus by Lèveillé, specimen in the herbarium of Professor Patouillard. The third from St. Domingo, named by Berkeley Favolus induratus. All are exactly the same plant.³⁵

HEXAGONA BIPINDIENSIS (Figs. 327 and 328).—Pileus thin, orbicular or reniform. Color reddish brown. Surface minutely velutinate, *strongly tessellate*. Pores pale, probably white when fresh, medium round, shallow.

³⁵ In a case of this kind, when Mr. Murrill uses the last name, induratus, his reasons are very puzzling to understand. As he has made so much fuss about "priority," we do not know whether he does not know it is the same plant or whether he thinks 1852 is prior to 1841.

This is represented by an abundant collection at Berlin. It came from Africa. It was named as *Hexagona bipindiensis* on the label by Hennings, but I do not know whether this was published. It is not his *Favolus bipindiensis*. A single specimen of what seems to me practically the same is found in Patouillard's herbarium under the name *Favolus velutinus* (Fig. 328). It came from Tonkin and has the same peculiar, velutinate surface. The pores are a little



Fig. 327



Fig. 328

Hexagona bipindiensis. Fig. 327 is type at Berlin.

larger and not so regular. I believe it to be, however, the same plant. The specific name *velutina* can not be used for a *Hexagona* as it is already occupied.

HEXAGONA MIRABILIS (Fig. 329).—Pileus white, smooth, thin, with a thin crust. Context none, the pores reaching the crust. Pores 1 to 2 mm. deep, 10-12 to cm., round or hexagonal, white.

This was a rare plant that I collected in Samoa in but one locality. When fresh it was pure white and a marked species, being so different from ordinary polyporoids. At that time I was not acquainted with any species of this group and was entirely at a loss to know where to place the plant. It is the only white *Hexagona* known in this section.




Fig. 329

Hexagona mirabilis. Photographed in Samoa.

SECTION 8, RESUPINATUS.

I believe there are no truly resupinate *Hexagonas*. The only one that has any claim is *heteropora*, and that is probably a resupinate form of something else. Some resupinate plants that have been named as *Hexagonas*, such as *carbonaria* and *Bartlettii*, are evidently so closely related to the ferruginous *Porias* that we shall so place them.



Fig. 330.

Hexagona heteropora. Types at Paris.

HEXAGONA HETEROPORA (Fig. 330).—Context pale. The remainder of the "description" can be made from our photograph. But three collections have been so named, which came from South America. We present photographs of all of them. The plant was named heteropora from the varying size of the pores, and it is evidently (from our figure) well named, if they are all the same species, which I doubt.

APPENDIX I.

NOMINA CONSERVANDA.

The following is an alphabetical list of the names of Hexagonas that we would "conserve." It does not have the formal sanction of our professional law-makers, but we think it has a better claim, namely, use, merit, and truth.

We give in addition to the name the country whence described, and as we think the name and country are the most important, we place them in heavy-face type. In lighter face type we summarize other details, viz.: the book citation, where published, and, what is more important, the museums where the type specimens are preserved. We also give what is of least importance of all, except to the parties concerned, the names of the wonderful discoverers.

ACULEATA—*South America*.—Ann. Sci. Nat. 2, vol. 13, p. 205. Montagne. Type, Museum at Paris.

AEQUALIS—*South America*.—Journ. de Bot., vol. 3, p. 258. Patouillard. Type in his herbarium.

ALBIDA—*Philippines*.—Jour. Linn. Soc., vol. 16, p. 47. Berkeley. Type at Kew.

AMPLEXENS—*New Caledonia*.—Bull. Soc. Myc. de France, vol. 18, p. 299. Patouillard. Type in his herbarium.

APIARIA—*East Indies*.—Voyage of Uranie, p. 169 (as Polyporus). Persoon. Type in museum at Paris.

ATROSANGUINEA—*Africa*.—Engler's Jahrb., vol. 23, p. 545. Hennings. Type at Berlin.

BIPINDIENSIS—*Africa*.—Not published, as far as I know. Hennings. Type at Berlin.

CAPILLACEA—*South America*.—Bull. Soc. Myc. de France, vol. 4, p. 36. Patouillard. Type in his herbarium.

CHARTACEA—*Africa*.—Bull. Soc. Myc. de France, vol. 9, p. 209. Patouillard. Type in museum at Paris.

CONCINNA—*Africa*.—Bull. Soc. Myc. France, vol. 9, p. 209. Patouillard. Type in museum at Paris.

CUCULLATA—*Cuba*.—Ann. Sci. Nat. 2, vol. 17, p. 125 (as Favolus). Montagne. Type in museum at Paris.

DESCHAMPSII—*Ceylon*.—Bull. Soc. Myc. France, vol. 7, p. 207. Hariot. Type in museum at Paris.

DISCOPODA—*Africa*.—Bull. Soc. Myc. France, vol. 9, p. 209. Patouillard. Type in museum at Paris.

DURISSIMA—*Ceylon*.—Jour. Linn. Soc., vol. 14, p. 57. Berkeley. Type at Kew.

DYBOWSKI—*Africa*.—Bull. Soc. Myc. France, vol. 8, p. 54. Patouillard. Type in museum at Paris.

ELEGANS.—Unknown, probably from Africa. Bull. Soc. Myc. France, vol. 7, p. 207. Hariot. Type in museum at Paris.

ERUBESCENS—Brazil.—Hooker's Jour. 1856, p. 237. Berkeley. Type at Kew.

GUNNII—Tasmania.—Flora of Tasmania, vol. 2, p. 255. Berkeley. Type at Kew.

HENSCHALLI—Java.—See page 11; not previously published. Berkeley. Type at Kew.

HETEROPORA—South America.—Jour. de Bot., vol. 3, p. 166. Patouillard. Type in museum at Paris.

HIRTA—Africa.—Flore d'Oware, p. 1, t. 1 (as Favolus). Palisot-de-Beauvois. Type at Geneva.

KURZII—India.—Trans. Linn. Soc., 2d ser., vol. 1, p. 126. Currey. Type at Kew, on sheet of polygramma.

LEPROSA—West Indies.—Nov. Symb., p. 101. Fries. Type in jar in museum at Upsala.

MACROTREMA—Java.—Nov. Symb., p. 101 (Junghuhn). Fries. Type in error in Box 42 of Persoon's herbarium at Leiden.

MIQUELII—South America.—Ann. Sci. Nat. 3, vol. 4, p. 357 (as Polyporus). Montagne. Type in museum at Paris.

MIRABILIS—Samoa.—Described on page 37. Type deposited at Kew.

NIAM-NIAMENSIS—Africa.—Engler's Jahrb., vol. 14, p. 348. Hennings. Type in museum at Berlin.

NITIDA—Algeria.—Sylloge, p. 170. Montagne. Type in museum at Paris.

OCHROLEUCA—India.—Ann. Sci. Nat. 3, vol. 5, p. 145 (as Sistrotrema). Léveillé. Type in museum at Paris. It is only a hexagonal form of a Lenzites.

PHAEOPHORA—China.—Bull. Soc. Myc., vol. 23, p. 74. Patouillard. Type in his herbarium.

POBEGUINI—Africa.—Bull. Soc. Myc. France, vol. 8, p. 28. Hariot. Type in museum at Paris. (Did not get in Saccardo's sweep net.)

POLYGRAMMA—Cuba.—Ann. Sci. Nat. 2, vol. 8, p. 365 (as Polyporus). Montagne. Type in museum at Paris.

PULCHELLA—Java.—Ann. Sci. Nat. 3, vol. 2, p. 200. Léveillé. Co-type in herbarium of Patouillard. It is only a small pored form of Hexagona tenuis.

RESINOSA—Philippines.—Bull. Torr. Club, vol. 35, p. 398. Murrill. Co-type in museum at Berlin.

RHOMBIPORA—South America.—Ann. Sci. Nat. 4, vol. 5, p. 370. Montagne. Type in museum at Paris.

RIGIDA—Pacific Islands.—Jour. Linn. Soc., vol. 16, p. 54. Berkeley. Type at Kew.

SACLEUXII—Africa.—Jour. de Bot., vol. 6, p. 19. Hariot. Type in museum at Paris.

SCUTIGERA—Brazil.—Elenchus Fung., vol. 1, p. 73 (as Polyporus). Fries. No type exists.

SEURATI—Pacific Islands.—Bull. Soc. Myc. France, vol. 22, p. 48. Patouillard. Type in his herbarium.

SIMILIS—Australia.—Hooker's Jour., 1846, p. 4. Berkeley. Type at Kew.

SPECIOSA—Africa.—Fungi Natalensis, p. 137. Fries. Type in a jar in museum at Upsala.

- SUBTENUIS**—India.—Not previously published. Berkeley. Type at Kew.
- SULCATA**—Ceylon.—Hooker's Jour., 1847, p. 510. Berkeley. There is no type at Kew, but a co-type in Montagne's herbarium, also one in that of Fries.
- TENUIS**—South America.—Kunth. Synopsis, vol. 1, p. 10 (as Boletus). Hooker. Type at Kew.
- UMBRINELLA**—Africa.—Fungi Natalensis, p. 137. Fries. Type in museum at Upsala.
- VARIEGATA**—Central America.—Proc. Amer. Acad., vol. 4, p. 122. Berkeley. There is no type so labeled, but there is no question that it is a specimen labeled *Hexagona papyracea* at Kew.
- VELUTINA**—Africa. Bull. Soc. Myc. France, vol. 9, p. 209. Patouillard. Type in museum at Paris.
- VESPACEA**—East Indies.—Voyage de Uranie, p. 170. Persoon. Only known from one little type in museum at Paris, another in Persoon's herbarium at Leiden.

APPENDIX II.

GEOGRAPHICAL DISTRIBUTION.

The real study of mycology, as I view it, is the study of species and their variations and the geographical distribution. Very little can be told of the geographical distribution at present, for much more abundant material will have to reach Europe before anything definite can be determined. We have arranged in the following tables the species found in the museums under nine geographical divisions, as follows:

- No. 1, United States and Canada.
- No. 2, Mexico, Central America, and West Indies.
- No. 3, South America.
- No. 4, Europe.
- No. 5, Africa.
- No. 6, India, Ceylon, and Malay Peninsula.
- No. 7, Japan and China.
- No. 8, Philippines and East Indies.
- No. 9, Australia, New Zealand, New Guinea, and Pacific Islands.

The sign † indicates localities from which we have seen one or but a few specimens; if a number of collections it is indicated by a heavy face **C**; if the species is only known from the type collection, we indicate that fact with a star (*). Species that are very closely related and perhaps better called sub-species are indicated by being indented under what we consider the "type" form.

	UNITED STATES.	WEST INDIES, ETC.	SOUTH AMERICA.	EUROPE	AFRICA	INDIA, ETC.	JAPAN & CHINA	EAST INDIES, ETC.	AUSTRALIA, ETC.
Group 1, Setosus.									
<i>Apiaria</i> ,		†						C	C
<i>Deschampsii</i> ,								C	
<i>hirta</i> ,						C			
<i>capillacea</i> ,									
<i>aculeata</i> ,			* *						
<i>elegans</i> (probably),									
<i>Henschalli</i> ,								*	
<i>Dybowski</i> ,									

	UNITED STATES.	WEST INDIES, ETC.	SOUTH AMERICA.	EUROPE.	AFRICA.	INDIA, ETC.	JAPAN & CHINA.	EAST INDIES, ETC.	AUSTRALIA, ETC.
Group 2, Velutinus.									
variegata,	C	C	.	.	†	.	.	.
scutigera,	C	C	.	.	†	.	.	.
velutina,
Group 3, Ungulaformis.									
nitida,	†	†	.	.	.	†
Gunnii,	*	.	.	.
sulcata,	†	.	†	.
durissima,	*	.
resinosa,
Group 4, Applanatus.									
Pobeguini,	C
niam-niamensis,	*
leprosa,	*
speciosa,	*
chartacea,	*
Kurzii,	*	†
erubescens,	*
amplexens,
Group 5, Tenuis.									
tenuis,	C	C	.	C	C	.	C	C
pulchella,	†	.
polygramma,	C	C	.	C	C	.	C	.
umbrinella,	C	C	.	.	.
discopoda,	C	C	.	.	.
subtenuis,	*	.	.	.
phaeophora,	†	.	.	†
rigida,	C
similis,	*
atrosanguinea,	*	.	.	.
Sacleuxii,	†
Group 6, Pallidus.									
albida,	C
macrotrema,	†	.	.	†	†
vespacea,	*	.
ochroleuca,	C	.	C	C
Seurati,	*
aequalis,	*
rhombipora,	*
Group 7, Pseudofavolus.									
cucullatus,	†	†
Miquelii,	†	†	†	.
bipindiensis,	†	.	†	.	.
mirabilis,	*
Group 8, Resupinatus.									
heteropora,	*

APPENDIX III.

SYNONYMS AND SPECIES IMPERFECTLY KNOWN.

The following is the list of specific names which we would refer to synonymy and our reasons for the same. We give also the countries from whence proposed, and the individuals responsible for them. We hold them responsible who published them, though in some cases the names were taken from and credit given to manuscript names. We also indicate a few manuscript names under which specimens are labeled in our principal museums. While of course a question in synonymy is largely a question of individual opinion, the following list (except in such cases as specially stated) is our conclusions as to authentic specimens examined. We have studied in the British Museum, the Museums of Kew, Paris, Berlin, and Upsala, which list embraces all the museums of Europe where much historic material is preserved, except the Java specimens at Leiden. We visited Leiden twice for this purpose, but both times found the Java specimens had been loaned. We would not pretend to publish as synonyms (as has recently been done) names that we have merely copied from others, nor would we perpetrate the fraud of pretending to pass upon specimens we never saw, and which in many cases do not exist. It is a fact well known to those who have investigated the subject that the usual description is a mere empty form. Plants can be recognized from systematic work in which those of a section or country are described by contrast, but it is impossible to describe a specimen as an isolated fact so that it can be surely recognized in one case out of a hundred. If the labels were removed from the type specimens in the museums I believe that not ten per cent of them could ever be replaced from anything that has been published about them, and I doubt if one per cent could. Under these conditions I feel it is useless to carry in our literature names and descriptions of specimens that do not exist. It is a part of the system of "science" to pretend to be able to judge from these descriptions as to the identity of the plants described, but I do not think that any one who has had experience really believes it (except in very exceptional cases), and I decline to subscribe to any such fiction.

If a plant has not acquired a name by use, or if it was not characteristically illustrated, and if authentic material does not exist in some museum or where it can be examined, there is little occasion to further encumber literature with it.

While the following list is specific names of plants placed in *Hexagona*, it does not follow they were all so placed by the authors stated. This may have been done by some one else, and who it was is immaterial and not worth recording. Nor does it follow that the species are all invalid in other genera where they belong, but not in my opinion in *Hexagona*.

Where we state "no type exists," we have been unable to find the type in the museum where it should be preserved, or authentic material in any other museum. We have made careful and systematic search, and taken time, and we believe the statements are literally true. Still we are aware there is always the possibility of the type turning up in some obscure place. Often we have found historic specimens in drawers or in cupboards, where the casual visitor would never think of looking.

The following is the list that we would refer to synonymy, and the reasons. We state also the name of the country whence described, and the names of the discoverers of these "new species." It is remarkable how many "discoveries" are made in "science," chiefly noteworthy from the fact that they are not true.

adelphica, Africa, Cooke=*Hexagona hirta*.

adnata, Ceylon, Berkeley=*an anomaly of some kind*.

affinis, Pacific Islands, (Published?) Berkeley=*Hexagona tenuis*.

arata, Pacific Islands, Berkeley. It is not a *Hexagona*, but a *Polyporus* related to *gilvus*.

auriculata, United States, Patouillard. Specimen not found for me, but I have little doubt it is *Hexagona cucullata*, which is the only species we have in the United States, I think.

Bartlettii, South America (Published?), Masee. Better classed as a ferruginous *Poria* related to *contigua*.

Blumei, Java, L veill . No type known to me.

Boneana, Africa, Patouillard. Too close to *umbrinella*.

brevis, Ceylon, Berkeley. No type exists.

Burchelli, Mss.=*umbrinella*.

carbonaria, United States, Berkeley. Better classed as a ferruginous *Poria*, close to *contigua*.

Casuarinae, New Caledonia, Patouillard=*Hexagona tenuis*.

cervino-plumbea, Java, Junghuhn=*Hexagona tenuis*.

Cesatii, Borneo, Cesati=*Hexagona albida*, a little cyclomycooid.

ciliata, Philippines, Klotzsch=*Polystictus versatilis*.

cingulata, West Indies, L veill . No type known to me.

crinigera, Africa, Fries=*Hexagona hirta*.

cladophora, Philippines, Berkeley. Not a *Hexagona* for me. A better *Trametes*.

Cookei, New Guinea, Saccardo. Change of *Hexagona favoloides* of Cooke, which being *Hexagona albida*, the change was not necessary.

coriacea, Brazil, Berkeley. Type inadequate to judge.

crassa, Africa, L veill =*Hexagona hirta*.

cruenta, South America, Montagne=*Trametes Persoonii*.

cyclophora, African island, L veill . No type exists.

decipiens, Australia, Berkeley. For me not a *Hexagona*. It has colored spores and is a better *Polyporus*.

dermatodes, Philippines, L veill . It is a *Polystictus-Trametes*.

discolor, Australia, Fries. No type exists.

Dregeana, Africa, L veill =*Hexagona umbrinella*.

fasciata, Pacific Islands, Berkeley. No type exists.

favoloides, Central America, Peck=*Hexagona tenuis*.

favoloides, New Guinea, Cooke=*Hexagona albida*.

favus, China, Linnaeus. No type exists. Supposed to be *Hexagona apiaria*.

flabelliformis, Philippines, Berkeley. Type material inadequate.

Friesiana, South America, Spegazzini=*Polystictus villosus*.

glabra, Africa, Palisot. No type is said to exist, but there is a good picture. Probably the same as *Hexagona umbrinella*.

glabra, India, L veill . A hexagonal form of *Lenzites ochroleucus*.

gracilis, Brazil, Berkeley. Belongs to a section of *Polyporus*.

inconcinna, Philippines, Berkeley (as *Daedalea*)=*Hexagona albida*.

induratus, West Indies, Berkeley=*Hexagona Miquelii*.

intermedia, Australia, Berkeley (as *Daedalea*)=*Hexagona albida*.

Klotzschii, Africa, Berkeley=*Hexagona hirta*.

Koenigii, Ceylon, Berkeley=*effete Hexagona apiaria*.

laevis, Pacific Island, Cooke, nondescript.

lurida, Java, L veill =*Hexagona glabra*.

Marcucciana, Italy, Baglietto=*Hexagona nitida*.

Molkenboeri, Java, L veill =*Hexagona macrotrema*, and based on same collection.

Mori, Italy, Marcucci (as *Favolus*)=*Hexagona nitida*.

Muelleri, Australia, Berkeley. Too close to *Hexagona rigida*.
nigro-cincta, Pacific Island, Patouillard=pale form of *rigida*.
orbiculata, Africa, Fries=*Hexagona tenuis*.
obversa, Africa, Patouillard. Too close to *chartacea*.
pallens, Mexico, Saccardo. Unknown to me.
pallida, African Islands, Schröter. Unknown to me.
papyracea, locality unknown, Berkeley=*Hexagona variegata*.
peltata, Africa, Fries. No type exists.
pergamenea, Ceylon, Berkeley. Not a *Hexagona*. Close to *Polystictus dermatodes*.

picta, East Indies, Berkeley. Type inadequate.
pustulosus, Java, Lévillé=*Hexagona Miquelii*.
sericea, United States, Fries=*Polystictus villosus*.
sericeo-hirsuta, United States, Klotzsch (as *Polyporus*)=*Polystictus villosus*.
sinensis, Africa, Klotzsch=*Hexagona hirta*.
sinensis, China, Fries. In reality merely a change of name of *Boletus favus*, of which no type exists. Fries states "v. s.," but the specimen he saw was from Klotzsch and was *Hexagona hirta*.

strigosa, Africa, (Mss. name) Cooke=*Hexagona hirta*. It was published as *Trametes adelphica*.

Stuhlmanni, Africa, Hennings=*Hexagona Pobeguini*.

subaculeata, Borneo, Cesati. Unknown to me.

subrigida, Philippines, Murrill. Unknown to me.

tabacina, Java, Lévillé Not a *Hexagona*, but the same as *Polystictus cichoriaceus*. Lévillé also discovered the same plant was another "new species" in another genus and called it *Polyporus fuscus*, but little matters of this kind did not bother Lévillé.

Taxodii, United States, Murrill=*Hexagona cucullata*, teste the author.

Thollonis, Africa, Patouillard. Unknown to me. Type is at Brussels, but not seen by me.

Thwaitesii, Japan, Berkeley=*Hexagona tenuis*. Thwaite collected in Ceylon, and had nothing to do with this plant from the island of Bonin. Why it was named after him I do not know.

tricolor, Africa, Fries. No type exists. From the description I think it is the same as *Hexagona discopoda*.

unicolor, Africa, Fries. No type exists.

velutina, China (published?), Patouillard=*Hexagona bipindiensis*.

versicolor (ascribed to Fries). No such plant was named or published, which, however, did not prevent Spegazzini from so determining specimens.

vitellina, Borneo, Cesati. Unknown to me, but I have not much faith in there being a yellow *Hexagona*.

vittata, Central America, Ellis=*Polystictus villosus*.

Welwitschii, Africa, Smith=*Hexagona Pobeguini*.

Wightii, India, Klotzsch=*Hexagona apiaria*, but in the museums mostly known under this name.

Note.—There is a variation in the spelling of the generic name. Some spell it *Hexagona*, others *Hexagonia*.

INDEX TO THE SPECIES OF HEXAGONA.

Those marked with a star (*) are perhaps better called sub-species or varieties or are of minor importance.

Aculeata	9	Miquelii	36
Aequalis*	33	Mirabilis	37
Albida	29	Niam-niamensis	18
Amplexens	22	Nitida	14
Apiaria	6	Ochroleuca	31
Atrosanguinea	27	Phaeophora*	26
Bipindiensis	36	Pobeguini	17
Capillacea	8	Polygramma*	25
Chartacea	20	Pulchella*	25
Concinna*	26	Resinosa	16
Cucullata	35	Rhombipora*	34
Deschampsii	9	Rigida	26
Discopoda*	26	Sacleuxii	27
Durissima*	16	Scutigera*	13
Dybowski	11	Seurati	33
Elegans	9	Similis*	27
Erubescens	21	Speciosa	21
Gunnii	15	Subtenuis*	26
Henschalli	11	Sulcata	15
Heteropora	39	Tenuis	23
Hirta	7	Umbrinella*	26
Kurzii	21	Variegata	12
Leprosa	20	Velutina*	14
Macrotrema*	30	Vespacea*	33

QK
600
-L79

BULLETIN No. 20.

1912.

MYCOLOGICAL SERIES, No. 6.

BULLETIN
of the
LLOYD LIBRARY
of
**BOTANY, PHARMACY AND
MATERIA MEDICA**

J. U. & C. G. LLOYD
CINCINNATI, OHIO

MYCOLOGICAL SERIES, No. 6

Synopsis
of the
Stipitate Polyporoids

By C. G. LLOYD



REV. G. BRESADOLA.

Who has in my opinion, the best critical knowledge of foreign Polyporoids and to whom I am indebted for many determinations and advice, I beg to dedicate this pamphlet in appreciation of the many kindnesses received from him.—C. G. LLOYD.

THE STIPITATE POLYPOROIDS.

The subject of the polyporoids is quite extensive, embracing as it does about three thousand alleged species. We have been engaged in the study now two or three years, but except in a general way have not been able to cover the entire field. We would divide them roughly into about five divisions as follows:

- 1st, The stipitate species, embraced in this pamphlet.
- 2d, The genus *Fomes*.
- 3d, The genus *Polyporus*, sessile section.
- 4th, The genus *Polystictus*, sessile section.
- 5th, The allied genera such as *Favolus*, *Laschia*, etc.

During the past two or three seasons we have visited all the museums of Europe and America where most of the historical material is preserved, and have made our studies, notes, and photographs of the type specimens. This embraces the museum at Kew, the British Museum at London, the museums of cryptogamic botany at Paris, Leiden, Berlin, and Upsala. We have looked over a small collection at Copenhagen, and some of the specimens in the private collection of Professor Patouillard at Paris. We did not have time to thoroughly work over Patouillard's species while in Paris, hence a number of his species are unknown to us. In America in our own collection we have more American specimens than there are in all the other museums combined. We have thoroughly studied the collection of Professor Peck and the specimens of Schweinitz at Philadelphia. The New York specimens we have not seen, as on both of our visits to New York Mr. Murrill was absent and we did not wish to work with his material in his absence.

Our final work on this pamphlet was done at Kew, during February, March, and April, 1911. There is no other institution in the world where one can work to such advantage as at Kew, where there is not only the largest collection of historical specimens, but the most perfect library, and where the conveniences are best.

The conclusions recorded in this pamphlet are our own, made on examination and study of authentic material. We have not indulged in the too common practice of passing on species we never saw. We may be mistaken in some of our opinions, but we have perpetrated no fraud. When we record a species as unknown, this means of course that it is unknown to us. In a few instances where we have not seen specimens we have adopted the opinion of the Rev. Bresadola, but in each such case this is distinctly stated.

The first and we think the best division of the pore species was made by Fries (1851) in his *Novae Symbolae*. At that time but relatively few plants were known, but Fries' divisions were based in the main on the most prominent characters, and of the eleven sections into which we have divided the stipitate species, nine of them have been taken mostly in their original signification from Fries' work. Professor Patouillard has outlined a plan of division which we think on the whole is not as good as that of Fries, but it embraced a few new ideas and two of them, the sections *Ganodermus* and *Amaurodermus* we have adopted.

In addition there has been no lack of men who have amused themselves by inventing new names for the sections of polypores. In the start we have Karsten who was the first to engage in such work. He discovered that most of Fries' sections were "new genera" and gave them names. The work had so little merit and had evidently so little originality as a whole that although proposed thirty years ago, no one except the author has followed it since, and it figures, when it has figured at all, chiefly in synonymy.

Monsieur Quélet, a leading French mycologist, learned the greater part of what he knew from Fries and his works, and in his first publication could not find words to express his appreciation of the "grand mycologue d'Upsala." After he had gotten a little insight into the subject he passed the latter part of his life juggling the names of his great master, and he did it so thoroughly that very few of his colleagues, even in France, have ever been disposed to use his work. This is unfortunate, for Quélet was a field mycologist and knew well the species that occur in France. As far as I have been able to decide there was no system or logic to his juggling, his only object apparently being to propose names in place of Fries' names.

Schroeter would divide the Polyporus species into three genera on the color of the spores and context, which while answering very well for the few species that he knew, if generally applied would bring the bulk of them, about a thousand, into one genus.

The last man to engage in this line of name changing is Mr. Murrill, who has no more trouble discovering "new genera" and concocting new names than if there had not been three men doing exactly the same thing with the same plants before him. I question if there is an institution or mycologist in Europe that attaches any importance or pays any attention to this kind of work, and very few in America. In my opinion such work is of little value or avail.

The principal work that these men do is to get up new "generic" names on various pretexts, and of course one can make a "genus" out of every species if he wants to. Their chief work, however, is to take the old sections of Fries' genera and then juggle up excuses to give them new names usually under the cover of some "rule." Such work, in my opinion, has so little to commend it that I do not consider it worth citing in detail even as synonyms.

When Fries proposed the divisions of the subject he knew but very few species, but in the years that have followed "new species" have been published in quantities, chiefly by Berkeley, who proposed so many of them that no one has been able to do much with them since. He was not, however, the only one to name foreign species, although he named a large part of them. Twenty-five per cent of the species considered good in this pamphlet were named by Berkeley.

In the early days Klotzsch and Junghuhn named quite a number. Then came Fries, Montagne, and Berkeley. Then Lévillé and Cooke, and Kalchbrenner. In the latter years we have Hennings, Patouillard, and Murrill. It would be more accurate to state that they named collections, for I do not think that any of them knew much about

what the others had done, and it has been very much of a haphazard proceeding from the beginning.

Junghuhn and Montagne, I think, did the best work, or rather the specimens they left are the best. Fries' foreign specimens have largely disappeared and many of them will never be known. Kalchbrenner did the worst work of anybody and renamed as "new species" the commonest, old, well-known things. He did not seem to have had the most elementary idea of the subject. L veill 's work was in the main very poor, and Murrill's recent work is almost as bad. Berkeley, Hennings, and Patouillard have named as new a large part of the collections that come into their hands. Naturally they got a number that are good, and many that I think are not. Spegazzini grinds the "new species" out by the wholesale from South America, but very few of his specimens reach Europe, and such as have are largely misnamed. I think no one knows what he is doing, not even he.

Very little can be told from any "description" that can be drawn from a Polyporus, and the most of the determinations that are made from "descriptions" are wrong. The only way to get names for the plants is to hunt them up in the various museums where they are preserved, and then it is often not satisfactory. One finds the same thing named over and over again. Names based on little frustules that never did give the slightest idea of any character and many other irregular things that would not be tolerated except in "Science." I believe Bresadola to be the only man in Europe who has made an earnest effort to hunt up and learn the characters of the "old species" of Polyporus in the various museums of Europe to-day. I do not always agree with him in all the details, but I think no two who endeavor to learn names for fungi from the fragmentary, indefinite, and conflicting specimens on which the names have been based will ever agree in all cases.

Cooke tried to arrange the names according to the Friesian system, but owing to the number of species and the hurried manner in which the work was done, it was very inaccurate and in its details was most erroneously done. This was not all Cooke's fault. Many of the "new species" are described in such a way that not only can nothing be told about their identity, but in many cases from the description one can not even place them in the section where they belong. In this pamphlet, when species stated to be unknown (to me) are placed in sections, I do not claim that such disposition is anything more than a guess.

Having nothing else to follow, Saccardo adopted Cooke's arrangement, which is quite unfortunate, as Saccardo is used as a basis of classification in most museums, and by this method species are brought into the same division that have little resemblance and often no relation.

In this pamphlet the stipitate species are divided into eleven sections, or genera if one so desires to call them, but we prefer to call them sections. We disclaim having discovered any "new genera" or anything else new in the classification. Nine of our divisions we have taken from the work of Fries and two from that of Patouillard. If

we have succeeded in arranging the species where they belong in these sections that will be enough "novelty" to satisfy us, for we think it has not been even approximately done before.

As this pamphlet is proposed simply as an arrangement of the species, we have given but very brief descriptions, in fact only the more salient points. We think very few species are ever learned except from specimens, and that the largest part of the bad Latin that is used in describing species is purely a waste of good printer's ink and of no avail whatever as far as identifying the species is concerned. We have introduced a number of photographs that will be found to be of more service in this respect than the most minute descriptions that could be written. We have not given in detail the source of these illustrations, but we believe them to all be true to name, and the greater part of them are made from the type specimens.

As to nomenclature we have employed the sectional name as the first binomial (except in one case where it would produce the barbarism called tautology) and these sectional names are all old and familiar and will not lead to any confusion. If these sectional names are taken as genera, it is absolutely senseless to record who used the sectional name first as a generic name for any particular species. Mycologists are so very busy recording in great detail *who* did this and *who* did that, and *who* called it this and *who* called it that, and *who* made this combination and *who* made that combination, that they often have little time left to consider *what* was done.

As to specific names, in the body of this work we have not added personal names to the specific names, believing that in the case of most of the plants the authors being dead, it would not serve the usual purpose of ministering to self-conceit. We have given them in our synonyms where we think they are quite appropriate. We have given these names also in our index, according to the wishes of the authors in most cases, although not all. Some writers are so selfish they wish to advertise only themselves, others divide the advertisement with a collector or with a friend. It has been suggested that it would be a gracious thing to give all the advertisement to the collector, and I think the latter is the best plan, at least I adopt it in this pamphlet where I am concerned. In several cases in arranging the species it occurs that sometimes two in the same section have the same specific name. We have made no change, merely indicating the second by the word *bis* and would prefer to leave the work of changing names to others. We have endeavored to make this pamphlet a practical summary of what is known (to us) on the subject, and have indicated by the size of the type the relative value of the species as they appeal to us. Those printed in the larger type are the leading marked, characteristic species which we believe have merit and value. Those in smaller type are forms or doubtful species or plants imperfectly known to us. Where I have not seen and studied a species I usually place it in the section "unknown" (to me) for I think there is nothing gained by my guessing a second time concerning what was in many cases largely an original guess. If I have done any guessing as to the identity of

those specimens I have not seen, I have plainly so indicated it in each instance. The eleven sections in which the species are divided being mostly the old, familiar sections, need little explanation. The two that are least known are the sections *Ganodermus* and *Amaurodermus*, which at our hands have undergone changes of gender in order to be uniform with the others.

The section *Ganodermus* was first proposed for the common *Polyporus lucidus* of Europe. There are but few species in temperate regions, but it is more common in the tropics. The main characters are usually a strongly laccate surface, colored context, and the real character is a spore character. The spore (Fig. 395) has a hyaline membrane or episore which is large and projects at the base beyond the colored endospore. This empty base usually collapses, then the spore becomes truncate at the base. It has been stated that this is not the base but the apex of the spore, a statement I do not believe.

Amaurodermus is a tropical section. All species have stems, usually central but also lateral. The surface is generally dull and the stems velutinate. The spores (Fig. 396) usually in abundance, are colored, globose or oblong, large, and the endospore fills the episore. The apiculus is rarely prominent. We have included in this section several thin species in which we have not found the spores and which we doubt really belong to it.

The other sections that we adopt are the well-known sections of Fries' system that need no special explanation other than our key. Some species present characters that would place them in two sections. In such cases we use our own judgment in placing them where we think they best belong. The names for the sections are mostly the same that Fries used. In one case, *Perennis*, we use another name, *Pelloporus*, for reasons we have previously stated, viz., the plants are not perennial.



Fig. 395.




Fig. 396.

KEY TO THE SECTIONS OF STIPITATE POLYPOROIDS.

There are included here only the stipitate species of the old genus *Polyporus*. The allied genera as *Favolus*, *Laschia*, etc., are not here considered.

Sub-woody.

With woody fibrils but not perennial and not having the pores in strata (except as to the first).

Pores in areas of growth (indistinctly stratified) *Fomes*.

Pores Not Stratified.

Spores colored (mostly elliptical) with a strong apiculus. Context colored. Surface of most species laccate. (Cfr. also p. 99).... *Ganodermus*.

Spores colored, mostly globose, with none or a small apiculus. Context colored. Surface of most species dull. (See p. 99)..... *Amaurodermus*.

Spores white. Context (except sec. 11) pale or white *Lignosus*.

Fleshy or Coriaceous.

Stipe lateral (Spores white) *Petaloides*.

Stipe branching and bearing several pileoli *Merismus*.

Stipe Central or Excentric (Rarely Lateral).

Flesh spongy, light (Spores white or colored) *Spongiosus*.

Spores colored. Fleshy or coriaceous *Pelloporus*.

Spores white. Fleshy, soft, usually terrestrial, with thick pilei *Ovinus*.

Spores white. Fleshy-pliant, coriaceous, usually thin pilei, and epixy-
lous *Lentus*.

Lentus with black stems *Melanopus*.

SECTION FOMES (STIPITATE.)

Although the first sixty-one species placed in *Fomes* in Saccardo (Vol. 6) have stems, I believe there is but one of them that can be so included on the definition there given and generally accepted for this genus, viz.: "perennis, successive strata nova gerens." Many are subligneous in texture, but are annuals in temperate regions, and in the tropics if they persist more than one season (which is doubtful) they do not produce successive pore strata. The following is the only one in which I have noticed the slightest indication of strata.

DIABOLICUS (Fig. 397).—The entire plant (except the pores and context) is black. Stem mesopodal, with pale, solid context, and black, smooth, dull crust (1-1½ x 8-10 cm.). Pileus (8-14 cm.) black, even, depressed in the center, with black, smooth, dull surface. Context (5 mm.) pale cinnamon color. Pores minute, at first cinnamon, but black when old. If not in layers at least in distinct areas of growth. Colored setae very abundant on the hymenium. Spores not found, doubtless white. This is a rigid, black plant, growing on wood in Brazil. It is known only from Spruce's original collection. In its context color, setae, and spores (probably) it is related to *Fomes pomaceus*, but there is no other similar stipitate species known.



Fig 397
Fomes diabolicus.
(Reduced)

SECTION GANODERMUS.

The section Ganodermus is characterized by peculiar spores (see page 99) and also usually the stipe and pileus are laccate (viz.: covered with a dark, resinous, shining crust). The context and spores are colored.

2. SPORES SMOOTH OR BUT SLIGHTLY ROUGH.

LUCIDUS.—Stipe and pileus strongly laccate. Context cinnamon or fulvous, varying lighter. Pores not stratified. Spores 6×10 , slightly rough. A strongly marked species of Europe and America and its forms (?) are found in the tropics. It is difficult to draw the line as to the tropical forms, although I am disposed to refer to lucidus all those with the same stem insertion and similar context color. The stipe is usually pleuopodal, rarely mesopodal, but the pileus is never in my opinion sessile. The three following I think are but forms of lucidus.

VALESIACUS.—Only a form with paler context, and not really a form at that, for lucidus varies much as to context color and is never very dark. The common American plant that corresponds to this European form has been called Ganodermus Thugae.

JAPONICUS.—Europeans usually refer the Japanese form to lucidus. In the Japanese lists it figures as Polyporus Japonicus. I think both are right. Forms that I have seen from Japan are blacker than the European plant but are surely the same species.

LAUTERBACHII.—A thin, tropical form of lucidus. It seems thinner and more rigid, but for me it is only a form.

OTHER FORMS.—Plants received from India I would refer to lucidus as forms. They are not so strongly laccate, more dull, and often mesopodal.

INCRUSTANS.—We have in the United States a curious form, or perhaps an abnormality, of lucidus which instead of taking the usual shape with a lateral stem is often thin, cup-shaped, with an indistinct stem. It has usually been referred to lucidus, but is quite different in its habits. It grows usually in grassy places, incrusting the blades of grass.

CURTISII.—Context, spores, and stipe as Polyporus lucidus, but not strongly laccate and color is yellowish. I have collections which are pale, almost white. It is a plant of a southern type in the United States, frequent in the South and extends up the coast to New Jersey and is rarely found in New England. In the museums of Europe there are several specimens exactly the same, from Africa.

AMBOINENSIS.—This is based on an old figure by Rumphius. Many specimens so named are in the museums, but I have seen but one that resembles the figure. This is a Philippine collection in the British Museum. The stem attachment is like that of fornicatus, but the stem is very slender and the plant appears to have grown erect as Rumphius shows it. The stem is not branched as originally shown, otherwise the specimen corresponds exactly. This was the first foreign species in this section to be illustrated. Fries referred to it pictures that appear

SECTION GANODERMUS.

quite different. The many specimens so named in the museums often have little resemblance to each other, or to the original picture.

COCHLEAR.—Plant with a black, laccate crust. Stipe thick (about an inch) 6-10 inches long, dorsally attached. Context cinnamon. Pores minute with white mouths. Spores 8×14 , smooth or slightly rough. This is a common species in Java and the East Indies, but we have seen no specimen from any other section. There are a number of collections at Leiden, and it was sent to us abundantly by Dr. Königsberger from Java. We take it in the sense of Bresadola's determination at Leiden, though we doubt if it is the same as Nees illustrated, especially as to the stipe. The plant has the same (*dorsal*) stem insertion as *fornicatus* and *amboinensis*, but much more obese stem.

AFRICANUS.—Pileus thick, obese, with a mesopodal, obese stem. Context dark, umber. Surface dull, resinous. Spores 7×10 , minutely rough. In its relationship, color of context, and spores this plant is close to the sessile species such as *applanatus* and widely departs from all others in this stipitate section. The type came from South Africa and was misnamed *Polyporus Umbraculum* by Kalchbrenner and fragments were so distributed (de Thümen, 708). It seems to have the same color characters as *fulvellus*, which is a sessile species, and the *exsiccatae* number was cited under that species.



Fig. 398
Ganoderma fornicatus.



Fig. 399.
Ganoderma Lingua.

SECTION GANODERMUS.

FORNICATUS (Fig. 398).—Pileus and stipe with black, laccate crust. Stipe slender, dorsally-lateral, attached. Context thin, cinnamon, fulvous. Pores *minute, hard, compact*. Mouths at first white (contrary to description), then *purplish* brown. Spores 6×10 , smooth. No type exists, but it is frequent in Brazil, the "type locality," according to numerous collections of Spruce (No. 48, 79, 172). It is characterized by the peculiar stipe attachment. In Australia are similar plants, but the spores are rougher. There is also a similar plant common in Ceylon (teste Petch), but I have seen no specimens. Specimens from New Caledonia determined as *amboinensis* I take to be the same.

MASTOPORUS.—Stipe thick, lateral, with a smooth laccate crust. Context thin, cinnamon, scanty. Pores hard, *minute, compact*, dark purplish brown. Spores 5×8 , smooth. Very similar to *fornicatus* as to the peculiar hard pores. Type from Singapore at Paris, but it comes to me frequently from Africa and is probably common throughout the East.

FLEXIPES (bis).—Pileus unilateral, attached, small (1-2 cm.) with a strongly laccate, black crust. Stipe slender, cylindrical, with a smooth strongly laccate, black crust. Pores small, pale cinnamon. Spores 5×10 , smooth. Known from one collection, from China, in the herbarium of Patouillard. It differs from all others in this group in its slender stem and habits. It has a general resemblance (except small pores) to our figure (411) of *Polyporus longipes*.

LINGUA (Fig. 399).—Pileus small, rarely more than two or three cm. wide, but deep (2-3 cm.) in proportion to its width. Attached by a dorsal-lateral, short stem. Surface dark reddish brown, laccate, sulcate. Pores long, reaching the crust, small. Context cinnamon. Not common in the museums, but specimens seen from Java, Sumatra, and New Guinea. Known from its small size and peculiar shape. Type has not been seen. We take the species in the sense of Montagne's determination. It does not exactly correspond to the original illustration, but we have seen no specimen that does.

BONINENSIS.—Stipe dorsally prolonged. Surface dull, ferruginous or cinnamon, not laccate. Context dark tabacinus. Spores 6×12 , smooth. Known only from the type at Paris from Bonin Island, collected by Wright and distributed (U. S. Expl. Exp.) as *Polyporus lucidus*. The corresponding collection at Kew is a different (laccate) plant.

REGULICOLOR.—Surface dull, reddish brown, not laccate. Stipe lateral, apparently proceeding from a rhizome or a rooting stem. Known from a single specimen at Kew stated to be from Cuba, but I think the locality is doubtful.

SECTION GANODERMUS.

3. SPORES DISTINCTLY ROUGH.

OCHROLACCATUS (Fig. 400).—Pileus small but deep, attached by a short rudimentary, dorsal stem. Crust *pale, ochraceous*, faintly laccate, rugulose, zoned. Pores medium with white mouths, long, not stratified but reaching the crust, very regular, arranged in lines. Spores *large*, 16×32 (!), with small apiculus, distinctly rough. A strongly distinct species, very rare and known only from the Philippines. Types at Paris and at Kew and the British Museum. These collections which are surely the same species vary some in external appearance. The type at Paris is our figure 400. That at Kew is almost white with a dull surface. That in the British Museum is sessile and has a pale, smooth, shiny crust as if waxed but not laccate.



Fig. 400

Ganoderma ochrolaccatus (pores enlarged X6).

PLACOPUS.—Pileus with an intense *black, shining, laccate* surface, becoming dull in old specimens. Stipe lateral, with similar crust. A small species, thin, an inch or two in diameter. Spores 8×12 , distinctly rough. Only known to me from Bresadola's naming from Java at Leiden. As I have found no types in any of the museums, I judge his determination was made from the description only.

EMINI (Fig. 402).—Pileus small, usually pleuropodal, rarely mesopodal. Stipe long, with black, strongly laccate, smooth crust, rooting at the base. Spores large with distinct apiculus 20×28 , rough. The pileus is not as strongly laccate as the stipe. A marked species known from abundant types at Berlin from Africa.

HENNINGSII (Fig. 401).—Pileus and stem strongly laccate, smooth, dark. Pileus 3-4 inches in diameter with a mesopodal, rooting stem. Pores small, pale. Spores 10×12 , rough, subglobose, but distinctly apiculate. Known from one collection at Berlin from Africa, made by Stuhlman and confused by Hennings with the preceding.

SECTION GANODERMUS.



Fig. 401

Ganoderms Henningsii.




Fig. 402

Ganoderms Emini.

OPACUS.—Pileus 3-4 cm. x 5-10 mm. thick, with a fragile, dull, brownish crust. Stipe mesopodal with similar crust. Context *pale* cinnamon, thin. Pores minute with concolorous mouths, 4 mm. long. Spores 8-10 oval, with small, hyaline apiculus and are *strongly rough*. This is known from two collections at Paris, one from Brazil, the other from Cuba.

SECTION GANODERMUS.

ALLUANDI.—Pileus with a smooth, black crust. Stem 1 cm. thick, 15 cm. long, laterally (dorsally) attached to the pileus, smooth, black crust. Pores small, round, some large and sinuate, with thin walls and concolorous mouths, long, reaching the crust. Context scanty, cinnamon. Spores 10 x 16-18 with a distinct apiculus and distinctly rough. Known from a single specimen at Paris (in the cupboard) from Africa.

UNNAMED.—Pileus with a thin crust, mat, minutely velvety, with a few darker, slightly metallic zones. Context very thin, pale cinnamon. Pores 1 cm. long, *minute*, pale cinnamon with concolorous mouths. Stipe mesopodal, 24 cm. long, sulcate, with sterile branches, covered with a smooth, black crust. Spores *strongly reticulate* (the only reticulate polyporoid spores known to me) obovate with small, apiculate base, 12 x 20, pale colored. Type found by me unnamed, without label, in a cupboard in the Museum at Paris, the origin unknown but probably from Africa. I do not name it as I presume they will wish to do so at Paris.

HILDEBRANDI.—Pileus, context, and stem exactly the same as the small form (*ramosii*) of *Polyporus rugosus*. Spores conidial, ovoid, 4-5 x 5-7, distinctly rough. Known from one specimen at Paris. I suspect it is a conidial form of *Polyporus Ramosii*.

4. ANOMALOUS SECTION WITH A FALSE STEM.

PISACHAPANI.—This is, I judge, an anomalous species. It is flat, branched like the fingers of a hand, and the stem is made of discs growing from each other, as if the plant started to produce a succession of pilei and then changed its mind and produced a false stem. The surface is smooth, laccate. Nees named and figured it from Java. I found a single specimen of this curious growth from Samoa. In my specimen the pores are not perfectly formed and I find no spores.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

I doubt if a more cumbersome, inaccurate, or impractical system could be devised for the naming of plants than the one that has been adopted by "Science" in the naming of fungi. The European work of Persoon and Fries was based for the most part on a practical knowledge that they had of the growing plants, and the greater part of their work was of the highest merit and will always stand. The only weak part is the species that were founded on old pictures, often inaccurate and erroneous, and the names were based often on the inaccuracies of the pictures. For many of them no plant is known that corresponds.

As to foreign (to Europe) species the whole subject has been a haphazard proceeding from the start. The earlier namers had very scanty material, but they based a "new species" on almost every collection that they received, and many of them were evidently but slight varieties or individual forms to which the same authors would have paid no attention had they seen the forms growing in their woods. As the years rolled by new "authorities" came into the field and each one has discovered a large part of the plants he got from foreign countries to be "new species" and gave them names, although not one of them, I think (except Bresadola), has made any serious endeavor to learn the names that others have given to largely the same plants.

SECTION GANODERMUS.

The result is a mess of about 3,000 names of Polyporei mostly compiled into Saccardo to date, and no one can tell anything whatever as to their identity from what has been written about them. The only way to learn the names is to hunt them up in the museums where they are preserved, and when they are not preserved, and many of them are not, nothing will ever be known about them. This hunting up process is rather difficult and ordinarily is not possible.

Of the 3,000 "species" of Polyporei that have (mostly) been scraped up into Saccardo I doubt if one-fourth of them represent anything of value, and it is more trouble to learn which are of value than the subject is worth. If I had spent the same time and work on something useful, that I have working over these old puzzles, I do not doubt that a great deal more good could have been accomplished. But with me it has been a recreation and a pleasure that the subject would probably not have yielded had it been exploited in an intelligent manner, and had anything definite been known about it.

There are about 700 alleged species (names) that are supposed to be stipitate and considered in this pamphlet. Of these I have seen about 500 authentic specimens, and of those seen 225 impress me as being good species and having merit and 58 others have been retained as having some possible value. The others I have seen to the number of about 215 chiefly reflect I believe the lack of knowledge or judgment on the part of the authors. In addition there are 65 stipitate species (dead) carried in our literature of which no authentic material can be found. Nothing will ever be really known about any of them, though it is the fashion nowadays for tourists to make running visits to the various museums and come home and tell just as big yarns about those that do not exist as they tell about those that do. As long as they can arrange a lot of Latin names in a row and give the dates, it seems to be immaterial whether there is any truth in the arrangement or not. There are 105 alleged species marked in this pamphlet unknown (to me) that do not exist in any of the principal museums. They may be found in some out of the way museum or private collection, though I doubt if they are worth the trouble to look them up. Still I presume they have the same possible value as those that are in the museums as they are all a very uncertain quantity.

The species of fungi are relatively few and widely distributed, a fact that is becoming more firmly fixed every day. What constitutes a species, however, can not be defined by words. It is only a matter of experience and individual opinions. The question of variation which is a large factor in the truth of the problem is hardly taken into account at all by the promoters of "new species." And it is a question of course in which there is room for many differences of opinion.

In the following lists of synonyms we have given our opinions of the specimens we have seen. We do not do it with the idea that it will settle the questions in any way, for that is impossible. "Science" nowadays consists in raking over these old "synonyms" and arranging them chronologically, and we suppose this process will be continued to the end of time. A "new species" is like a spot of ink. It may not have the slightest merit or value, and may be based solely on the fact that its author was not informed on his subject, but there is no way to ever get rid of it.

albo-cinctus, Congo, Patouillard. Unknown to me. Seems to have the stipe attachment of fornicatus.

argillaceus, Cuba, Murrill. Unknown.

asperulatus, Philippines, Murrill. Unknown.

avellaneus, Central America, Murrill. Unknown.

coffeatus, West Indies, Berkeley. Type is a few fragments from which nothing can be learned.

Currani, Philippines, Murrill. Unknown.

declivis, Pacific Island, Kalchbrenner. Unknown to me, but the description reads like fornicatus.

flaviporus, West Indies, Murrill. Unknown.

SECTION GANODERMUS.

formosissimus, South America, Spegazzini. Only known to me from Rick's determination, which (teste Bresadola) is the same as renidens.

Haenslerianus, New Zealand, Hennings. No specimen found by me at Berlin.

incrustatus, Central America, Fries. No type exists.

Javanicus Java, L veill . Type at Leiden in very bad condition, but I think belongs to the section Ganodermus and has no relation whatever to Polyporus varius, of which it was given as a "variety" by L veill .

neglectus, Central America, Patouillard. Type is a mere fragment from which little can be told. The species was based on large, globose spores, which are not the normal spores of the plant. The basidial spores are typically those of the section Ganodermus (not Amaurodermus, as named).

nutans, Central America, Fries. No type exists and its identity is unknown. The determinations at Paris, Berlin, and London are all different from each other and all are probably wrong. Murrill's elaborate account was only worked up from Fries. He tells "spores not examined," which was not strange as he never saw an authentic specimen, and I do not see how he could have examined the spores of a specimen he never saw.

perzonatus, Cuba, Murrill. Unknown.

Pes-simiae, Brazil, Berkeley. No type exists. From the description it seems to be Pisachapani.

praelongus, Cuba, Murrill. Unknown.

pulverulentus, West Indies, Murrill. Unknown.

stipitatus, Central America, Murrill. Unknown.

subamboinensis, Brazil, Hennings. Same as Lauterbachii, and both are but tropical forms of lucidus.

subformicatus, Central America, Murrill. Unknown.

subincrustatus, West Indies, Murrill. Unknown.

Tsugae, United States, Murrill. Same as the common Polyporus lucidus of Europe, the distinction given being that it has paler context and that Polyporus lucidus has "one to many-layered strata varying in distinctness," all of which was chiefly imagination on the part of the author. Polyporus lucidus is an annual and never has strata of pores, though as it has been called "Fomes" Mr. Murrill was undoubtedly right in thinking if it did not have strata it ought to have. As to the paler context the same form had been named Valesiacus in Europe, but it is not even a distinct form of lucidus.

SECTION AMAURODERMUS.

The section Amaurodermus is quite close to the preceding section, but is distinguished by large, globose, oval, colored spores, which usually have no distinct apiculus. All are stipitate plants with usually dull (not laccate) surface and often velutinate stems. Species rarely have smooth, laccate stems. Context and hyphae are colored. All are plants of the warm countries, no species being known in temperate regions. (Cfr. also p. 99).

5. POLYPORUS. SPORES SMOOTH OR BUT SLIGHTLY ROUGH. STEM SLENDER, USUALLY MESOPODAL.

RUGOSUS.—Pileus dark brown, rugulose with a dull, mat surface. Stem olive brown with a dull, minutely velutinate surface. Context pale cinnamon, when freshly cut it turns reddish. Pores small with thin walls. Spores globose, smooth, 6-8 or 8-10. This is quite a common species in the East and numerous specimens from Java and Ceylon are at Kew. I have not seen the type but Nees gave a good figure of it, and specimens from Ceylon (Thwaite, No. 728) exactly



Fig. 403

Amaurodermus rudis



Fig. 404

Amaurodermus Sprucei with enlargement of pore mouths.

SECTION AMAURODERMUS.

accord with this figure. The fresh plant when bruised turns dark and herbarium specimens are usually black.

RAMOSII.—Bresadola endorses this as a synonym for rugosus, and I think it is a slender form. The spores and other characters are in the main the same, but the plants are more slender and the context thinner. It occurs over the same regions as rugosus and also the Philippines.

RUDIS (Fig. 403).—Pileus strongly rugulose with mat, dull surface, minutely velutinate, light in color. Stems with olive, velutinate surface. Pores medium, with thin walls. Context light cinnamon. Spores globose, 9-12, with thick walls, minutely rough. The type of rudis I have not found, but there are abundant collections so named by Berkeley from Australia, where it seems to be common. It is close (too close perhaps) to rugosus of the East, but seems to be more rugulose, has larger pores and spores, and when mature retains its color. Young specimens, however, turn black in drying.

SPRUCEI (Fig. 404).—(Change of Porothelium rugosum of Berkeley.) Pileus dark brown, rugulose, with narrow, concentric zones. Surface mat. Stipe pleuropodal with mat surface, concolorous. Pores and context pale, the pore mouths pustular, hence put in the genus Porothelium (sic) when originally named. Spores globose, 8 mic., smooth, very pale. Known only from the (abundant) types collected by Spruce in Brazil. It departs from others of this section in its spores and context being paler.

VARIABILIS (Fig. 405).—Pileus from 2 to 6 cm. broad and about 5 mm. thick, with a lateral, slender stipe. Color pale alutaceous, both pileus and context. Surface dull. Spores 9 x 12, oval, smooth. This is quite a distinct species, characterized by its pale color and oval spores. It is known from two collections (Nos. 57 and 183 part) made by Spruce in Brazil. It is badly named for it is quite uniform, but there was confused (and figured) with it quite a different plant (cfr. Polyporus unilaterus, in the next section).

CALCIGENUS.—Pileus about an inch in diameter, with a reddish brown, laccate crust. Context pale olive. Stem mesopodal, slender, with brown, mat surface. Spores abundant, oval, large, 12 x 16, deeply colored, smooth. Quite distinct but known from a single specimen at Kew from Spruce, Brazil.

RIVULOSUS.—Pileus glabrous, rugulose (not rivulose, I think), reddish brown with paler margin. Stem pleuropodal, branched, sometimes bearing two pilei, with a smooth, dark reddish crust. Context thin, ligneous, pale cinnamon. Pores minute, pale but darker than the context. Spores globose, 14 mic. faintly reticulate, with thick walls. Known only from the type in the Museum at Paris from Java. It has the general appearance of a Ganodermus, but from its spore characters belongs to Amaurodermus.

SECTION AMAURODERMUS.



Fig. 405
Amaurodermus variabilis.



Fig. 406
Amaurodermus Chaperi.
Reduced one-half.

CHAPERI (Fig. 406).—Pileus 15 cm. broad, with a smooth, dark crust, but not laccate. Stem hollow, with sterile branches, with smooth, pale grayish surface. Pores minute, short with dark mouths. Spores globose, smooth, or minutely punctate, 8-12. Known from a single and quite old specimen in a cupboard at Paris. It was first referred (through error) to *Polyporus scleropodius* of Africa, then described and named as above.

INTERMEDIUS.—Pileus soft, velutinate (now wrinkled) black. Stipe mesopodal with a hard, black, smooth, laccate crust. Spores abundant, globose, smooth, apiculate, 12 mic. Known from a single specimen kept in a drawer at Berlin, from Africa. It was originally sent in glycerin and is still soft and sticky.

SECTION AMAURODERMUS.



Fig 407

Amaurodermus Auriscalpium.



Fig 408

Amaurodermus praetervisus.




Fig. 409

Amaurodermus omphalodes.

AURISCALPIUM (Fig. 407).—Pileus reniform, the upper surface rugulose, zonate, mat. Stipe lateral with a dull crust. Color reddish brown. Spores 6-8, globose, smooth, pale colored. The original Persoonian specimen from Brazil is still well preserved at Paris. Montagne referred here several collections from South America which Berkeley and Patouillard have held to be different.

PRAETERVISUS (Fig. 408).—This is based on a single specimen from Brazil, referred to the preceding by Montagne. As to shape, size, context, color, and spores it is very much the same, but it has larger pores and a thicker, blacker, harder crust.

OMPHALODES (Fig. 409).—Pileus orbicular, with a mesopodal stem, but rarely perfect, being usually more or less lobed and divided to the stem. Surface glabrous but not laccate, rugulose, more or less zoned. Stipe slender with a mat, dull surface. Pores small, 2 mm. deep. Spores globose, 12 mic., smooth, pale. Abundant specimens were sent to Kew from Spruce, Brazil. All have very much the same general size and stature as our figure. At Paris are much larger, obese (but otherwise apparently the same) specimens from South America, called "var. fulvaster."

SECTION AMAURODERMUS.

6. POLYPORUS. SPORES DISTINCTLY ROUGH. STEM
SLENDER, USUALLY MESOPODAL.

a. STIPE SMOOTH, NOT LACCATE SURFACE.

ANGUSTUS (Fig. 410).—Pileus large (10 inches in diameter) with glabrous, strongly rugulose, zoned, dull, grayish brown, not laccate, crust. Stipe mesopodal, an inch thick, with spongy context and hard, smooth, gray, not laccate crust. Pores minute, soft, *pale isabel-*



Fig. 410

Amaurodermus angustus. (Reduced more than one-half).

SECTION AMAURODERMUS.

line, as is the context. Spores (only conidial, I think) globose, colored, strongly tubercular, rough, 8 mic. A remarkable species known from one specimen at Kew collected by Spruce in Brazil. It is the largest, mesopodal polyporoid I have noted and grew on wood. I suspect that its normal spores would be found to be quite different, and that the plant is not well classed in this section. No other plant in this section, I believe, grows on wood, and most of them have subterranean rhizomes.

b. STIPE WITH A SMOOTH, LACCATE CRUST.

LEPTOPUS.—Pileus 5 cm. broad, 1½ cm. thick, with a dark, smooth crust. Stem, almost gone now, but enough remains to show that it was mesopodal, about 12 cm. long, one cm. thick, and had a dark, smooth, shiny, laccate crust. Context pale cinnamon. Pores small, about one cm. long, pale cinnamon. Spores globose, 12 mic., strongly rough. The species was referred by Fries, who never saw it, to *umbraculus*, of which no type exists, and I think there are no grounds for accepting the reference. It was so accepted by Patouillard, however, who drew his characters from Persoon's specimens and knew nothing whatever about Fries' plant. The species is only known from the original, Persoonian specimen, preserved at Paris. It was from the island of Rawak.

LONGIPES (Fig. 411).—Pileus unilateral, attached, with a reddish brown crust which is not polished (laccate) as the stipe. Stipe slender, with a black, shiny, laccate crust. Pores large, *pale* cinnamon, reaching the crust. Spores unique, globose, 12-14 mic., strongly rough, having the *asperities arranged in distinct bands or areas*. This is a very peculiar species, known only from one quite abundant collection from French Guiana. Collector unknown. Leprieur, who made large collections from the same locality, never found it.

RENIDENS.—Pileus dull reddish brown. Stipe lateral, smooth, laccate crust. Context scanty, cinnamon. Pores and pore mouths concolorous. Spores globose, 8-9, rough. Known to me only from the type at Berlin, from Brazil, collected by Moeller. (Plants distributed by Rick as *formosissimus* are said to be the same. I have not examined their spores.) Except as to the spores this species has the general appearance and character of *Polyporus lucidus*.

BASILAPIDOIDES (as *Laccocephalum*).—Pileus brownish fawn, with strongly pitted surface. Context whitish. Stem short, thick, mesopodal, forming at the base a large, hard, false sclerotium, consisting of agglutinated grains of sand fixed by the mycelium. Spores globose, orange yellow, echinulate, "44-50 in." (mic.?) in diameter. This Australian species, called the "stone making fungus," is only known in Europe from the description and figures in an Australian publication. It was proposed as a "new genus," but I judge from its spore characters it should be classed here.

SECTION AMAURODERMUS.



Fig. 411
Amaurodermus longipes.



Fig. 412
Amaurodermus insularis.




Fig. 413
Amaurodermus unilateralis.

SECTION AMAURODERMUS.

c. STIPE WITH DULL, MINUTELY VELUTINATE SURFACE.

UNILATERUS (Fig. 413).—Pileus small (1-1½ cm.) reddish brown, dull surface, unilaterally attached. Stem slender (1½-2 mm. x 7-9 cm.) with dull surface. Pores minute, 5-8 mm. deep, pale cinnamon with white mouths. Spores large, globose (or subglobose) 20 mic., distinctly rough. The types at Kew (Spruce, No. 207) from Brazil were named by Berkeley in manuscript "ellipticus," but when published they were included and figured as part of *Polyporus variabilis*. They differ from *variabilis* not only in different stem insertion but have very different spores.

FASCICULATUS.—Pileus unicolorous, pale fauve in some specimens, fuliginous in others, marked with prominent, raised, narrow, concentric zones. Context cinnamon. Stem with dull, velutinate surface, light brown color. Pores minute, 3-4 mm. deep, darker color than the context, the mouths stuffed, isabelline. Spores subglobose, 12-14, strongly rough, pale colored. A strongly marked species known only from two collections, both from Congo, Africa. The original is in the herbarium of Patouillard at Paris, others sent me by Édouard Luja, Congo Belge. A character of both of these collections is that each pileus is borne on two or more distinct stems, or perhaps the pilei of two or more stems are consolidated into one, but they do not seem to have that appearance.

INSULARIS (Fig. 412).—Pileus 3 cm. with a strongly wrinkled, dull, mat surface. Pores large, pale cinnamon, in the "type" mostly torn and destroyed. Stipe mesopodal with mat, finely velutinate surface. Spores large, oval, 12 x 16, minutely but distinctly rough. Known from a single specimen at Paris from New Caledonia.

7. POLYSTICTUS, PLANTS WITH THIN PILEI AND PORE LAYERS.

Spores of some species said to be globose, colored, but I have rarely found them, and hence can not state from my own knowledge. Context and pores colored, brown. Hyphae colored. I suspect some at least have hyaline spores.

a. PORES LARGE.

GRACILIS (Fig. 414).—Pileus lateral (or unilateral) thin, dark reddish brown, with dull surface. Stipe slender (1-2 mm. thick by 5-15 cm. long) with a dull surface, proceeding from a rhizome or buried rootstalk. Pores *large*, 1 mm. in diameter, round or hexagonal. Spores not found by me. This is a unique species only known from the original collection, Spruce, Brazil. It was classed by Berkeley in the genus *Hexagona* where it really belongs on its pore characters alone. However, there is no other similar species in the genus *Hexagona*, and in its habits, context, surface, also spores probably, it is evidently so close to this section *Amaurodermus* that it should be placed here.

SECTION AMAURODERMUS.




Fig. 414

Amaurodermus gracilis. (One specimen enlarged X6).



Fig. 415

Amaurodermus ocellatus.



Fig. 416

Amaurodermus Schomburghii

SECTION AMAURODERMUS.

b. PORES SMALL.

OCELLATUS (Fig. 415).—Pileus thin but *rigid*. Surface smooth, faintly zoned. Pores very minute, rigid with thick walls. The colored hyphae have thick walls so that a cross section appears like thickened cells. Stipe central or lateral, slender, mat, light brown. Spores not found. Known only from Brazil collections (Spruce). The pores are so minute that they are hardly visible, but it is all a mistake that “the pore mouths are contracted, etc.”

MACER.—I did not cut the single specimen that represents it (Spruce, Brazil) but as to the pileus and pores it seems to me the same as the preceding. The lateral stem, however, is blacker and I think it a different species. The determinations, “macer, Berkeley,” both at Berlin and Paris have no resemblance to it.

SCHOMBURGKII (Fig. 416).—Pileus mesopodal or pleuropodal, thin with zonate, smooth surface. Pores concolorous, minute. Stipe dull, slender. Named from specimens from British Guiana in Hooker's herbarium, but afterwards found by Spruce in Brazil.



Fig 417
Amaurodermus sericatus.
119

SECTION AMAURODERMUS.

SERICATUS (Fig. 417).—Pileus thin, seal-brown, depressed in the center, coriaceous, with a smooth, minutely velvety, shiny, satiny surface. Stipe mesopodal, long (15-20 cm.), slender (3-5 mm. thick) with dark, dull velvety surface. Pores minute ($1-1\frac{1}{2}$ mm. deep) clear seal-brown, with concolorous mouths. Hyphae deeply colored. Setae none. Spores not found, probably white. A single specimen is at Kew, collected by J. H. Holland at Old Calabar, Africa, and referred by error to rugosus.

HETEROMORPHUS.—Pileus depressed in the center, brown, sub-zonate. Pores with white mouths, at least when young. Stipe mesopodal, with dull surface. The types both at Paris and at Kew are young collections. Several specimens in the museums I think are referred here in error.

RENATUS.—Pileus thin, reniform, with a lateral stipe, dark reddish brown. Pores minute, white then brown. Stipe slender ($1\frac{1}{2}$ -2 mm.) with dull surface. Known from Spruce's collection in Brazil. Ellis also determined it from Nicaragua and I think correctly. It seems to me very close to the next, but is larger and pleuropodal.




Fig. 418

Amaurodermus exilis.

Fig. 419

Amaurodermus marasmioides.

SECTION AMAURODERMUS.

JURIENSIS.—Pileus thin, rigid with a minutely velutinate, rugulose, dark, zonate pileus. Pores minute, white, bruising brownish. Stipe mesopodal, slender with dark, dull, velutinate surface. Spores globose, pale colored, $3\frac{1}{2}$ -4 mic. Collected in Brazil and called *Polystictus sacer* var. *juriensis* by Hennings. It has no resemblance or remote relationship even to *Polyporus sacer*.

EXILIS (Fig. 418).—Pileus thin, mostly mesopodal, with smooth, faintly zonate surface, small, 1-1½ cm. Pores minute. Stipe filiform (1 mm. thick) long, wiry. Known from Spruce's collections from Brazil. Placed in *Fomes* (sic) by Cooke.

MARASMIOIDES (Fig. 419).—The smallest species in this section, but with the exception of size it has the same characters as *exilis*, and I think is only a small *Polyporus exilis*. The specimens sent to Paris by Berkeley have a little filiform stem, not over 2 cm. long, with pilei about ½ a cm.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

auriscalpioides, Brazil, Hennings=*auriscalpium*.

bataanensis, Philippines, Murrill. Unknown.

boleticus (Bull. Soc. Myc.) misprint for *boleticeps*.

Boleticeps, South America, Patouillard. Unknown to me except from illustration (Bull. Soc. Myc. France, 1888, pl. 12). Seems close to *omphalodes* and came from same region.

cassiaeolor, Brazil, Berkeley. A single specimen so named which, though thick, I believe to be a thick specimen of *Schomburgkii*.

Clemensiae, Philippines, Murrill. Unknown.

Elmerianus, Philippines, Murrill. Unknown.

nigripes, Brazil, Fries. No type exists. Unknown. The description reads much like *leptopus*.

Pala, South America, Léveillé. Unknown. No type exists.

Parmula, Brazil, Berkeley=*exilis*.

passerinus, Brazil, Berkeley=*renatus*.

procerus, Brazil, Berkeley. Only two specimens so named, both immature. One specimen has quite a long stipe. I think both are heteromorphous.

pulcher, Africa, Fries. No type exists. Figure (Afz. 19) seems to be in this section although it has a laccate stipe.

pullatus, China, Cooke. This is a manuscript name that Berkeley gave to an old specimen from Hong Kong, but afterward concluded that it was *rudis* of Australia and did not publish it. Cooke afterward dug it up and published it. I do not think the old specimen is *rudis*, but it was too poor to publish.

rufobadius, South America, Patouillard. Unknown to me except from illustration (Bull. Soc. Myc. France, 1889, pl. 10). Seems to me to be too close to *omphalodes*.

rugosus, Berkeley, Brazil (as *Porothelium*). The specific name being a duplicate, was changed to *Spraguei*.

subrenatus, Central America, Murrill. Unknown.

subrugosus, Samoa, Bresadola=*rugosus*.

Umbraculum, Africa, Fries. No type exists. Unknown. Used by Patouillard as a substitute for *leptopus* of Persoon which was not justifiable as he knew what *leptopus* was and did not know as to *Umbraculum*. Specimen determined by Kalchbrenner and distributed (de Thümen 708) has no possible resemblance to Fries' description or the figure cited.

xylodes, Brazil, Berkeley=*Schomburgkii*.

SECTION LIGNOSUS.

This section embraces stipitate species, that are subligneous but not perennial. In texture they are similar to the preceding. They are never soft and fleshy. The hyphae and spores are pale, in which characters they differ from the two preceding sections. Most of them are included in *Fomes* in Saccardo, but none of them are *Fomes* according to the definition of the genus that Saccardo gives.

8. PLANTS WHICH FORM A SCLEROTIUM. CONTEXT PALE OR ISABELLINE. SPORES PROBABLY WHITE.

SACER (Fig. 420).—Pileus thin, with minutely velutinate, zonate surface. Color pale to dark brown. Stipe mesopodal, dull, pale surface, proceeding from an underground sclerotium. Context and pores isabelline. Pores medium small with thin walls. *Polyporus sacer* is represented in the museums by a number of collections, all from Africa. It first reached Fries and was of much interest from the fact of having a sclerotium. The name "sacer" refers to some superstition that the negroes are said to attach to it.

RHINOCEROTIS.—Pileus glabrous, rugulose zoned, at first thin, then thicker and indurated. Stipe mesopodal with a dull surface, not "laccate" as erroneously described, proceeding from a sclerotium two or three inches in diameter. Context pale. Pores *minute*. This plant is so close to *Polyporus sacer* that our photograph (Fig. 420) could well represent either. It is quite different however in its *minute* pores, and the tissue of old specimens becomes more thick, hard, and woody. It was known for many years only from the imperfect type from Malay, but recently a fine specimen was sent to Kew from Perak, and Professor Petch has made one collection in Ceylon.

9. PILEI UNILATERAL AND SUPERIMPOSED. CONTEXT PALE.

SUPERPOSITUS (Fig. 421).—Pileus unilateral and superimposed, arranged one above the other or on one side of stem, like shelves. Surface pale isabelline, smooth. Context pale isabelline. Hyphae pale. Spores not found, doubtless white. This species is most curious in the peculiar arrangement of the pilei. It is known from three collections, all at Kew. First it reached Berkeley from "New England," Australia (not "Amer. Bor." as Saccardo incorrectly compiles it) then Cooke got a collection from Perak and one from New Guinea. These three from widely remote localities are all I have seen of this most peculiar species.

10. STIPE MESOPODAL OR PLEUROPODAL. CONTEXT WHITE OR PALE. SPORES WHITE.

CORRUGIS (Fig. 423).—Pileus with pleuropodal (rarely mesopodal) stipe. Surface of pileus and stipe minutely velvety-brown with

SECTION LIGNOSUS.



Fig. 420

Lignosus sacer, the plant (reduced) and pore surface (natural size).

faint, metallic zones. Context white. Pores medium, firm. This is a rare species of Europe known only from the Alps and rare there. I have seen but three collections. First, at Berlin, collected by Morthier at Neuchatel and determined as "triqueter, Fr." Second, sent to me by Dr. Butignot, Switzerland, and third, in the herbarium of Boudier, also from Dr. Butignot (and labeled *Trametes Butignotii*, not published?). In its context, color, and pores it closely resembles *Polyporus benzoinus* (a sessile species). It was named *Polyporus rugosus* by Trog, then changed to *Polyporus Trogii* by Fries (Cfr. Sacc. 6, 82) but when Fries published it, Hym. Eur. p. 536, he called it *Polyporus corrugis*. No type exists in Fries' herbarium, nor I am told in that of Trog.

SECTION LIGNOSUS.



Fig 421

Lignosus superpositus.

PREUSSII.—Pileus with a dark, nearly black, rugulose surface. Stipe mesopodal. Context and pores pale isabelline, rigid. Pores minute. Spores not found, but I think are white. Known from one specimen at Berlin from Africa. The photograph would closely resemble *Polyporus rugosus*, but the plant is quite different. It is close to the preceding if not the same. It was named by Hennings as *Ganoderma*, but does not belong to that section though it might be called a *Trametes*.

DEALBATUS (Fig. 422).—Pileus lateral or unilateral with a stem 3-8 mm. thick. Surface pale, smooth, dull with a minutely velutinate coat (compared in its naming to a coat of whitewash). Context pale. Spores globose or compressed-globose, 5-6, smooth. This

SECTION LIGNOSUS.



Fig. 422
Lignosus dealbatus.



Fig. 423
Lignosus corrugis.

species has been badly confused (see page 190). It was originally collected by Ravenel in South Carolina and Curtis also found it in North Carolina, but I think no specimens exist except these original types. It is found in Saccardo (page 159) as *Fomes* and also (page 218) as *Polystictus*, and it is neither.

PANSUS.—Pileus rugulose, with a dull surface, strongly zoned with brown and darker zones. Stipe mesopodal with a mat, dull surface. Context thin, almost none, pale. Pores minute, $1\frac{1}{2}$ -2 mm. long, isabelline color which may be due to age, as it is described as white. Spores abundant, globose, 8 mic., pale colored, which may also be due to age. This is a strongly marked species of which several specimens have reached Kew from northern South America.

DUBIOPANSUS.—This has all the characters of the preceding except that the pores have *orange mouths*. This is quite uniform and appears to me a natural

SECTION LIGNOSUS.

color. I am told that it is caused by a Hypomyces, but I am unable to detect the mycelial threads of a parasite in the tissue and it does not seem to explain it to me. The spores which I think are conidial are subglobose, hyaline, apiculate, and distinctly rough. I have a specimen from L. Damazio, Brazil, and there is one at Kew from Georgetown, British Guiana.

PAULENSIS.—Known by a single specimen from Brazil at Berlin. If not the same it is quite close to pansus. It has a well developed, ligneous, white context and *hyaline* globose, 6-7, smooth spores. Otherwise it seems the same as pansus, particularly in its peculiar, zoned surface.

HYPOPLASTUS.—Surface dark, almost black, faintly zonate. Stipe black, smooth, with a resinous crust. Context pale isabelline. Spores not found, but I think they are white. The type is a mesopodal plant, but Berkeley refers here (and I think correctly) two flabelliform specimens. All are from northern South America. This plant differs from all others in this section in its laccate stem. While I have found no spores, I believe it does not belong to the section Ganodermus.

CAMERARIUS.—Pileus reniform, smooth, even, glabrous, beautifully zoned with narrow, regular, concentric, brown zones. Stipe is pleuropodal (in one specimen, probably the same, it is mesopodal) with a dull crust. Context pale isabelline, probably white when fresh, 2-3 mm. deep, almost reaching the crust. Spores not found, probably white. Several specimens of this are at Kew, all from northern South America.

ARENATUS.—Pileus flabelliform, subligneous, incurved in drying. Surface gray, strongly zoned. Context white. Stipe lateral, short, thick. Pores minute, rigid, pale. A strongly marked species from New Guinea found in the Museum at Paris.

RHIZOMATOPHORUS.—Pileus flabelliform, thin, with smooth, pale isabelline surface. Pores minute, concolorous. Stipe slender, long, attached to a slender, long rhizome. A single specimen of this is at Berlin, from Brazil. It is endorsed “=Trametes Rhizophorae” which is surely an error.

PUDENS.—Known from a single, young, half specimen at Kew, from India. It has a long rhizome and in some respects it resembles the preceding. I think not much can be ascertained from this single, immature type, but it may be recognized through comparison if found again.

POLYDACTYLUS.—This is known from one apparently abnormal specimen from Brazil. It has white context and a lateral stipe which divides and bears on the ends of the branches little, orbicular, disc-like pilei. The surface is minutely velutinate, brown, and marked with metallic zones. In its general nature I think it is related to corrugis of Europe.

ATRO-PURPUREUS.—This is also known from a single specimen from Brazil, and has the same context color and surface marking as the preceding. The pilei are thinner and borne in a different manner. The pore mouths are white, but when bruised are reddish. I think the plant is badly named.

SECTION LIGNOSUS.

II. CONTEXT BROWN OR GILVUS. SPORES WHITE
(PROBABLY.)

a. STIPE NOT BLACK.

BRUNNEO-PICTUS.—Pileus suborbicular or reniform with a smooth, brown zoned surface. Context *brown*, hard, with hyphae deep yellow under the glass. Pores minute with pale mouths but brown context. Stipe lateral, hard, with dull brown surface. Spores not found, but I think without question are white. This is a rigid, well-marked species, known from several collections of Spruce, all from Brazil. When young it is thin, but becomes thicker with age. The thin, young specimens contract strongly in drying and were called *Polyporus semiclausus* by Berkeley.



Fig. 424

Lignosus Zambesianus.
(Top of Pileus)

SECTION LIGNOSUS.

ZAMBESIANUS (Fig. 424).—Pileus strongly rugulose with a distinct, dull crust, brown, zonate. Context gilvus, rigid. Stipe mesopodal (but not preserved with the type). Pores small, 8 mm. deep, gilvus context and brown mouths. Hyphae bright yellow under the microscope. Setae none. Spores not found, but doubtless white. This is known from a single specimen, preserved at Kew and collected in Zambesi in 1881. It was misreferred to rudis. It is the only mesopodal polyporoid I ever saw with *gilvus* context.



Fig. 425

Lignosus scopulosus.

b. STIPE BLACK—MELANOPUS.

SCOPULOSUS (Fig. 425).—Pileus with a smooth, pale, thin crust. Stipe lateral, black, with a black crust. Context punky, isabelline, with slender, pale hyphae. Pores minute, isabelline, with concolorous mouths. Spores hyaline. This is a frequent plant in the East and has been well illustrated by Reichardt under the name *Trametes Rhizophorae*, under which name it has been well known to me for a number of years. It is a noteworthy plant with its black stem, and smooth, pale pileus. It is given in Saccardo as a *Fomes*, but is a ligneous *Polyporus* or might be classed as a *Trametes*.

SECTION LIGNOSUS.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

- Butignoti, Europe, Boudier (as *Trametes*, published?)=*corrugis*.
canalium, China, Loureiro. Too ancient and vague to be entitled to a place even in synonymy. "Described" 120 years ago and never seen since. Said to be "white and viscid."
- hemibaphus, Brazil, Berkeley=*obese camerarius*.
obsoletus, Brazil, Fries. No type exists. I judge it is similar to *brunneo-pictus*.
pallidus, Brazil, Berkeley=*camerarius*.
Rhizophorae, East Indies, Reichardt (as *Trametes*)=*scopulosus*. It was well illustrated and has generally been called under this name.
rhodophaeus, Java, L veill . This was described as having a short, lateral stipe (or sessile). The type specimen (at Leiden) is sessile and is now referred to *semilaccatus*, which I believe always a sessile species.
scleropodius, Africa, L veill =*sacer*.
semiclausus, Brazil, Berkeley=*brunneo pictus*.
Trogii, Europe, Fries. This was a name proposed for *rugosus* of Trog, but when Fries published it he called it *Polyporus corrugis*.
triqueter in the sense of Qu let is *corrugis*. Not the same as in the sense of Romell and Bresadola.

SECTION PETALOIDES.

We include in this section most species that have lateral stipes, embracing for the most part the divisions *Petaloides* of *Polyporus* and *Discipedes* of *Polystictus* as found in Saccardo. The more woody species (*Lignosus*) are classed in the preceding sections. Also in the section *Melanopus* are found those species with black, lateral stems.



Fig. 426

Petaloides hirtus.

SECTION PETALOIDES.

12. **CARNOSUS. FLESHY, SOFT, THICK SPECIES.**

HIRTUS (Fig. 426).—Surface brown, velutinate or hirsute. Flesh white. Stipe lateral. Spores peculiar, *fusiform*, $6 \times 14-16$, hyaline, smooth. A very rare plant both in Europe and the United States.

AQUOSUS.—This has a strong, lateral stipe. Flesh white, soft, watery, thick, drying thin. Pores small, white. Known from two collections, both from Brazil, also one (?) from India.

RUTROSUS.—A good picture of a petaloid, fleshy, white species with medium large pores, published by Rostkovius seventy years ago. Found or imagined in Germany, but known to no one since.



Fig. 427

Petaloides fusco-maculatus

FUSCO-MACULATUS (Fig. 427).—Flesh soft, watery, not “papyraceus-membranaceous” but so drying. Surface spotted with minute spots. Pores large. Spores oblong, $3-4 \times 8-10$. Found by me in Samoa. Has no relation to squamosus to which it was compared.

GLUTINIFER.—Known from a single, sliced specimen at Kew, and is I think probably the same as the preceding. It is said to have come from Mauritius, but it was more probably from Australia.

Osseus. See Addenda, page 191.

SECTION PETALOIDES.

13. POLYPORUS. FLESHY, THIN SPECIES, COLOR WHITE OR PALE. PORES SMALL.

ANNULATUS (Fig. 428).—A small, white species with a short stipe expanded into a disc at the base. Originally from Java (well illustrated by Junghuhn). Found by me in Samoa. It grows attached to sticks on the ground.

RHIPIDIUM.—Stipe lateral, expanding above. Color white (?) when fresh, reddish when dried. Pores medium with thin walls. The type form is rare in the United States. It is large, but otherwise the same as the little form, called pusillus by Persoon, which occurs frequently throughout the tropical world.

FRACTIPES.—White, with a lateral, white stipe. Pores small, slightly rose colored. Surface dull. Rare in the United States. Also found in Brazil, by Rev. J. Rick.



Fig. 428

Petaloides annulatus.



Fig. 429

Petaloides nivicolor.



Fig. 430

Petaloides biokoensis.

NIVICOLOR (Fig. 429).—Pure white, with smooth surface. Pores small, white. Not truly stipitate, but the pileus extends behind into a stipe-like prolongation. Known only from New Zealand, but there are abundant specimens at Kew.

BIOKOENSIS (Fig. 430).—Pileus clear yellow when fresh, bleaching out to white in drying. Surface smooth, faintly zoned. Stipe short, lateral, concolorous. Pores minute, yellow when fresh, isabelline when dry. Spores (conidial?) globose, hyaline, smooth, 4-5 mic. The plant contracts in drying and the color change is unusual and marked. Very rare in Samoa, collected by me but once. Named by Bresadola. Type unknown to me. (Published?)

SECTION PETALOIDES.

PENETRALIS.—Pileus spathulate, tapering to a long stem. Color pale. Pores small. Grew on stem of tree ferns in greenhouse in England. No type in the museums, but it is well illustrated. (Jour. Bot., 1875, t. 162.)

CANDIDUS.—Pure white with a lateral stipe. Pores small. Seems to be well illustrated (Persoon, Myc. Europe, t. 15), but is unknown from specimens in any museums or recent collection. Referred as being a form of *chioneus* by Fries, but it is surely not.

OBLIQUUS.—Context pale, darker in drying. Pores $1\frac{1}{2}$ -2 mm. long. Pileus with a long, lateral stem. Known from one collection, New Guinea, at Kew. With the exception of long pores it has a resemblance to *obovatus*.

14. POLYPORUS. FLESHY, THIN SPECIES, COLOR WHITE OR PALE. PORES LARGE, FAVOLOID.

JANSEANUS (Fig. 431).—Pileus thin, fleshy, pure white, tapering to a long stem. Pores large, favoloid. Known from one collection from Java, preserved in alcohol at Berlin.



Fig. 431

Petaloides Janseanus.



Fig. 432

Petaloides brachyporus.



BRACHYPORUS (Fig. 432).—Pileus thin, tapering to a lateral stipe. Pores large, *shallow*. Dried specimens dark and brittle, but I judge white or pale when fresh. Originally from French Guiana, in the herbarium of Montagne. There is also a collection at Berlin from Brazil, which Hennings has named as a “new species,” but I have mislaid my memorandum as to what he called it.

SECTION PETALOIDES.

15. **POLYPORUS. THIN, RIGID. COLOR PALE ROSE OR REDDISH BROWN. NOT ZONATE OR ONLY FAINTLY ZONATE.**

MODESTUS (Fig. 433).—Color when fresh pale cinnamon or rose, becoming in old specimens reddish brown. Pores minute. Surface dull. Rarely distinctly stiped, but reduced to a short stipe-like base. Appears to be frequent in tropical America and usually named by Berkeley *albo-cervinus*.

RUBIDUS.—This from the East is close to *modestus*, and I know of no marked difference. It is thicker, not so spatulate, and has but a faint indication of a stipe. I am told by Professor Petch that when old, dark discolored patches usually appear on the top. It is common to Ceylon and I think in other parts of the East.



Fig 433

Petaloides modestus.



Fig 434

Petaloides Didrichsenii.

BRUNNEOLUS.—The best specimen is in the British Museum. Those at Kew are poor. It is quite close to *rubidus*, but the context and general color are more brown. It seems to be common in the Philippines, and in recent determinations under the name *atypus* is confused with *rubidus*.

PETALODES.—Surface dark reddish brown with appressed fibrils. Context pale. Pores minute. Stipe lateral, thick. Known from a single specimen at Kew, collected in Brazil by Spruce.

DIDRICHSENII (Fig. 434).—Very similar to *modestus*, but with distinctly larger pores. Only type known, from Borabora (Society Is.), is at Kew. It seems to be frequent in the East and was received abundantly from Ceylon and called *Menziesii* by Berkeley.

BRUNNEO-MACULATUS.—Abundant specimens are at Kew, named *brunneo-pictus* by Cooke from Malay. They are light brown, with medium pores and subzonate, slightly rugulose pilei, marked with darker brown spots. It is the basis of the record of

SECTION PETALOIDES.

brunneo-pictus from Malay, in Saccardo, but has no resemblance to the original from Brazil.

MARIANUS.—Known only from the original. It is close to modestus, but not the same. The color is not the same and it is more rigid. The two following are close to Marianus.

ASPERULUS.—From New Caledonia. Type at Paris.

BRACHYPUS.—From West Indies. Type at Paris.

Konipi.—The only type at the British Museum from Ceylon comes in this section, but I have not a very clear idea of it.

16. POLYPORUS. THIN, RIGID. STRONGLY ZONATE WITH GRAYISH ZONES.

GALLOPAVONIS.—Pileus rigid, thin, usually orbicular or reniform with short, lateral stem. Surface with narrow, concentric, gray zones. Pores minute, pale yellowish. Very common in the East, in Java, the Philippines, Samoa, and Australia.



Fig 435

Petaloides Gaudichaudii.

GAUDICHAUDII (Fig. 435).—Pileus thin, with short, lateral stipe. Surface with narrow, concentric, gray zones. Pores medium. Close to the preceding but thinner, flexible, and the pores are distinctly larger. Specimen distributed in Zollinger's collection as *Blumei* belongs here.

INCURVUS.—Pileus thin, rigid, incurved in drying. Surface with strongly cinereous or fuliginous zones. Stipe lateral, from 2-8 cm. long. Pores small, dark. Specimens from Java and Malay.

SECTION PETALOIDES.

17. **PILEUS YELLOWISH BROWN, GILVUS. HYPHAE DEEP YELLOW UNDER THE MICROSCOPE.**

MALIENCIS.—Pileus dark tobacco-brown, rigid, foliaceous, the type being crenately lobed. Stipe short, thick, lateral. Pores short, dark, small. Setae none. There are liberal collections at Kew from Perak.

ARATOIDES.—Close to the preceding and so referred by Bresadola. The type from New Caledonia is more even, not crenate. However, it is probably the same plant.

DISCIPES.—Close to maliencis, but *longer pores*, not foliaceous, and the type is not so dark in color. Type at Kew is from Ceylon.

GLAZIOVII.—Pileus orbicular, thin, brown, with long, lateral stem, in the same plane. Pores minute, brown. Context is thin, brown. Hyphae yellow (but not deep yellow, hence not truly in this section). Stipe lateral, dull surface. Spores not found, but I believe are white. Only known from specimens at Kew from Brazil.



Fig. 436

Petaloides musashiensis.

MUSASHIENSIS (Fig. 436).—Entire plant brown, gilvus. Pileus orbicular, thin, dry, rigid, with soft, velutinate surface. Stipe laterally-dorsally attached, concolorous. Pores minute, concolorous, with soft, velutinate pore mouths. Colored setae rare. Spores not

SECTION PETALOIDES.

found but surely white. As to its context, color, velutinate surface and colored setae this corresponds to *Fomes pomaceus*, but the presence of a stipe removes it from all species with similar structure. Specimen collected by S. Kawamura, Japan. Referred to Hennings' species on the description only as there was no type in the cover when I visited Berlin.

18. GRAMMOCEPHALUS GROUP. PILEUS MARKED WITH RAISED LINES.

A varying assortment of plants that could be referred to one species, and still have marked differences. Those with large pores run into *Favolus*.

a. PORES SMALL. SETAE NONE.

GRAMMOCEPHALUS (Fig. 437).—The type form from the Philippines is orbicular or reniform, reddish brown with medium small pores. This is quite a frequent plant in several countries and seems to vary, so it is hard to decide what to consider as its varieties.



Fig 437

Petaloides grammocephalus.

PERVERSUS (Fig. 438).—Only a variety of *grammocephalus*, more spathulate and of darker color. Determined, published, and distributed (Copeland, No. 18) as *Polyporus coracinus* which, teste the description, viz.: with colored setae, and teste Bresadola, is quite a different plant. I have not seen *coracinus* except this evidently mis-named specimen. This form of *grammocephalus* is common in Samoa.

CAYENNENSIS (Fig. 440).—The American form from South America. Pores minute. Color pale, form spathulate.

SECTION PETALOIDES.



Fig. 438

Petaloides perversus.



Fig. 439

Petaloides albellus.



Fig. 440

Petaloides Cayennensis.

ALBELLUS (Fig. 439).—Similar to the preceding but has larger pores. White. From India, also from Chas. O'Connor, Mauritius.

MACULATUS.—A spotted form (?), dark, known only from a single specimen, Malay.

PLATOTIS.—Known from a partial specimen (Australia). It seems from the color, surface, and pores to be *thick* grammocephalus, but is much too thick, and apparently does not belong to the section.

b. PORES LARGE, FAVOLOID, RUNNING INTO FAVOLUS.

EMERICI.—Pileus the same as that of the type form of grammocephalus, but the pores are larger. From India.

FUSCO-LINEATUS.—This is an *obese* form of grammocephalus with a short, thick stem, 1-1½ cm. thick. The pores are larger than the type form. The surface is smooth, but lined. It was figured in the Trans. Linnean Soc., 2d series, vol. 1.

FAVOLOIDES.—(As a form of grammocephalus.) Pileus corresponds to the type form of grammocephalus, but the pores are large, favoloid. I think it is a better Favolus. Known from Africa.

DORCADIDEUS.—Color rich cinnamon brown. Surface velvety-tomentose with soft, brown hairs. Marked with branched, vein-like reticulations. Pores large, 2-3 mm. deep. Stipe short, lateral. The surface is covered with simple, colored hairs, which have no relation to those of russiceps. There are no cystidia on the hymenium. This strongly marked species is known from but one specimen at the British Museum, from Australia.

SECTION PETALOIDES.

c. SETAEFERA.

Bearing on the pileus and in one species on the hymenium, very peculiar colored spiny or branched setae. See Fig. 441.

CINNAMOMEO-SQUAMULOSUS (Fig. 441).—Pileus and pores dark cinnamon brown, both densely covered with peculiar, colored, branched setae. Pores small. A most striking species known from collections of Dr. Zenker, Camerun, Africa.



Fig. 441

Petaloides cinnamomeo-squamulosus with two types of peculiar cystidia found on the hymenium. (Drawing by Miss Wakefield.)

RUSSICEPS.—Color of pileus dark cinnamon brown, same color and peculiar setae as the preceding, but in this species the setae are absent from the pores. Pores small, pale. Not a form of *grammocephalus* as given, but closely allied to the preceding. Only known from Ceylon.

19. **POLYPORUS. SPECIES DARK COLORED, ALMOST BLACK AT LEAST WHEN DRY.**

a. SETAEFERA.

MEGALOPORUS (Fig. 442).—Pileus dark, spotted, with short, lateral stipe. Pores large, subfavoloid. Hymenium with numerous very peculiar setae (same nature as those of the preceding section). Known from one specimen in Montagne's herbarium from South America. It is close to *Favolus princeps* in its peculiar setae, but otherwise I think it is different.

SECTION PETALOIDES.



Fig. 442

Petaloides megaloporus with the peculiar cystidia. (Drawing by Miss Wakefield).

b. WITHOUT SETAE.

COCHLEARIFORMIS.—Plant black now but has probably changed in drying. Spathulate with a long stipe. Context thin, of peculiar, thick-walled hyphae cells such as have not been otherwise noted by me. Pores small, dark now. Spores globose, 5 mic., white. Only known from types at Kew, from Malay.

TRISTICULUS.—Color dark, almost black, thin, smooth. Pores small. Stipe short, lateral. Known from but two specimens, one at Paris and the other at Kew.

(Cfr. also *stereinus* and *holotephrus* in Section 22.)

20. POLYPORUS. COLORED CONTEXT AND SPORES.

(All of the preceding are supposed to have white spores.)

I do not know that there are any species with colored context and spores and lateral stems except in the section *Ganodermus*. The cotype specimen at Kew of *Polyporus sideroides* has these characters, but the type specimen at Leiden has a thick, spongy, pleuropodal stipe and would be looked for in section 35 of *Spongiosus*.

SECTION PETALOIDES.

21. ABERRANT SPECIES FORMING "NEW GENERA."

POCULA (Fig. 443).—A little species, the smallest known, and for many years supposed to be a *Sphaeria*. A full account is given in *Myc. Notes*, Pol. Issue p. 44. It is not rare in the United States and occurs also in South America, Australia, and Japan.




Fig. 443

Petaloides Pocula. Natural size; also two specimens (X6), and the face of pores (X6).




Fig. 444

Petaloides pusiolus. Enlarged six diameters. Also drawing by Miss Wakefield showing cystidia much magnified and a section of the pores enlarged.

PUSIOLUS (Fig. 444).—Pileus obconic, pendant, 3-4 mm. in diameter, with a short stipe. Surface brown, appearing velutinate from the enlarged projecting ends of the hyphae tissue. Context isabelline,

SECTION PETALOIDES.

the hyphae pale colored. Spores not found but on the hymenium are peculiar, horn shaped, acute, *hyaline* cystidia. This diminutive species came from Sarawak and there are co-types at Kew. I was much pleased to find in it an analogue of the little *Polyporus pocula* of the United States, heretofore supposed to be unique. It figures in our "literature" as a *Fomes* (sic). It is the antithesis of *Fomes*.



Fig. 445

Petaloides obovatus.



Fig. 446

Petaloides mutabilis.

**22. POLYSTICTUS. PILEUS PALE (IN ONE SPECIES DRY-
ING BLACK), USUALLY SPATHULATE OR FLABEL-
LIFORM, THIN. PORES IN A THIN LAYER,
WHITE OR PALE, MINUTE.**

MUTABILIS (Fig. 446).—Pileus thin, marked with ochraceous or grayish zones. Very common in Brazil and also occurs in southern United States.

OBOVATUS (Fig. 445).—Very close to the preceding and I am not sure that it is distinct. It is not so strongly zoned. Usually tapering to the stem. Seems common in Java and the East. I have

SECTION PETALOIDES.

seen no authentic specimens of *obovatus*, but teste Bresadola, *dilatatus* (bis) of Berkeley is a synonym and there is an abundance of that at Kew.

PETALIFORMIS.—Usually cuneate, tapering to the base. I judge from specimens I saw at Berlin that it has a rooting stem. Usually faintly zonate and marked with striations. Close to *mutabilis* and of the same distribution, but is quite different I think.

STEREINUS.—Pileus thin, attenuate behind and sometimes with a short stem, evidently soft and watery when fresh but drying thin, rigid, incurved, and turning black. It seems very common in tropical America, also from the East.

HOLOTEPHRUS.—Spathulate, attenuate at the base but hardly stipitate. Pores minute. Color almost black with metallic zones, and I think has not changed in drying. Known from the type at Kew from Cuba.

ARMENICOLOR.—Pileus thin, minutely pubescent, brown-zoned, tapering to a short but distinct stipe. Pores small, white. Except as to its distinct stem this plant is more closely allied to the *versicolor* group. It is known from one collection at Kew from Cuba.

ANTILOPUS.—This is quite a frequent plant in the museums, and I found it also in Samoa, that had been misdetermined either as *russogramme* or *rasipes* or *palensis*. It is referred by Bresadola to "*vibecinus* var. *antilopus*, Kalch." and it agrees with the cotype at Kew. No specimens of *vibecinus* are preserved and I think no one knows what it was.

23. POLYSTICTUS-MICROPORUS. LATERAL STEM.

(Cfr. also pleuropodal species page 173.)

The section *Microporus*, which has been held to be a genus, is characterized by its thin, rigid context, reddish brown color and minute pores in a very thin layer. We have published a "Synopsis of the section *Microporus*" with illustrations of the species.

AFFINIS (Fig. 447).—Stipe lateral, smooth, dark bay or black. Pileus smooth. Frequent throughout the East.

LUTEUS.—Same as *affinis* but more obese and thicker. Frequent and runs into *affinis*.

MAKUENSIS.—Same as *luteus* but with distinctly larger pores. Referred in my Synopsis as a synonym for *luteus*, which I think on re-examining the type is an error. Known from one collection at Kew from Africa.

PORPHYRITIS.—Probably the same as *luteus*, but from America, where all of this section are rare.

SECTION PETALOIDES.



Fig. 447

Petaloides affinis.




Fig. 448

Petaloides vernicipes.

CARNEO-NIGER.—Same characters as *luteus* except its black color. Occurs in the East. It is the same plant as *microloma*, an earlier name for it.

FLABELLIFORMIS.—Stipe black, lateral. Pileus with pubescent zones. Same as *luteus* except the pubescent zones of the pileus. It varies in all degrees however as to this character. It is the most abundant species in Africa, Samoa, and the East in general.

PTERYGODES.—Pileus sessile; hence does not belong to this section but placed here from analogy, as it is surely a sessile plant of the same nature as the other and perhaps simply a sessile condition of *xanthopus*.

SECTION PETALOIDES.

24. POLYPORUS (CORRESPONDING TO POLYSTICTUS, BUT THICKER) WITH COLOR AND PORES OF THE SECTION MICROPORUS.

VERNICIPES (Fig. 448).—Pileus smooth, rugulose, faintly zonate, shiny, thick, 2-3 mm., rigid. Pores minute. Specimens from Japan, Philippines, and Africa.

SUBFULVUS.—Plant smooth, rigid, pale ochraceous, smooth, with a short, thick, sublateral stem. Pores concolorous, minute, rigid, 2-3 mm. long. Specimens (Wright 135 & 355) at Kew from Cuba, published as ochrotinctus of Bonin Island. Subfulvus was Berkeley's manuscript name and a good name for it.

SIENNAECOLOR.—Comparable to a thick specimen of Polystictus luteus. Same color and pores, but on the Polyporus order with a thick, short, dorsal stem. Known from one specimen from Ceylon. The Brazilian specimen cited was something else I think.

25. RED SPECIES.

SANGUINEUS.—Perhaps no other one species is as abundant in the museums as this. It is the common red species that grows in every warm country of the world. It is strangely rare in Samoa, however. A short, lateral stem, often disciform at the base, is a feature of most collections, but not always, as museum specimens are sessile and even dimidiate. It is close to Polystictus cinnabarinus, the red species of temperate regions, but typically it is thinner and smoother and I think cinnabarinus is never stalked. There is no other bright red species of the tropical world that is likely to be confused with Polystictus sanguineus. Bleached specimens are sometimes collected that have lost all their red color.

CINNABARINUS.—The red species of the temperate world, very similar to the above but thicker and not so brightly colored. It rarely if ever has a distinct stalk, hence does not belong in this section, but we so place it from its evident close relation to the preceding. While not stipitate, it is attached by a reduced base, rarely dimidiate, hence it is related even in its attachment to the stipitate species. Polystictus cinnabarinus is rather rare in Europe, usually on birch, very common in America and favors especially cherry and beech. While ordinarily easily distinguished from the southern species, sanguineus, specimens occur in intermediate territory that are hard to refer.

(Miniatus of Java from its shape might be sought in this section. The brittle, caseous flesh so closely allies it to sulphureus that we place it in that section.)

SECTION PETALOIDES.

26. POLYSTICTUS. WHITE SPECIES.

CONCHIFER.—Pileus white, with a short, lateral stem. Bearing usually secondary, abortive pilei. For a full account see *Polyporoid Issue* page 41. A common plant, always on elm branches, in the United States.

Note.—A number of species of *Polystictus* such as the form *elongatus* of *pergamenus*, have the pilei usually reduced to an attenuate base and might be sought in this section.

Note.—*Glaberrimus*, South America, Montagne (as *Irpex*) and *nepalensis*, India, Berkeley and Pocos, Japan, Berkeley, each known from a single collection, have short, lateral stems, and would be sought here. *Glaberrimus* is close to *biformis* as to its pores. The types of *nepalensis* are quite poor and it may be the same as *glaberrimus*. Pocos has *hirsute* pileus and medium large pores.

SPECIES UNKNOWN TO ME.

Except *leiodermus* no specimens of any of these have been found by me in the museums of Europe

Agaricon, Java, Zollinger.
albo-luteus, Asia, Rostrup.
atro-albus, Africa, Hennings.
bambusicola, India, Hennings.
Baurii, Africa, Kalchbrenner.
cotyledoneus, South America, Spegazzini.
cuneatiformis, Philippines, Murrill.
decrescens, Java, Zollinger.
dilatatus, Java, Léveillé.
discifer, Java, Patouillard.
evanido-squamulosus, Africa, Hennings.
Gregonii, Africa, Smith.
Gualaensis, South America, Patouillard.
hirto-lineatus, Java, Patouillard.
incompletus, Borneo, Cesati.
labiatus, West Indies, Patouillard.
leiodermus, South America, Montagne. My photograph and notes as to this plant do not agree, hence I can not refer it at present.
manubriatus, Sumatra, Léveillé.
monachus, South America, Spegazzini.
olivascens, Asia, Rostrup.
parvmarginatus, South America, Spegazzini.
Pentzkei, Australia, Kalchbrenner.
prostratus, China, Patouillard.
subhydrophilus, Brazil, Spegazzini.
subpendulus, United States, Atkinson.
substereinus, Cuba, Murrill.
tigrinus, Asia, Rostrup.
udus, Java, Junghuhn.
vitiensis, Pacific Island, Reichardt.

SYNONYMS AND REJECTED SPECIES.

Adami, Ceylon, Cooke. Change of *dilatatus* (bis) which is *obovatus*.
albo-cervinus, Brazil, Berkeley=*modestus*. This is the name generally used by Berkeley, who took *modestus* correctly at first but afterwards changed on the evidence of a specimen in Kunze's *exsiccatae* which is not the same as the specimen in the same *exsiccatae* at Upsala.
anisoporus, Europe, Montagne. Type inadequate, probably a little, undeveloped specimen of *Favolus europaeus*.
annularius, Java, Fries. Merely an unauthorized and unnecessary change of *annulatus* of Junghuhn.
apophysatus, Europe, Rostkovius. Only known from an old picture which is probably something abnormal.
atro-cervinus, Brazil. Error in Saccardo for *albo-cervinus*.
atypus, Java, Léveillé. No type exists as far as I could find at Leiden. In the sense of Bresadola it is *brunneolus*. Determinations of Murrill are largely *rubidus*.
aurora, Borneo, Cesati (as *Trametes*). Not seen by me, but stated by Murrill to be a synonym for *atypus*, in which case it is probably *rubidus*, as many of Murrill's determinations of *atypus* are.
bomfinensis, Brazil, Hennings (as *Fomes*, sic)=*Polystictus mutabilis*.

SECTION PETALOIDES.

- caryophyllaceus, South America, Cooke. Type a little remnant, inadequate but probably or possibly mutabilis.
- celebicus, East Indies, Hennings=*carneo-niger*.
- cervicornis, West Indies, Cooke. Something abnormal, but not a synonym for mutabilis as stated.
- cervino-nitens, South America, Berkeley (Schweinitz mss.)=*modestus* and was at first so referred.
- cinerascens, East Indies, L veill . No type found by me. In the sense of Bresadola it is the same as *incurvus*, of which nice specimens are at Kew.
- confundens, Borneo, Cesati=*gallopavonis*.
- coracinus, Philippines, Murrill. Unknown to me. The specimen distributed to Kew (Copeland No. 18) and cited by the author is a form of *grammocephalus*, which does not have cystidia and is an entirely different species from the "type," teste Bresadola in a letter. I have not seen the type, but it was described as having "branched cystidia" and, teste Bresadola, is close to *cinnamomeo-squamosus* if not the same.
- crenatus, Ceylon, Berkeley. This at best is a form of *flabelliformis* with the pubescence covering a small area at the base of the stipe rather than in zones. Only types are at the British Museum. Specimens of L veill 's naming at Leiden and Paris are both wrong.
- cretatus, United States, Cooke. Change of *Ravenelii* (bis) which being in his scense mutabilis was not worth changing.
- cupuliformis, United States, Berkeley=*pocula*.
- Currani, Philippines, Murrill=*vernicipes*.
- decolor, Brazil, Berkeley. Type inadequate.
- delicatus, United States, Berkeley=*a small specimen of fractipes*.
- dendriticus, Mexico, Fries. No type exists. Fries cites Curtis' number 1481 which I do not find at Kew, but on Curtis' notes Berkeley has endorsed "arcticus, Klotzsch."
- dilatatus (bis), Ceylon, Berkeley=*obovatus*.
- diminutus, Australia, Masee. Type is not preserved, but from the figure and description I have no doubt it was based on *rhipidium*.
- eripus, Borneo, Cesati. Unknown to me. Seems from the description to be *flabelliformis*.
- fibro-radians, South America, Montagne=*mutabilis* or close.
- flabellato-lobatum, Africa, Hennings. Teste Bresadola=*cinerescens* (*brunneolus*). I found no type at Berlin.
- gallinaceus, Brazil, Berkeley=*mutabilis*.
- geminella, Moeller, Brazil (as new genus, "Henningsii")=*petaliformis*.
- hispidellus, United States, Peck=*hirtus* of Europe.
- Holstii, Africa, Hennings, also in my Synopsis,=*incomptus*.
- Hostmanni, South America, Berkeley. Type inadequate. There is a better specimen at the British Museum.
- humilis, United States, Peck=*fractipes*.
- hydrophilus, Cuba, Berkeley. Type inadequate. Compare *stereinus*.
- inconspicuus, Africa, Miquel. Said by the author to be the same as *Hostmanni*.
- intonsus, Tasmania, Berkeley. No type exists.
- involutus, Europe, Britzelmayer. Not worth the trouble to bother with his cartoons.
- Kurzianus, Java, Cooke=*Blumei*.
- lacer, Java, Saccardo or Cooke, change of *lacerus* (why?).
- lacerus, Java, Junghuhn=*obovatus* probably, more thin and tapering it seems to me.
- languidus, Africa, Fries. (Fomes in Saccardo.) No type exists. Stated by Fries to be the same as monochrous, which if true is the same as *modestus* and surely not a Fomes.
- lenzitoides, Brazil, Berkeley. Same plant as *aquosus*, which is a much better name for it.

SECTION PETALOIDES.

- Leprieurii* (bis), South America, Montagne (as *Enslenia*)=*pocula*.
Léveilléi, Java, Cooke. Change of *cinerescens*, then changed back.
Libum, Australia, Berkeley. Type inadequate.
licmophorus, India, Masee=*pale* form of *affinis* or possibly a dark form of *obovatus*.
Liebmanni, Mexico, Fries. Type at Upsala, inadequate, a little piece about the size of an oyster cracker. Teste Bresadola it is the same as *stereinus*. It is black and curved in drying, but I think is rather thick for *stereinus*.
ligoniformis, Europe, Bonorden. Unknown. Alleged to be yellow, small pores, white flesh, reddish when broken.
liturarius, Pacific Island, Berkeley. No type exists.
malacensis, error in Saccardo for *maliensis*.
Meleagris, Pacific Island, Berkeley,=*gallo-pavonis*.
Menziesii, Sumatra, Berkeley=*Didrichsenii*.
microloma, Philippines, Léveillé=*carneo-niger*, and an earlier name for it. I have seen the type since my Synopsis was published, it having been placed in its cover since.
minutissimus, Asia, Rostrup. I have not seen this, but I judge from the description it is *rhipidium*.
Mollerianus, Africa, Saccardo. Teste Bresadola, this is a stipitate form of *Polyporus vinosus*. I have seen no stipitate forms. Bresadola refers *vinosus* to *badius* of Junghuhn, not Berkeley.
monochrous, South America, Montagne,=*modestus*. It was first referred to Feei. This is in the sense of Berkeley. "Monochrous, Mont." is quite a different plant.
murinus, Java, Léveillé. No type found by me at Leiden. Teste Fries and Bresadola it is same as *brunneolus*. Most of Murrill's Philippine determinations are *gallopavonis*.
Muelleri, Australia, Kalchbrenner is *grammocephalus* or a form at best.
nanus, Australia, Masee=*rhipidium*.
nigrescens, Brazil, Cooke=*stereinus* (??)
notopus, Java, Léveillé. (Nothopus in Saccardo.) The type at Paris is a little specimen, too inadequate to form an opinion.
palensis, Philippines, Murrill. I have seen more than one species so named by the author, but most of them I would refer to *antilopus*.
peltatus, Central America, Fries. No type exists.
pendula, an alleged synonym for *pocula* used as a juggle. Not based on any evidence but is contrary to the specimens of the author.
peroxydatus, Australia, Berkeley. No type exists.
petaloides, Europe, Fries. No specimens or figures exist. It was based on one collection sent from Pomerania. Unknown now.
phlebophorus, New Zealand, Berkeley. Same as *nicivolor*, which is a much better name for it.
polygrammus, Cuba, Berkeley=*petaliformis*.
pseudo-cinerascens, New Guinea, Hennings=*gallo-pavonis*.
puellaris, Pacific Island, Kalchbrenner=*atypus*, teste Bresadola on the label at Berlin. For me *atypus* in sense of Bresadola is *brunneolus*.
pusillus, West Indies, Léveillé (or Persoon, mss.)=*rhipidium*.
putidus, Central America, Fries. No type exists.
rasipes, East Indies, Berkeley=*obovatus*.
Ravenelii (bis), United States, Berkeley. No type found by me. In the sense of Cooke it=*mutabilis*. There is a cotype in the British Museum from Ravenel, which is same as *mutabilis*, but has no resemblance to *dealbatus* as erroneously stated.
rigescens, Perak, Cooke=*stereinus*.
rufo-ochraceus, South America, Patouillard=*mutabilis*.
russogramme, East Indies, Berkeley. Type inadequate. It has large pores and seems to be something well marked.

SECTION PETALOIDES.

- squamaeformis, Borneo, Berkeley. No type exists.
stereoides, Cuba, Berkeley. Not published but is a manuscript name for plants published as stereinus. The reference in Saccardo, p. 219, probably refers to a Brazilian collection which is mutabilis.
sterinoides, Brazil, Hennings=*petaliformis*.
Stuckertianus, South America, Spegazzini. Seems from the description to be rhipidium, the large, type form.
subflabellum, Africa, Hennings. The types are in alcohol and I can not form much of an opinion of them.
subpulverulentus, Cuba, Berkeley. A form of rhipidium at the best.
subverniceps, Philippines, Murrill=*pterygodes*.
subvernicosus, Brazil, Hennings=*porphyritis* or close to it. It seems to be a slightly thicker plant.
subzonalis, Australia, Cooke=*gallo-pavonis*, pale form.
tomohomiensis, East Indies, Hennings=*grammocephalus*.
torquescens, Africa, Saccardo=*biokoensis*, teste Bresadola=*zonalis*, teste Patouillard. Unknown to me.
unguicularis, Mexico, Fries. No type exists. Judging from the description it is probably the same as mutabilis. Mr. Murrill informs us that it is "only known from the type locality" and that he did not examine the spores. As he evidently never saw a specimen, as none exists, it would have been much more strange and worthy of record under the circumstances if he had examined the spores.
vernicifluus, Tasmania, Berkeley. Type inadequate.
vibecinus, Africa, Fries. No type exists. From the description it is close to *grammocephalus*.
virginus, United States, Schweinitz=*conchifer*.
virax, India, Berkeley. Types at Upsala seem the same as *Liebmanni* to me on comparison, but I am not so certain that it is the same as *stereinus*.

SECTION MERISMUS.

The section *Merismus* embraces plants that have numerous pilei proceeding from the branching of a common stem or rootstalk. Some of them form very large clusters. We also include here the section *Conglobatus* where the pilei proceed from a common tubercular core. As a matter of truth the section *Conglobatus* is quite different from *Merismus* in its manner of growth, but we include it here in order to reduce the number of sectional names.

27. SPORES GLOBOSE, ECHINULATE.

Plants of this section having echinulate, globose spores form a very natural group.

BERKELEYI.—Pilei imbricate, arising from a short, thick stem or root stalk. Surface pale, dull, slightly tomentose and obscurely zoned. Context ($\frac{1}{2}$ -1 inch thick), white becoming isabelline in old specimens, brittle when dry. Pores large, unequal, white. Spores globose, 8 mic., distinctly echinulate. This is the largest and a quite frequent species in the United States, growing usually at the base of a tree.

MONTANUS.—This is the European analogue of *Polyporus Berkeleyi*, and has the same surface, context, spores, and general character. It is much smaller and simpler and more regular. It is quite a rare plant in Europe and very few specimens are in the museums. It occurs in the Alpine regions of France, extending east.

SECTION MERISMUS.

Dickinsii.—This, which is known from a single pileolus at Kew, from Japan, is thinner but seems to have the same context and spores as Berkeley's and is probably the same plant.

ZELANDICUS.—Pileus thin, lentus. Pores large and lacerate. Spores globose, hyaline, echinulate. Known from one collection at Kew from New Zealand. As to pilei it is very similar to *giganteus*, and as to spores it is similar to Berkeleyi.



Fig. 449

Merismus Talpae.

TALPAE (Fig. 449).—Pilei very large, forming a clump several feet in diameter. Surface dark, dull, minutely velutinate, soft to the touch. Pores in the dried specimen small, cinereous. Spores globose, hyaline, 8 mic., slightly rough. This is the largest known species of fleshy *Polyporus*. A specimen from Dutch Guiana at Leiden measures $7\frac{2}{3}$ feet in circumference. It occurs only as far as known in Brazil and other parts of northern South America.

SECTION MERISMUS.

28. SPORES SMOOTH, HYALINE. PLANTS FLESHY.

UMBELLATUS (Fig. 450).—Stem dividing into many branches, each bearing a small pileus centrally attached. Flesh white. Pores decurrent on the branches of the stem, with angular mouths. Spores 3×10 , hyaline, smooth. The stem is said to arise from an underground, thickened rootstalk or sclerotium. This is a most striking and peculiar species and very rare both in Europe and the United States.



Fig. 450

Merismus umbellatus. (Reduced more than half).

FRONDOSUS.—Pileoli very numerous, imbricate, dimidiate or spatulate, fuliginous gray, with white, decurrent pores. They proceed from a common root stalk. This species, which is quite common both in Europe and the United States, sometimes forms large clusters two feet in diameter. It usually grows at the base of a tree or stump. It can be readily known from the more rare species (*umbellatus*) by the insertion of the pileoli, though in their general habits they are very similar.

WYNNEI.—Pileus merismatoid, imbricate, irregular, semi-incrusting, and in habits of growth resembles somewhat *Thelephora terrestris*. Upper surface yellowish brown, smooth but uneven, rugulose. Context thin or a mere pellicle. Pores medium, elongated, round,

SECTION MERISMUS.

4 mm. long. Spores globose, 3 mic., hyaline, smooth. When fresh this is soft and would hardly be sought in *Polystictus* where it is placed in our text books. It is quite a rare plant, known from England, France, Germany. It is not a true *Merismus*, but is more close to this section than any other in its appearance and habits.



Fig. 451

Merismus cremeo-tomentosus.



Fig. 452

Merismus Ridleyi.

GIGANTEUS.—Pilei thin, tough with a dark brown surface and white pores that turn dark when bruised or in drying. Spores globose, 5-6 mic., hyaline, smooth. This which is large, but hardly large enough to be called “giganteus,” is of frequent occurrence in both Europe and the United States. It is similar to *frondosus* but thinner with larger pilei and is tougher.

SECTION MERISMUS.

ANTHRACOPHILUS.—Pilei arising from a hard, woody root-stalk, flabelliform, tapering at the base. Surface rugulose, dark. Pores white when fresh. Spores 4 x 5 subglobose, hyaline, smooth. This plant is very similar to *giganteus* in some respects but is smaller and the dried specimens are hard and subwoody. It is only known from one or two collections at Kew, from Australia.

CREMEO-TOMENTOSUS (Fig. 451).—Pilei thin, flabelliform, tapering to the base and proceeding from the apex of a woody root-stalk, contracted and curved in drying. Surface soft, velutinate. Pores minute. The entire plant is a pale isabelline color. The several pilei proceeding from the apex of a woody rootstalk might be treated as simple pilei and classed in Section 12. It was described by Hennings as a *Fomes*. It never was a *Fomes*. Known from a single specimen from Ule, Brazil.

MULTIPLEX.—Pilei numerous, small, imbricate, with very much the appearance of being small *Polyporus frondosus* with similar general habits and pores. At the base, however, there are numerous white, mycelial fibrils, and it grew on rotten wood, totally at variance with the method of growth of *frondosus*. Known from a single specimen at Kew from Mueller, Australia, but I believe was not formally published.

LITHOPHYLLOIDES.—Only known from the types from Japan at Paris. They are black now, probably discolored from having been sent in alcohol. The small, imperfect pilei proceed from a thick, rooting system. It is quite different in appearance from all others and was compared by the author to the genus *Lithophyllum*, which seems to be a genus of seaweed.

29. MERISMUS-POLYSTICTUS. THIN PLANTS HAVING THE HABITS OF THE SECTION MERISMUS.

RIDLEYI (Fig. 452).—Pileus thin, flabelliform, tapering to the base and proceeding from a rootstalk. Surface smooth, even, gray and beautifully zoned. Pores white, rather large, shallow, elongated. This is a fine species, having the same texture and color and zones as Section 16, page 134, but is merismatoid in its habits of growth. Known from one collection at Kew from Tasmania.

COLENSOI.—Pileoli very numerous, much branched and crispid, thin with dark surface. Pores probably white when fresh, large, shallow. Known from a single rather poor collection at Kew from New Zealand. In general appearance it resembles *Polyporus frondosus* but is much thinner and has smaller pileoli.

FIMBRIATUS (Fig. 453).—Pileus thin, usually imbricate-multiplex, but often more simple, variously cut and lobed. Color pale, dark when dried. Pores white, shallow, usually imperfectly developed and incomplete, the portions of the undeveloped pore walls resembling

SECTION MERISMUS.




Fig. 463

Merismus fimbriatus with section of the hymenium enlarged.

in a faint degree a *Hydnum*. Spores ovate, 4×5 , hyaline, smooth. This is a common species in Brazil and abundant specimens have been sent to Europe, particularly by Glaziou. Owing to the peculiar hymenium it has been variously classed as *Polyporus*, *Polystictus*, *Hydnum*, *Thelephora*, *Craterellus* and *Beccariella*, with a corresponding number of specific names.

30. *MERISMUS CASEOSUS*.

I am not sure that the plants listed here are all merismatoid. The common species of Europe and America, *Polyporus sulphureus*, when growing at the base of a stump usually has a common stem or tubercle, but on the side of a tree it is often sessile, in several imbricate layers. Sometimes on logs it occurs that it has a single pilei, each with its lateral stem. The feature common to all the following species is the caseous, brittle flesh, light and crumbling when dry.

SULPHUREUS.—Pileus bright reddish yellow and when in its prime furnished with a yellowish juice. When old it loses its bright color and becomes dry, light and crumbling. Pores minute, bright sulphur yellow when in prime condition. This is a common species in both Europe and America and occurs in Ceylon, Mauritius, and probably many countries. On the oaks where it habitually grows in Sweden it forms large, conspicuous masses noticeable from a distance. In the United States a form with a stem is quite common at the base of stumps. I have also collected it growing with simple, flabelliform pilei, each with its own short stem.

Polyporus *sp.* — This specimen is distinct by the type from *Polyporus* *sp.* which has a more or less distinct central core, and is not so much as the *Polyporus* *sp.* which has the same brittle flesh and the same pores, but it has a small, smooth, flat form of *Polyporus* *sp.*



Fig. 41

Polyporus *sp.* and *Polyporus* *sp.*

Polyporus *sp.* — This is from a single small collection from Chile. It has the same in the texture, pores, color (in discolored specimens) and surface and in the shape as the *Polyporus* *sp.* *Polyporus* *sp.* The pores are a little larger than the *Polyporus* *sp.* which, as far as I can note, is the only real difference.

Polyporus *sp.* — This form is distinct by the appearance of being sulphureous with larger pores and more abundant. I think it will prove to be only a form.

31. CONGLOBATUS CARNOSUS.

I believe there is no other fleshy species with central core, but that *Polyporus* *sp.* is perhaps the only one. Such a specimen was described collected by T. G. Smith under the name *Polyporus* *sp.* I judge it is the same plant that was named *Polyporus* *sp.* by Peck.

32. CONGLOBATUS FOMES.

But one species of *Fomes* is known with imbricate pilei proceeding from a central core. (See *Bot. Beech* p. 48)

CONGLOBATUS Fig. 455 — Pilei numerous, densely imbricate *over* *the* *core* *and* *the* *margin*. Context and pores brown *and* *of* *a* *hard* *woody* *texture*. Pores minute with darker mouths. *Spores* *are* *found* *by* *me*, *but* *I* *think* *are* *hyaline*. A unique species *known* *only* *from* *the* *United* *States*. It is not common and found only *so* *far* *as* *known* *on* *oak* *or* *beech*.

SECTION MERISMUS.



Fig. 455
Fomes graveolens.
155

SECTION MERISMUS.

MINIATUS (Fig. 454).—This is represented at Leiden by the type from Java also a colored drawing. It is simple, thin, with a short, lateral stipe, and is brick-red when fresh. It loses its color with age and has the same brittle flesh and other characters, and I take it to be only a small, simple, thin form of *Polyporus sulphureus*.



Fig 454

Merismus (Caseosus) miniatus.

SORDULENTUS.—Named from a single, small collection from Chile. It has the same habits, texture, context color (of discolored specimens) and surface, and in my opinion is only a form of *Polyporus sulphureus*. The pores are distinctly larger than the European form which, as far as I can note, is the only real difference.

RETIPORUS.—This from Australia has the appearance of being *sulphureus* with larger pores and firmer context. I think it will prove to be only a form.

31. CONGLOBATUS CARNOSUS.

I believe there is no true fleshy species with central core, but that *Polyporus sulphureus* rarely takes this form. Such a specimen was distributed collected by Toldt in Tirol under the name *Polyporus imbricatus*. I judge it is the same plant that was named *Polyporus flabellatus* by Bresadola.

32. CONGLOBATUS FOMES.

But one species of *Fomes* is known with imbricate pilei proceeding from a central core. (Cfr. Pol. Issue, p. 43.)

GRAVEOLENS (Fig. 455).—Pilei numerous, densely imbricate like tiles, on a hard, woody, central core. Context and pores brown and of a hard, woody texture. Pores minute with darker mouths. Spores not found by me, but I think are hyaline. A unique species known only from the United States. It is not common and found only so far as known on oak or beech.

SECTION MERISMUS.



Fig. 465

Fomes graveolens.

155

SECTION MERISMUS.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

acanthoides, Europe, Bulliard. In the sense of Fries a flabelliform specimen of *rufescens*. In the sense of Bulliard a poor picture of *giganteus*. Used as a juggle for *giganteus* by Quélet.

alligatus, Europe, Fries. Based on Sowerby, t. 422, but no such species known. The picture probably represents an unusual development of *rufescens*.
amygdalinus, United States, Berkeley. The type is very poor, so poor I doubt that it could be recognized on comparison. Said by Ravenel, the collector, to have a strong odor of vanilla or almond. It does not belong in the section *Merismus*.

anax, United States, Cooke. Berkeley's manuscript name for the plant that had been named *Polyporus Berkeleyi*. The species was one of Cooke's posthumous varieties.

Barrelieri, Europe, Viviani. Plate 28 cited by the author is a good picture of *sulphureus*. Plate 36, cited by Fries, is *frondosus*.

Beatiei, United States, Peck=*Berkeleyi*.

bonariensis, South America, Spegazzini. Unknown.

botryooides, "incog.," Lévillé=*Fomes graveolens*, which being such an obvious fact, was a long time being found out. I dug the type out of a cupboard at Paris and at once recognized it.

caespitosa, Brazil, Cooke (as *Beccariella*)=*fimbriatus*.

candidus, Europe, Roth. Unknown to me. No illustration. No specimens in museums. Bresadola told me that he considers it a good species, but I only know what he told me in conversation. I have never seen the plant.

casearius, Europe, Fries. Generally admitted to be only a discolored form of *sulphureus*.

Ceratoniae, Europe, Risso. Based on Barla's *Icon*. t. 30, f. 1-3, which is surely only *sulphureus*.

Cinnatus, United States, Morgan=*sulphureus*, a bright colored form that grows in great abundance at the base of stumps, at Preston, Ohio.

conglobatus, United States, Berkeley=*graveolens*.

discolor, Mauritius, Klotzsch=*sulphureus*.

eurocephalus, Ceylon, Berkeley. The type is much decayed and full of globose, strongly asperate, hyaline spores. Teste Petch, these are spores of a *Hypomyces*, and in viewing them in that light I think it is correct, though I had no suspicion of it when I examined it and mistook them for the spores of the plant.

flabellatus, Europe, Bresadola. From the description I think it is the same plant as has been distributed from Tirol by Toldt as *Polyporus imbricatus*. In my opinion it is an abnormal development of *Polyporus sulphureus*.

Glaziovii, Brazil, Berkeley. This was included in Cooke's *Praeuresores* twice, No. 166 and No. 394, the first as a *Polyporus*, the second as a "Fomes," and both with the *same citation*. The second as a "Fomes" is a *Polyporus* (cfr. page 135) and the first as a *Polyporus* in section *Merismus* is an illusion or error of some kind.

Glaziovii (bis), Brazil, Hennings=*Talpa*.

helopus, Exotic, Patouillard. This is based on a single specimen preserved in the museums at Paris. It is probably abnormal and surely adventitious and was found in the *Jardin des Plantes*.

imbricatus, Europe, Bulliard. Said by Fries to be rare and local in Sweden and is unknown to any one now. I think it was based on *intybaceus* that grew horizontal, hence the lobes are more flat. The common plant called *intybaceus* in England is surely *frondosus*. Most modern books carry both, but I think no one knows two different plants to correspond.

irregularis, England, Sowerby. The *Icones* 423 was referred to *amorphus* by Fries. The color is not right for *amorphus*. When Berkeley first met *Polyporus Wynnei* he referred it to this picture and sent specimens to Montagne. It has a general resemblance to *Wynnei*, but Sowerby's mention of "shallow pores" does not accord.

SECTION MERISMUS.

- lactifluus*, United States, Peck=*Berkeleyi*.
lobatus, Europe, Hudson. Unknown. Fries cites Schaeffer, t. 316 & 317, which are too crude to even be cited. Although attributed to Hudson, this plant was never known to English mycologists.
Merrittii, Philippines, Murrill=*sordulentus*.
multiceps, South America, Patouillard. Unknown to me.
multifida, Portorico, Klotzsch (as *Thelephora*)=*fimbriatus*, teste Bresadola on a label.
Olea, Europe, Panizzi. Unknown to any one I think.
Oxyporus, Europe, Sauter. Unknown to any one I think.
Pauletii, Europe, Fries. Based on an old crude figure from which nothing whatever can be told.
plumarium, Cuba, Berkeley (as *Hydnum*)=*fimbriatus*. Some of the "plumes" are sterile, proliferous pilei on this particular type specimen. While there are many specimens of the plant in the museums under many names, I think the "plumes" are "only known from the type locality."
ramosissimus. An old name often used as a juggle for *umbellatus*.
ramosus, United States, Schweinitz. Published? Poor specimens, but authentic, are *frondosus*.
Rostafinskii, Europe, Blonski. Unknown to me, but the description seems to be *sulphureus*.
rubricus, India, Berkeley. Based on decayed, discolored specimens of *sulphureus*.
scabriusculus, Australia, Berkeley. No type exists.
sparassioides, South America, Spegazzini (as *Craterellus*)=*fimbriatus*.
speciosus. An ancient relic of Europe, 1755, alleged to be the same and used as a cheap juggle for *Polyporus sulphureus*.
subgiganteus, United States, Berkeley,=*Berkeleyi*, as Berkeley did not seem to know his own namesake.
Sumstinei, United States, Murrill,=*the common Polyporus giganteus* both of Europe and the United States and which has not the slightest difference as it grows in either country. I should think Mr. Sumstine would feel quite proud of the honor.
Todari, Europe, Inzenga=*sulphureus*.
trichrous, United States, Berkeley. No type exists.
Warmingii, Brazil, Berkeley=*fimbriatus*.

SECTION SPONGIOSUS.

The section *Spongiosus* embraces those species with soft, light, spongy flesh. These characters are more strongly evident in the dried specimens.

33. CONTEXT PALE OR WHITE. SPORES WHITE.

RUFESCENS (Fig. 456).—Pileus soft, spongy, hirsute. Pores large, daedaloid, pale flesh color when fresh. Spores are globose, 8 mic., hyaline, smooth. Also usually abundant, conidial spores 4 x 6, hyaline, smooth, oval. Not rare in Europe and quite variable. When well developed with a mesopodal stem as shown in Fig. 456 and Persoon's *Icones Pictae* t. 6, often more pleuropodal stem, Sow. t. 191, or lateral stem or even dimidiate (var. *flabelliforme* of Persoon). In the United States perfect forms occur but very rarely. A distorted form is more frequent, called *Polyporus distortus*. The pores of the European form are large and daedaloid and in France it is often called *Daedalea biennis*. In the United States there is not such a strong daedaloid tendency. The two following should be held as forms.

SECTION SPONGIOSUS.



Fig. 466

Spongiosus rufescens.

HETEROPORUS (Fig. 457).—The flabelliform or dimidiate form of *Polyporus rufescens*. Frequent in Europe. Apparently absent from the United States.

DISTORTUS (Fig. 458).—A frequent plant in the United States which I believe to be only a distorted form of *Polyporus rufescens*. Rarely perfect forms are also found in the United States.

ANTHELMINTICUS.—A plant said to be used in India as an anthelmintic is not represented in the museums of Europe by specimens good enough to tell much about its classification. It was compared to *rufescens* by Berkeley.

HYSTRICULUS (Fig. 459).—Known from a single specimen at Kew from Australia, but strongly marked. Surface with rigid, dense, dark hairs. Flesh white, soft. Pores round, medium, flesh color. Spores probably conical, globose, hyaline.

SECTION SPONGIOSUS.



Fig. 467

Spongiosus heteroporus.



Fig. 468

Spongiosus distortus.



Fig. 469

Spongiosus hystriculus.

34. CONTEXT DEEPLY COLORED. SPORES SUPPOSED TO BE WHITE.

SCHWEINITZII.—Pileus dark brown covered with matted tomentum. Context brown, soft, spongy when fresh, brittle when dry. Stipe usually short and thick, rarely central, usually excentric, sometimes wanting. Pores large at first, meruloid, shallow, becoming longer, irregular and often lacerate when old. Spores white in mass, elliptical 4×6 , hyaline, smooth. A frequent plant in both Europe and the United States, but growing usually in pine woods. Sometimes quite large, one to two feet in diameter.

REPSOLDI.—Described from Brazil as growing on trunks and having a "gigantic" stem. Spores 5×7 , hyaline. Not found by me at Berlin, but the description is close to Schweinitzii except the pores are *minute*.

PACHYPUS.—Known only from unsatisfactory specimens in the Herbarium of Montagne. It came from Cuba and seems to me to be closely related to Schweinitzii.

35. CONTEXT DEEPLY COLORED. SPORES COLORED BUT OFTEN BUT FAINTLY.

a. WITH COLORED SETAE ON THE HYMENIUM.

CIRCINATUS.—Context thick, spongy, deeply colored. Stipe mesopodal in the type form, usually pleuropodal. Setae curved.

SECTION SPONGIOSUS.

Spores pale color, 7×12 (or 3×5 in the American plant). The type form which is mesopodal is only known in Europe from Fries' *Icones* t. 3. The pleuropodal form occurs but is also rare. In the United States the mesopodal form is not rare in New England and the pleuropodal form is still more common.

TOMENTOSUS.—Same as the preceding plant but thin, the upper, spongy context layer being very slightly developed. Same color and setae. Frequent in the pine woods of Sweden. If it occurs in central Europe it is rare, and it is unknown from the United States.

The following two are not stipitate, but we mention them here on account of their evident close relationship to the preceding.

TRIQUETER.—In the sense of Fries (?) and Romell, a thick, sessile form of *circinatus*. Same context, color, and setae. It is rare in Europe. In the original sense of Persoon it is in my opinion the same as *cuticularis*.

LEPORINUS.—A thin, dimidiate form of the same plant. Rare in Europe and the United States.

b. SETAE NONE.

SIDEROIDES.—Context ferruginous, spongy. Spores abundant, colored, globose, 8-9 mic. Stipe in the type form thick, pleuropodal, spongy. This species is represented at Leiden by several collections from Java, but not in other museums (except one cotype at Kew). The most perfect forms have a general resemblance in color and shape to *Polyporus Schweinitzii*. Thin forms occur with lateral stipes, and the type of *Polyporus Korthalsii*, at Leiden, appears to be a sessile form.

PUIGGARIANUS.—Context spongy, soft, brown. Spores abundant, *conidial*, globose, minutely rough. According to the collector's notes a large, infundibuliform species. Known from a piece of the pileus at Berlin, from Brazil.

ALBERTINII (Fig. 460).—Pileus mesopodal, with thick, obese stem. Surface and context, pores and stem concolorous, ferruginous. Context soft, spongy with large, inflated hyphae. Pores large, angular, decurrent on the stem. Setae none. Spores abundant, colored, 6×8 , smooth. Specimen at Kew from Endeavor River, Australia, referred by Cooke to *Schweinitzii*, which it closely resembles in general appearance. Named for Albertini, who was the tutor of Schweinitz.

MONTAGNEI.—Pileus obese, ferruginous, with uneven surface. Pores concolorous, medium large, decurrent. Spores 8×10 , pale colored, smooth. Setae none. Rare in both the United States and Europe. This plant is usually placed in the next section, but is rather obese for the allied plants of that section.

FRAGILISSIMUS.—Context soft, spongy, cinnamon-ferruginous. Pores concolorous. Spores $3-4 \times 4-5$, deeply colored, smooth. Stipe mesopodal. Known only from pieces of the pileus in the herbarium of Montagne from French Guiana.

SECTION SPONGIOSUS.



Fig. 460

Spongiosus Albertinii.

SUB-BULBIPES.—Imperfectly known from a half specimen at Berlin from Brazil. It has a soft, spongy context and I judge from my photograph a thin crust. Spores I did not find, but Hennings records them as globose, $3\frac{1}{2}$ -4, light yellow. The plant may belong in Section 5 of *Amaurodermus*.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

abortivus, United States, Peck=*distortus*.
acanthoides, Europe. In the sense of Fries=*rufescens*. In the sense of Bulliard=*poor picture of giganteus*.
benquetensis, Philippines, Murrill=*circinatus*. (?)
biennis, Europe, Bulliard=*rufescens*. Often called *Daedalea biennis*.
conglobatus, Europe, Karsten. Unknown to me, but the description reads much like *distortus*.
dualis, United States, Peck. (Also Polyp. Issue, p. 4)=*leporinus* of Europe.
hispidoides, United States, Peck=*Schweinitzii*.
holophaeus, Europe, Montagne=*Schweinitzii*.
Kalchbrenneri, Europe, Fries. Based on small specimen of *tomentosus* now labeled *perennis* through error of R. Fries.
Korthalsii, Java, L veill . The type specimen appears to be a sessile form of *Polyporus sideroides*, same spores, context, texture, and color. In other namings of L veill  (viz., Zoll. 872) it is *Fomes Harkarlii*, which has no relations whatever to the type at Leiden.
maximus, Europe, Brotero. No one knows and no one doubts but that it equals *Schweinitzii*.
Memmingeri, United States, Murrill. Unknown to me. Seems close to *Montagnei*.

SECTION SPONGIOSUS.

obesus, United States, Ellis (also Polyp. Issue, p. 11)=Montagnei.
occultus, Europe, Lasch. I judge from the little co-type frustule I have seen (Rabh. Exsic. 617) that it is rufescens.
platyporus, India, Berkeley. Type very scanty and inadequate but=I think rufescens, form heteroporus.
proteiporus, Australia, Cooke=rufescens.
Sahranpurennis, India, Hennings. Not found by me at Berlin, but from description seems to be Schweinitzii.
scutiger, Europe, Kalchbrenner. Changed by Fries to Kalchbrenneri and based on small specimens of tomentosus.
sericellus, Europe, Saccardo=rufescens, form heteroporus.
Sistotrema, an old synonym for Schweinitzii, often used as a juggle.
spectabilis, United States, Fries=Schweinitzii.
spongia, Europe, Fries=Schweinitzii. The only type is at Kew.
tabulaeformis, United States, Berkeley=Schweinitzii.
tubulaeformis, United States, Saccardo, misprint for tabulaeformis.

SECTION PELLOPORUS.

Context dry, ferruginous, or yellowish brown with deeply colored hyphae. Setae rare. Pores concolorous. Spores colored, pale in most species. Plants growing in the ground and usually concolorous. Rarely epixyloous. This section is practically the same as the section Perennes of Fries, but we do not use the name as it is misleading for the plants are not perennial.

36. PELLOPORUS POLYPORUS. CONTEXT FLESHY, TOUGH, RATHER BRITTLE, MOSTLY MORE OBESE THAN THE NEXT SECTION.

INDICUS (Fig. 461).—Pileus rugulose, dark brown, zonate. Flesh 5 mm. thick, pale rhubarb color. Hyphae deep yellow. Stipe mesopodal, subligneous, irregular. Pores small, round, 5-8 mm. long, darker than the context with pale mouths. Spores abundant, globose, 5 mic., smooth, deeply colored, mostly guttulate. Known from specimen sent me by B. S. Cavanagh, Baroda, India.

CUMINGII.—Context thin. Pores minute, 2 mm. long, concolorous. Stipe slightly spongy. Spores 3-3½, pale colored. Plant mesopodal. Only known from types at Kew from Philippines (also Mexico?).

VALLATUS (Berkeley).—Context thick, subligneous, bright color. Surface dull, concolorous. Pores small. Known from two specimens at Kew from India. One seems pleuropodal, the other mesopodal.

The two following are much smaller than those that precede, but the rigid, brittle flesh more closely allies them than to the next section.

LUTEO-NITIDUS (Fig. 462).—Pileus irregular, mesopodal or pleuropodal. Stipe slightly spongy, often long rooting. Context thin. Pores minute. Very similar but a larger species than the next. Seems frequent in tropical South America.

SECTION PELLOPORUS.



Fig. 461

Pelloporus indicus. (Top of pileus). Reduced



Fig. 462

Pelloporus luteo-nitidus.



Fig. 463

Pelloporus multiformis.

MULTIFORMIS (Fig. 463).—Stipe lateral, rooting. Pileus flabelliform, thin. Surface striate. Spores 3×4 , colored. Very similar in color and habits to preceding and in the same region. It is smaller and not disposed to take mesopodal forms. Known from three collections all at Paris. It may be only small forms of the previous.

SECTION PELLOPORUS.

37. PELLOPORUS POLYSTICTUS. CONTEXT THIN, FLEXIBLE.

Small plants growing in the ground with mesopodal stems. This is the old section *Perennes* of Fries.

a. COLOR DULL CINNAMON.

PERENNIS.—The most frequent species of the section in both Europe and the United States. Known from its dull, zonate, cinnamon color. Pores small. Spores 4-5 x 8-10, pale colored.

FOCICOLA.—Very similar to the preceding species but with larger pores (1 mm. or more). Frequent in the *southern* United States and there replaces the perennis of the northern states. Unknown from Europe.

DECURRENS.—A rare plant if not a form, based on one collection, from Massachusetts (Cfr. Pol. Issue, p. 12).

PICTUS.—It is a little, slender species known only from specimen in the Herbarium of Fries. It has very thin context, $\frac{1}{2}$ mm., pores 2 mm. long, and the color now is black. Distinct but rare. Spores 6 x 8. The reference Bulliard No. 254 is an error as also are the French records of this plant.

b. COLOR BRIGHT, FERRUGINOUS CINNAMON.

CINNAMOMEUS.—A uniformly bright colored plant with silky, shining, appressed, radiating fibrils. Very rare in Europe, more frequent in the United States. Pores small. Spores 5-6 x 7-10, pale colored under the microscope. The forms from Ceylon and India are otherwise the same but have more globose spores, 6-7 x 8.

OBLECTANS.—The Australian plant (and it is evidently very common in Australia) differs from the European in having usually larger pores and more erect fibrils on the pileus. Spores are 5 x 8, pale. The color is the same and the plants have been held to be the same, but I feel that the Australian plant is entitled to a name.

OBLECTABILIS (Fig. 464).—This, which is based on specimens collected in Brazil, distributed by Ule (No. 48) has been referred to oblectans of Australia. It is similar in color but is more slender, has larger, more shallow pores. The margin is thin and fimbriate and the spores more narrow and more pointed. The spores are 4 x 10, pale colored and tapering at one end.

OBLIVIONIS (Fig. 465).—Entire plant unicolor, of a bright cinnamon color. Pileus soft, subzonate, appressed, fibrillose. Context thin, less than 1 mm. Pores minute, 2 mm. deep. Stipe slender, 3-4 mm. thick, soft tomentose, 7 to 10 cm. long. Spores abundant, elliptical, 7-8 x 10-12 mic., colored. This is a beautiful species known from one collection at Kew from Brazil. It is much larger than any other brightly colored species of this section.

SECTION PELLOPORUS.



Fig. 464

Peltoporus oblectabilis.



Fig. 465

Peltoporus oblivionis.

CUTICULARIS.—Pileus very thin with large, long pores, and was published in Pol. Issue page 11, and has since been received from the same locality. It is a very rare species of the New England States. It is badly named as it is liable to be confused with *Polyporus cuticularis*, to which it has no resemblance.

DEPENDENS.—A most curious little species which hangs pendant from the under side of logs somewhat in the manner of a wasp's nest. Entire plant bright cinnamon color. Spores colored, 5×7 . It is very rare in the United States.

See *Polyporus hamatus*, Addenda, p. 195.

(*Polyporus Montagnei*, see Section 35, is a thicker, more obese plant than others of this section, but its relations are undoubtedly closer here than where we have placed it.)

SECTION PELLOPORUS.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

- bulbipes, Australia, Fries. No type exists=oblectans, teste Fries, but he claimed to have named it in manuscript first.
- carbonarius, Europe, Fries. Based on an old picture (Micheli) and unknown. Said to have white pores, hence can not belong in this section.
- Cladonia, Australia, Berkeley. Types very young=oblectans, young, I think.
- connatus, United States, Schweinitz=perennis.
- connatus, United States, most writers.=focicola.
- Ehrenreichii, Brazil, Hennings. Type inadequate.
- Euphorbiae, China, Patouillard. Unknown to me.
- fimbriatus, Europe, Bulliard=perennis.
- parvulus, British Columbia, Klotzsch=cinnamomeus.
- parvulus, United States, most authors=focicola.
- perdurans, Tasmania, Kalchbrenner. Nothing authentic has been seen by me, but the determinations at Berlin are oblectans.
- peronatus, Europe, Schulzer. Only known from a drawing, showing a volva (sic) surely inaccurate.
- proliferus, United States, Lloyd. Something abnormal.
- Salpincta, New Zealand, Cooke. Types inadequate. Probably an abnormal oblectans. The illustration in the Handbook is largely made up.
- saxatilis, Europe, Britzelmayr. It is purely a waste of time to bother with his work.
- scutellatus, Siberia, Borszczow. Seems from the description to be focicola which, however, is not known excepting in America.
- scutigera, Europe, Kalchbrenner. Changed by Fries to Kalchbrenneri. It was based on a small specimen of tomentosus.
- simillimus, United States, Peck. At best a form of perennis, but it is not possible to maintain it even as a form.
- spatulatus, South America, Hooker. Type in two little fragments. It is probably the same as multiformis.
- splendens, United States, Peck=cinnamomeus.
- subsericeus, United States, Peck=cinnamomeus.
- Verae-crucis, Mexico, Cooke. Known from but one collection from Mexico which I think on comparison is the same as Polyporus Cumingii, known only from the Philippines.

SECTION OVINUS.

In the section Ovinus we give very scanty accounts, as the section has been recently considered and illustrated in full by us in a separate pamphlet. Ovinus embraces the thick, fleshy species of Polyporus with mesopodal or pleuropodal or rarely lateral stems.

38. WITH SCLEROTIUM.

TUBERASTER.—Sclerotium (false) of earth, agglutinate with mycelium. Pores small, white. In Italy, Switzerland.

GOETZII.—Sclerotium small, 2-3 inches. Plant mesopodal. Known from one specimen at Berlin from Africa.

SAPUREMA.—Sclerotium large, bearing several plants. Specimen in alcohol at Berlin from Brazil.

SECTION OVINUS.

MYLITTAE.—Sclerotium, the well known “native bread” of Australia. In the museums I have visited there are no specimens with the fruit, but a photograph of a sclerotium with the Polyporus is at the British Museum.

39. STIPE USUALLY MESOPODAL. PORES SMALL.

OVINUS.—White. Pores small. Spores $3\frac{1}{2}$ -4. Common in Sweden and is also found in Alpine regions of central Europe. Its record in the United States is not certain.

LEUCOMELAS.—Pileus fuliginous. Pores pale. Spores tubercular. Rare in Europe. Not known from the United States.

GRISEUS.—Pileus and pores smoky gray. Spores tubercular. Frequent in the United States. Rare in Europe.

CAERULIPORUS.—Pileus and pores bright blue when in prime. Brown when dry. Very rare in the United States.

POLITUS.—Color dark reddish. Very rare in Europe. Only known from Fries' Icones and a specimen at Kew. Possibly it is a small mesopodial form of confluens.

Peckianus, cfr. Lentus, p. 171.

40. STIPE CENTRAL. PORES LARGE.

(Compare tuberaster in 38.)

41. STIPE USUALLY EXCENTRIC OR IRREGULAR. PORES SMALL.

CRISTATUS.—Color greenish yellow. Frequent in the United States. Rare in Europe.

CONFLUENS.—Color pale reddish, becoming deeper red in drying. Often confluent and irregular. Frequent in Europe and eastern United States.

DISCOIDEUS.—White, becoming isabelline in drying. Grows on logs. Brazil and the type from Cuba.

POPANOIDES.—White or yellowish when dry. No distinct cuticle. With short, thick stipe near one side. Known only from one collection from Mauritius at Kew.

42. STIPE EXCENTRIC. PORES LARGE.

PES CAPRAE.—Surface with small, fasciculate scales. Alpine regions and southern Europe. Very rare in the United States.

SECTION OVINUS.

ELLISII.—Pale yellow with large floccose scales. Very rare in the southern United States.

SQUAMATUS.—With large scales. Dark reddish when dry. Known from one collection from Hungary at Berlin.

(Cfr. Boucheanus in next section.)

43. MELANOPUS. PORES LARGE.

SQUAMOSUS.—Pileus scaly. Pores favoloid. Stipes usually excentric. Common in Europe, rare in the United States.

ROSTKOVII.—Smooth form of squamosus. Rare in Europe and the United States.

BOUCHEANUS.—Small, smoothish form of squamosus with uncolored stipes. Rare in Europe.

LENTINOIDES.—Tropical, smooth form of squamosus.

TUMULOSUS.—Pileus with a smooth, thin cuticle, recalling betulinus. Stipe short, central. Supposed to form large, mycelial masses. Known from the type at Kew from Australia.

TASMANICUS.—Pileus turbinate. Stipe short. Known from one collection from Tasmania at Kew.

44. MELANOPUS. PORES SMALL.

RADICATUS (Fig. 465 bis.).—Mesopodal with a long, rooting base. Not rare in the United States.

HARTMANNI.—Brown, velutinate, with short, thick, excentric stipe. Two collections from Australia at Kew.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

alpinus, Europe, Sauter. From the description seems to be Rostkowii.
asprellus, Europe, Léveillé. Based on a crude figure of *Pes caprae*.
bulbipes, Europe, Beck. Known only from an illustration and is probably the same as Boucheanus. Spores seem a little different, but that is all.
cadaverinus, Europe, Schulzer. Some abnormality.
Campbelli, India, Berkeley. Type is inadequate.
caudicinus. A cheap juggle of *Polyporus squamosus*. It originated in Europe, but has been copied in the United States.
Clusianus, Europe, Britzelmayer. Unknowable.
decurrens, United States, Underwood. Unknown.
Earlei, United States, Underwood=*griseus*.
flavo-squamosus, United States, Underwood=*Ellisii*.
flavo-virens, United States, Berkeley=*cristatus*.
Forquignoni, Europe, Quélet=*Boucheanus*.
fuliginus, Europe, Fries. Unknown, based on an old figure.
holocyaneus, United States, Atkinson=*caeruliporus*.
Kansensis, United States, Ellis. Not seen by me, but is probably *Polyporus melanopus*.



Fig. 465 bls.

Ovinus radicans (reduced about one-fourth). Pore details natural size.

SECTION OVINUS.

- laeticolor, United States, Murrill. Preoccupied. Changed to luteo-luteus by McGinty.
- luteo-luteus, United States, McGinty. Unknown to me.
- Michellii, Europe, Fries. Unknown, based on an old figure.
- Morganii, United States, Peck=*radicatus*.
- myclodes, Australia, Kalchbrenner. Unknown.
- nodipes, India, Berkeley. No type exists.
- novo-guineensis, New Guinea, Hennings. Nondescript.
- olivaceo-fuscus, Ceylon, Berkeley. Type consists of two sections from which nothing can be told. It is probably a young *Boletus*.
- pallidus, Europe, Schulzer=*squamosus* with small scales.
- poripes, United States, Fries. Unknown. No type exists.
- pseudoboletus, South America, Spegazzini. Unknown.
- punctiporus, Europe, Britzelmayr. All of his species are unrecognizable.
- retipes, United States, Underwood=*Pes caprae*.
- Schweinfurthianus, Africa, Hennings. Not a *Polyporus* but a *Boletus*.
- scobinaceus, Europe. Used as a juggle for *tuberaster*.
- subradicatus, United States, Murrill. Probably=*radicatus*, which the author does not seem to know very well.
- subsquamosus, Europe, Linnaeus. Unknown. Probably=*griseus*.
- tessulatus, Europe, Fries. Unknown. Based on an old picture.
- violaceo-maculatus, China, Patouillard. Unknown to me.
- virellus, Europe, Fries. Based on picture=*cristatus* sans doubt.
- viscosus, Europe, Persoon. Not a *Polyporus* but a *Boletus*.
- Whiteae, United States, Murrill. Unknown to me.
- xoilopus, Europe, Rostkovius. Unknown except from a doubtful picture.

SECTION LENTUS.

This section generally has mesopodial stipes. They are thinner, more pliant, or coriaceous than the section *Ovinus*. Also they are mostly epixylous in habitat. All as far as known have pale context hyphae and white spores.

45. LENTUS. PORES SMALL.

a. WHITE.

TRICHOLOMA (Fig. 466).—White, strongly marked with ciliate hairs on the margin when young, but they are detersive. Pileus depressed in the center. Spores 4×6 , hyaline, smooth. Frequent in tropical America.

CRYPTOPUS (Fig. 467).—Growing attached to grass stems in western United States. Similar to *rhizophilus* of Tunis, but has smaller pores and spores.

CORYLINUS.—Only known from illustration (Viv. t. 1) from Italy, but seems very distinct.

LEPTOCEPHALUS.—Only known from an old illustration (Jacq. Misc. 1, t. 12) but seems quite characteristic. In short, it is *elegans* *without* a black stipe. (albiceps, see page 180. Specimens with uncolored stems would be sought here.)

b. GRAYISH OR FULGINOUS BROWN.

BRUMALIS.—Stipe and pileus fuliginous. Pores small but elongated, white. Common both in Europe and the United States, on branches, late in the season.

SECTION LENTUS.



Fig. 466

Lentus tricholoma. With section enlarged, showing marginal hairs.



Fig. 467

Lentus cryptopus.

CILIATUS.—Very close to *brumalis* but lighter color and more slender and grows often attached to buried sticks. It is rare in Europe.

LEPIDEUS.—Close to *brumalis* but of a different color, pale yellowish, fuliginous. Very rare in Europe, known only on birch.

SCABRICEPS.—Known from one specimen at Kew from Cuba. Close to *brumalis* but has more scabrous pileus. Still it is not well named.

GUARANITICUS.—Close to *brumalis* as to shape and size. Close to *lepideus* as to color, but the pores are small and *round*.

VERNALIS.—Close to *brumalis* but more slender and yellowish. Rare in Europe and only known to me from the figure by Quélet.

UMBILICATUS.—Close to *brumalis* but more smooth and *rigid*. Known from the type at Kew from India. I have also a specimen from India from Rev. Theissen.

c. COLOR YELLOW OR REDDISH BROWN.

FUSCIDULUS.—Rare, and the only collection known is in Cooke's herbarium and was collected in England. Figured by Bolton (t. 170). When fresh seems to have white flesh and *yellow* pores.

PECKIANUS.—Pileus thin, infundibuliform, with a central stem. Pale yellow when fresh as are the small pores. Grows in the ground, hence might be sought in *Ovinus*, but is too thin for that section.

SECTION LENTUS.



Fig. 468

Lentus virgatus

VIRGATUS (Fig. 468).—Known from one abundant collection from Cuba. The type is at Kew. Pileus of peculiar reddish brown color and appressed fibrils. Context and pores also reddish brown.

SUBVIRGATUS.—A plant of India very close to *virgatus* of the American tropics as to color, shape, and size, but is devoid of the virgate fibrils and the pores are not at all irregular.

IRINUS.—Known to me only from the figure (Bull. Myc. France, 1888, t. XII). Seems to be a reddish plant with a zonate pileus.

REPANDO-LOBATUS.—Specimen at Paris which is very close to *virgatus* and endorsed by Patouillard on the label as same, but which seems to me to be a little different.

Unknown to me Except from Description.

Guarapiensis, South America.
pauperculus, South America.
tucumanensis, South America.
fuegianus, South America.
Braziliensis, South America.
dictyoporus, West Indies.
depressus, South America.

SECTION LENTUS.

d. MICROPORUS. THIN, RIGID WITH MINUTE, WHITE PORES IN A VERY THIN LAYER. COLOR REDDISH BAY OR SIENNA BROWN.

A natural section (or perhaps one species) which runs also to forms with lateral stems (Cfr. page 142). (See also Synopsis of Microporus section published, 1910.)

XANTHOPUS (Fig. 469).—Stipe smooth, yellow. Pileus smooth. A very common species, particularly in Africa, but it occurs also in the East in general. It seems to be absent from the American tropics.

FLORIDEUS.—At best only a dark form of xanthopus with a short stipe.



Fig. 469

Lentus xanthopus.

CONCINNUS.—Stem slender, black. Pileus covered with a uniform, fine, downy, velvety pubescence. This is a rare form in Africa.

INCOMPTUS.—Stem dark or black. Pileus with pubescent zones. This appears to be a common form in Africa and is frequently sent to Europe with xanthopus. However, I believe they are distinct. It was included in my Synopsis under the name *Holstii*.

SECTION LENTUS.

PSEUDO-PERENNIS.—Stipe dark or black. Pileus densely covered with appressed pubescence, faintly zoned, with narrow, glabrous zones. Known from a single collection at Berlin from Africa. Though the plant has no relations, it has a general resemblance to *Polystictus perennis*.

46. PORES LARGE.

Some of these might be placed in *Favolus* to probably better classification.

a. WHITE.

FAVOLOIDES (Fig. 470).—Known only from one specimen from Africa, in alcohol at Berlin. A very marked species with thin, white, umbilicate pileus, long, slender stem, and large, white, favoloid pores.



Fig. 470
Lentus favoloides.



Fig. 471
Lentus partitus.



Fig. 472
Lentus nanus.

NANUS (Fig. 472).—Known from one collection from Algeria at Paris. A little species growing in the sand. Spores are globose, 4-5, hyaline, smooth. Color seems to me to have been white though described as pale yellowish.

RHIZOPHILUS.—Collected by Patouillard. Common in Tunis, attached to grass culms. Very similar to *cryptopus* of the United States but has larger pores and spores.

SECTION LENTUS.



Fig. 473
Lentus Marmellosensis.



Fig 475
Lentus ciliatus (bis).



Fig. 474
Lentus orbicularis.

PARTITUS (Fig. 471).—Pileus thin, mesopodal and usually more or less parted or lobed, reddish brown, rigid. Pores thin, large, shallow with thin walls, white. Stipe mesopodal, lateral from the imperfect formation of the pileus, slender, dull, dark surface. Known from several collections from Spruce, Brazil, at Kew, also one collection at Berlin.

b. COLOR BRIGHT YELLOW.

TILIAE.—Known only from *Icones*, Kalchbrenner, t. 38. All parts of the plant are bright yellow. Very peculiar if correctly depicted, but probably an exaggeration or imagination of some kind, as the author was much given to drawing imaginary pictures. No such plant is now known in Europe.

c. COLOR BROWN OR BROWNISH.

ARCULARIUS.—Pores large, favoloid, white. Pileus brownish, often infundibulform, more or less scaly. A widely distributed plant

SECTION LENTUS.

in most countries of the world. In the United States it is common in the spring. In Europe it is of a southern range only. Frequent in the tropics.

SPECIES WHICH ARE CLOSE IF NOT THE SAME AS ARULARIUS.

squamiger (as Favolus), Australia.
cremoricolor, India.
semulans, Cuba.
maculatus, India.
tunetanus Algeria. Placed in Melanopus, but I think belongs here.
arculariellus, United States. The late summer form of arcularius.
arculariformis, United States.

CILIARIS (as Favolus) (Fig. 475).—This plant occurs in tropical America and is very close to Polyporus tricholoma but *with much larger pores* and close to Polyporus arcularius, but *with smaller pores*. It was distributed by Ule as a variety of Polyporus tricholoma.

ORBICULARIS (Fig. 474).—Known to me from one collection at Berlin made in that vicinity, referred by Hennings to Boucheanus but surely not that. It has a sublateral stipe and large, favoloid pores. It seems to answer the description, but of course that is only a guess.

LENTUS.—Very much the same as arcularius excepting its color which is paler and the plant is more tough. Formerly collected, apparently in abundance, on old stems of gorse (Ulex) in England, but not in recent years. Recorded in error from the United States.

MARMELLOSENSIS (Fig. 473).—Known from but one specimen from Brazil at Berlin, which is most peculiar. Thin, dark, reddish brown with *large*, white, *round* pores. So thin it might be sought in Polystictus or really in Hexagona from its pore shapes. It is the only similar plant known.

(Boucheanus, see Ovinus, page 168, probably better classed here.)

LENTUS SPECIES UNKNOWN TO ME AND NOT FIGURED.

Incendiarius.—Smooth, white, said to be "copious" in Russia, but unknown in the museums of Europe.

Penningtonia. South America.

Velutipes, China. Said to have a viscid pileus and contorted pores.

Fagicola, United States. Known only from the type locality.

Variiporus, West Indies. Said to resemble Tricholoma, but has large pores.

47. LENTUS. SUBGELATINOUS WHEN FRESH (TENDING TOWARD LASCHIA.)

GRACILIS.—Smooth, reddish brown, small pores, with smooth, reddish stem. When fresh it is subgelatinous. Several collections are known, all from tropical America. Original at Kew from the West Indies. The color of the dried specimens is reddish brown, but we suspect that when fresh it is white.

FORM.—We have from Rev. Rick of Brazil a collection which we think is the same species as the preceding but with notably larger pores.

SECTION LENTUS.

48. LENTUS. ABERRANT SPECIES AS TO SHAPE. INFUNDIBULIFORM, GIBBOUS, OR VERY MINUTE.

CRATERELLUS.—Infundibuliform, brown now, described as white or yellow, and it may have been white. Spores abundant, 4×5 , hyaline, smooth. Only known from original collection at Kew from Cuba.

CONFUSUS (Fig. 476).—In shape, size, and color apparently the same as the preceding and so originally determined. Spores $3-4 \times 12-14$. Known from a collection from Louisiana, sent by Ellis to Kew.



Fig. 476
Lentus confusus.



Fig. 477
Lentus Tuba.



Fig. 478
Lentus acicula
(enlarged six times).

TUBA (Fig. 477).—Peculiar, gibbon shape which has been inaccurately described as cup-shaped. Known only from the original collection at Kew from Cuba.

ACICULA (Fig. 478).—If it is a Polyporus, and I can not say it is not, it is unique in its small size, not much larger than a pin head and known from a single specimen from Cuba at Kew.

SECTION LENTUS.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

- agriceus, Ceylon, Berkeley=*arcularius* as Berkeley himself referred.
alveolaris, United States, Bosc. Figure is a crude representation of *arcularius*.
apalus, Brazil, Berkeley=*gracilis*.
Armitii, Australia, Cooke. No type exists. It was figured in Grevillea but I do not know as to its accuracy. Referred afterwards to *stipitarius*, but that surely was an error if the figure at all represents it.
bataviensis, Java, Holtermann. From the figure I judge it is *Laschia caespitosa*.
Binnendykei, Java, Cooke. No specimen exists. Based on an old drawing, probably pale *arcularius*.
cachoeriacensis, Brazil, Hennings=*partitus*.
callochrous, Léveillé. Neither specimen nor locality known.
clypeatus, South America, Patouillard (as *Laschia*)=*gracilis*.
collybioides, Australia, Kalchbrenner. Type inadequate.
Columbiensis, United States, Berkeley. Oregon, not South Carolina as inaccurately compiled by Mr. Murrill. The type is a little discolored frustule that tells nothing and which should never have been named.
conspicabilis, Europe, Britzelmayr. Not worth the trouble of bothering with his crude cartoons.
Cowellii, West Indies, Murrill. From the description I judge it is the same as *gracilis*, which the author does not seem to know although there is a good type at Kew.
crassipes, India, Curry=*xanthopus*.
cupreo-nitens, Australia, Kalchbrenner=*xanthopus*.
Curtisii, United States, Berkeley (as *Favolus*), a late form of *arcularius*, changed to *arculariellus* by Murrill.
cyathiformis, Léveillé, West Indies. No type exists. Probably the same as *Polyporus craterellus*.
dibaphrus, United States, Berkeley=*brumalis*.
esculentus, Europe, Britzelmayr. Cartoon.
favularis, East Indies, Fries. No type exists.
flavidus (bis), United States, Peck. Changed to *Peckianus*.
flexipes, Brazil, Fries. No type exists. Supposed by Fries to be the same as *gracilis*, but I have little doubt it was the same as *Polyporus Tricholoma*.
floccopus, Europe, Rostkovius. Seems from the crude picture to be the same as *lentus*.
fuligineo-albus, Europe, Trog. Unknown to me.
hapalus, Brazil, Saccardo. A variant spelling of *apalus*.
Humphreyi, West Indies, Hennings. No specimen in the cover at Berlin, but said by Murrill to be the same as *Polyporus Tricholoma*. Whether he saw it or merely guessed at it he does not state.
Katui, Marshall Island, Ehrenberg=*xanthopus*. It was finely illustrated under the name *Katui* and on its merits this name should be used.
luridus, United States, Berkeley=*brumalis*.
meizoporus, Cuba, Berkeley (in Saccardo as a variety of *stipitarius*). I find no type.
melanocephalus, Japan, Patouillard. Type small and inadequate. Probably discolored by alcohol.
Mildbachii (Mss. at Berlin), Africa,=*concinus*.
mycenoides, New Caledonia, Patouillard=*Laschia caespitosa*.
obolus, Central America, Ellis. I have not seen this, but the description indicates *gracilis*.
obscura, China, Kalchbrenner. Unknown. Probably *arcularius*.
paraguayensis, South America, Spegazzini. Same as *guarapiensis*, but more pleuropodal.
Perula, Africa, Palisot. Picture probably represents deformed specimen of *xanthopus*.

SECTION LENTUS.

phaeoxanthus, United States, Montagne. Type is a mere frustule (cfr. Myc. Notes, p. 492).

pisiformis, Australia, Kalchbrenner. "Type" is a little incipient sessile, undeveloped pad, about the size and appearance of a wart. Should never have been named at all and most certainly should never have been put in the section Lentus of *stipitate* fungi where Cooke placed it.

planus, Europe, Wallroth. Unknown. As it grew in the ground it may not belong to the section Lentus.

platensis, South America, Spegazzini. Unknown.

Polyporus Polyporus. This gibberish sounds to me more like the college yell of the Carlisle Indians than "Latin," but is alleged to be a Latin name for *Polyporus brumalis*. I doubt if a Roman barmaid would have been guilty of employing such silly language, and yet men who claim to be "scientific" have the assurance to go into print with such names under the pretext that they are employing Latin.

Puiggarii, Brazil, Spegazzini. Unknown.

quadrans, Australia, Berkeley. No type exists. From the description it seems to be *xanthopus*.

rubripes, Europe, Rostkovius. Known only from an old picture which is probably inaccurate.

rubro-maculatus, Europe, Britzelmayr. Cartoon.

saccatus, Rawak, Persoon=*xanthopus*.

similis, Brazil, Berkeley. Type very scanty but probably=*Polyporus Tricholoma*.

squamoso-maculatus, India, Saccardo. Change of *maculatus* of Berkeley because Peck, twenty years later, published another under the same name. Berkeley's name does not seem to have needed changing very badly on this account.

stipitarius, Cuba, Berkeley=*Polyporus Tricholoma*.

substriatus, Europe, Rostkovius. Only known from an old illustration which appears to me to be the same as *lepideus*.

tubarius, Europe, Quélet. Unknown except from his figure, which seems too close to *vernalis*.

umbilicatus, Java, Junghuhn=*arcularius*, teste Fries. Bresadola claims it is different from *arcularius*, but by such slight structural differences that I can not grasp them. The types at Leiden I should refer to *arcularius*.

Vossii, Europe, Kalchbrenner=*brumalis*, teste Bresadola.

Weddelii, Brazil, Montagne. No type exists.

Zenkeri, Africa, Hennings. No specimens found at Berlin. Seems from description to be pale *xanthopus*.

Zollingeri, Java, Saccardo. Unknown and unknowable from such descriptions.

SECTION MELANOPUS.

Plants that have black stems are called *Melanopus*, but we include in this section only those with pilei that would be classed as *Lentus*. The soft, fleshy species are included in *Ovinus*, Sections 43 and 44. Others with black stems will be found in Section *Petaloides*, Section 23, and in *Microporus*, Section 45d.

49. STIPE PLEUROPODAL OR CENTRAL, RARELY LATERAL. PORES MINUTE.

VARIUS.—Pores very minute, white. Pileus dark bay to almost black, smooth. Flesh firm. Type form is common in Europe and it occurs under varying forms in many countries. The following eight could easily be considered as forms.

ALBICEPS.—Pileus white, smooth, with firm, compact, white flesh. Pores very minute, white, decurrent. Stem mesopodal, white, rarely slightly black at the very base. The stem is not always black at the base hence the plant may not be sought for in this section. Its firm flesh, small pores, and other characters bring it very close to *Polyporus varius* even if occasional specimens were not found showing a "black stem." The plant is rare in the United States.

ELEGANS.—This has all the characters of *varius* except the small size which is so constant that on this one character it is generally held to be a good species.

PICIPES.—Name applied to the black form of *varius* with more velvety stems. Frequent in England. In the United States this name is applied to a thin dark form of *varius*.

LEPRODES.—A deformed, irregular, submerismatoid form of *varius*. Rare in Europe and the United States. Given in Fries as a variety of *Polyporus melanopus*.

ADMIRABILIS.—A large, *white*, thick plant with a rudimentary stem. Very different from *varius* in some features but very close in the essentials. Rare, and occurs in the extreme eastern part of the United States.

BLANCHETIANUS.—For me this is a tropical, reduced form of *varius*. Short stem, small size, but in all essential characters it is the same as *varius*.

DICTYOPUS.—For me the tropical form of *picipes-varius*. Color *black* and it is smaller than the form of temperate regions. Usually known as *infernalis* of Berkeley which is the same thing. Widespread in tropical countries, Brazil, Africa, Ceylon.

PAUCHERI (Fig. 482).—A form of the preceding with striate pileus. Common in Australia and at Kew it is referred to *infernalis*.

MELANOPUS.—Pileus villose or rarely scaly. Growing in the ground attached to buried sticks by which habits it can be told from *varius* which usually grows on trunks or decayed spots of living trees. Rather rare in both Europe and the United States.

SECTION MELANOPUS.



Fig. 479

Melanopus Rhizomorphus.



Fig. 480

Melanopus veluticeps.



Fig. 481

Melanopus Lepreurii



Fig. 482

Melanopus Paucheri.

SECTION MELANOPUS.



Fig. 483

Melanopus rubrocastaneus.



Fig. 484

Melanopus hemicapnodes.




Fig. 485

Melanopus Guyanensis (with pores enlarged X6).

RHIZOMORPHUS (Fig. 479).—Produced by long, black, woody rhizomes which I judge are aerial. Known from abundant material in Montagne's herbarium from tropical America.

VERNICOSUS.—Mesopodal, black. Quite peculiar structure, having the dark hyphae prolonged into protruding pointed setae in the pores. Known from a single specimen at Kew from Brazil.

VELUTICEPS (Fig. 480).—Quite different from all others of this section, the pileus having the general appearance and color of *Polystictus perennis*. Stem black, spores doubtless white. Not related to *perennis* however. Known from one collection from Africa at Kew.

HEMICAPNODES (Fig. 484).—Slender, smooth, sometimes infundibuliform at first. The slender, black stipe is sometimes mesopodal and lateral in the same collection. Originally from Ceylon, common in Samoa and seems to occur in the East generally.

SECTION MELANOPUS.

LEPRIEURII (Fig. 481).—Entire plant including the minute pores is dark, fuliginous. Stipe mesopodal or in most of the specimens lateral. Known from very abundant collections sent Montagne from French Guiana.

HYDNICEPS.—Stipe short, rudimentary but black. The “hydroid” processes of the pileus from which the plant was named are much exaggerated. Known from three apparently undeveloped specimens at Kew, from Cuba. I should not be surprised if it develops into the section *Merisma* when it is well known.

RUBRO-CASTANEUS (Fig. 483).—Stem short, black, mesopodal. Pileus infundibuliform, reddish brown, smooth. Pores small, decurrent. Hyphae pale colored. Spores not found. This is the only truly infundibuliform species with black stem. The pileus is the same peculiar reddish color as *Polyporus virgatus*. Specimens from Malacca (Malay) at Kew.

50. STIPE PLEUROPODAL OR CENTRAL. PORES MEDIUM.

GUYANENSIS (Fig. 485).—Slender with a dark, slender stipe. Pores white, medium, favoloid in shape. Seems to be frequent in tropical America.

PODLACHICUS.—Unknown to me, but described as similar to *elegans* but *larger pores*. Rare in Europe, no doubt.

51. STIPE PLEUROPODAL OR CENTRAL. PORES LARGE, FAVOLOID.

(Probably all better classed in the genus *Favolus*.)

PUTTEMANSII (Fig. 486).—Pileus white, with dull, smooth surface. Pores large, favoloid, white. Stipe all black. Known from a half specimen at Berlin from Brazil. I have also a specimen from Rev. Rick.

WRIGHTII (Fig. 487).—Pileus white with striate surface. Pores large, white. Stipe mesopodal, black, abruptly enlarged at the base. Type from Cuba (Wright 201) but not found at Kew nor cited by Berkeley. Known to me from a specimen from Rev. Rick, Brazil, which seems to accord with the description, but of course I can not say that it is correct.

VADOSUS.—Pileus rigid, pale, with smooth or slightly virgate pileus. Stipe mesopodal, black, with a rooting base. Pores large, favoloid, shallow. Based on a specimen at Berlin, collected in Guadelupe by Duss and determined, evidently in error, as “*Favolus dermoporus*, Pers.” This may be the same as *marasmioides*, which is unknown to me.

SECTION MELANOPUS.



Fig. 486
Melanopus Puttemansii.



Fig. 487
Melanopus Wrightii.



Fig. 488
Melanopus palpebralis.

PALPEBRALIS (Fig. 488).—Pileus dark, thin, with a minutely tomentose surface. Margin ciliate with short hairs. Pores dark, favoloid. Stipe mesopodal, slender, black. Based on a specimen at Paris from French Guiana, labeled by Montagne “ciliaris vel affinis.” It is close to ciliaris but differs in its dark color and *black* stipe.

MARASMIOIDES.—Not seen by me, but it seems peculiar in its slender form and habitat (seed of *Melioma*). Perhaps it is the same as *vadosus*.

SECTION MELANOPUS.

52. STIPE LATERAL, BUT THE PILEUS IS NOT SPATHULATE. PORES MINUTE.

LATERATUS.—Pileus reniform, white, firm. Pores minute, now dark. Stipe lateral, black. Named by Persoon from a specimen from Rawak now preserved in good condition at Paris. Referred by Fries, who never saw it, to *Polyporus affinis*, to which it has no affinity, and compiled in error in Saccardo as a synonym. In my opinion it is a *lateral* stemmed form of *elegans* of Europe.



Fig. 489

Melanopus nephridius.



Fig. 490

Polyporus melanopus (bis).

NEPHRIDIIUS (Fig. 489).—Pileus bay brown, with a slightly scabrous surface. Stipe lateral, short, black. Pores white, minute. Known from a few collections from South America. It is a plant intermediate between this section and the section *Microporus*, but is closer to this.

GAYANUS.—Pileus rigid, firm, smooth. Context pale. Stipe black, lateral, rudimentary. Known from one collection made in Chile by Gay, which has another name, *cycliscus*.

(*Dictyopus* and *hemicapnoides* in the preceding sections sometimes have lateral stipes.)

53. STIPE LATERAL BUT PILEUS NOT SPATHULATE. PORES MEDIUM OR LARGE.

PUSILLUS (bis or tris).—A minute, little plant, known from *one single specimen*, less than 1 cm. in diameter, at Upsala, collected in Brazil, eighty years ago. Stipe lateral, short, round, black. Pores favoloid. Referred by Fries to his then new genus, *Favolus*, but is quite distinct from the type of plants that have since become known under that name.

SECTION MELANOPUS.

MELANOPUS (bis) (Fig. 490).—Pileus reniform, smooth. Pores medium favoloid. Stipe lateral, smooth, black. Known from two little specimens in Montagne's herbarium from South America. Classed by Montagne in Favolus but its relations are entirely with this section.

54. (PETALOIDES) STIPE LATERAL. PILEUS SPATHULATE, TAPERING TO THE STIPE.

These plants could be put in a section *Melanopus* of Section *Petaloides* with equal propriety.

GUILFOYLEI (Fig. 491).—Pileus thick, rigid, smooth, pale or yellowish. Pores very minute. Pileus tapering to a short, black stipe. Originally from Australia. Specimens at Kew from Australia, Samoa, Philippines, Malay, and Mexico.



Fig 491

Melanopus Guilfoylei.



Fig. 492

Melanopus Warburgianus.

WARBURGIANUS (Fig. 492).—Pileus thick, rigid, smooth, *dark*. Pores minute, *dark*. Stipe short, black but there is no distinct ring at the base as shown in Hennings' figure. Known from a single specimen at Berlin from the Celebes. Classed by Hennings and found in Saccardo as *Fomes* (sic). Murrill's Philippine references are all wrong.

XEROPHYLLUS (Fig. 493).—Pileus subreniform, but tapering to the short, black stipe. Surface strongly raised, striate. Pores minute. Known from a single specimen at Kew from New Zealand.

SECTION MELANOPUS.



Fig 493

Melanopus xerophyllus.



Fig 494

Melanopus radiato-scruposus.



Fig 495

Melanopus malnominus.

RADIATO-SCRUPOSUS (Fig. 494).—Pileus spathulate, dark brown, strongly rugulose, striate, tapering to a short, lateral black stipe. Pores small. Known only from the type specimens at Berlin from Brazil.

MALNOMINUS (Fig. 495).—Pileus spathulate, reddish brown, smooth, with crenate, lobed, thin margin, tapering to short, smooth, lateral, black stipe. Pores minute, probably white when fresh. Specimens at Paris from Mexico which had been sent to Cooke and named "Teysmanni, Berk." which does not exist, and the name is not appropriate for a Mexican plant.

SYNONYMS, REJECTED AND UNKNOWN SPECIES.

- atratus*, Mexico, Fries. No type exists.
- atripes*, Asia, Rostrup. Unknown.
- atrofuscus*, South America, Léveillé. Type inadequate. Probably only a short stemmed specimen of *lateralis*.
- Beccarianus*, Bornco, Cesati. Unknown.
- Calyculus*, South America, Patouillard. Unknown to me, but from the figure I judge it is *hemicapnodes*.
- cyathoides*, Europe, Swartz. Unknown in the museums and referred to a small form of *melanopus*.
- cycliscus*, South America, Montagne=*Gyanus* and based on the same collection.
- diabolicus* (bis), Brazil, Spegazzini. Unknown.
- dimorphus*, Malay, Cooke=*hemicapnodes* with a lateral stipe.
- fissus*, United States, Berkeley. Type inadequate. Probably *depauperate picipes* and not *fissile*.
- glabratus*, Australia, Kalchbrenner. Unknown.
- infernalis*, Brazil, Berkeley=*dictyopus*, but most of the specimens in the museums bear this name.
- Juranensis*, Brazil, Hennings (as var. of *Leprieurii*)=*Guyanensis*.
- maculosus*, Central America, Murrill. Unknown.

SECTION MELANOPUS.

- minimus*, Europe, Fries. A tiny, little form of *elegans*. No specimens known.
- nephelodes*, South America, Léveillé. No type exists. Said by Fries to be the same as *infernalis*.
- nigripes*, Africa, Masee=*hemicapnodes*.
- nummularius*, Europe, Bulliard=*elegans*.
- pertenuis*, Asia, Kalchbrenner. Unknown.
- pusillus*, Asia, Rostrup. Unknown to me. Said to be zoned, hence probably does not belong in this section.
- rufo-atratus*, Brazil, Berkeley=*rhizomorphus* (sans rhizomes).
- scabellus*, West Indies, Patouillard=*nephridius*.
- seminigrita*, Brazil, Berkeley=*Guyanensis*.
- Strangerii*, Australia, Mueller. Unknown. Seems from the description to be *dictyopus*.
- subelegans*, West Indies, Murrill. Unknown to me. The description suggests *xerophyllus* or *radiato-scruposus*.
- tephromelas*, South America, Montagne=*Leprieurii*.
- Teysmanni*, Mexico, Cooke. (Mss.) Changed to *malnominus*.
- trachypus*, United States, Montagne. Based on an abortive *picipes*.
- tubaeformis*, Europe, Karsten. Evidently close to *varius* and given as a form in Saccardo.
- Underwoodii*, United States, Murrill. I think will prove to be the same as *admirabilis*.
- versiformis*, India, Berkeley. Based on two little specimens, one of which seems to be *melanopus*, the other I think is different.




Fig. 496
Petaloides osseus.

ADDENDA.

NOTES. COMMENTS AND OMISSIONS.

The following points have come up since this pamphlet has been in type. We include here some comments on Romell's article on Brazilian fungi, a critical and valuable paper which we did not have at Kew, when the main portion of this pamphlet was written.

MESOPODIAL AND PLEUROPODIAL.

These terms which are really coined words, meaning central and excentric stemmed, are for the most part spelled Mesopodal and Pleuropodal in the body of the work. The correct spelling is probably Mesopodial and Pleuropodial to correspond with the word podial as spelled in the Century Dictionary.

SECTION GANODERMUS 2.

LUCIDUS.—Several collections received from C. D. Mahaluxmirala shows that this is one of the most puzzling and variable species in the tropics. The tropical form so referred are not as strongly laccate as the European form. The stem is usually shorter, thicker, and often mesopodial. It is usually pleuropodial in temperate regions. It varies in the tropics also greatly as to the color. Numerous intermediate species connect it with Curtisii.

SECTION AMAURODERMUS 5.

AURISCALPIUM.—A fine collection has recently been received from Gustav Peckolt, Brazil. It is evidently quite a common plant in Brazil. The stipe proceeds from a deep rooting rhizome and probably connected with a creeping rhizome, though none of these specimens show it.

Most of the specimens are "auriscalpium" in shape, though some are mesopodial. I expect in time that Auriscalpium, omphalodes, praetervisus, boleticeps, and rufobadius will all prove to be one and the same species.

SECTION AMAURODERMUS 5.

CHAPERI (Page 112, fig. 406).—We have recently received a specimen from Gustav Peckolt, Brazil, which is the second specimen known. The original in the museum at Paris was supposed to come from Cuba.

It has a character that I have noted in but very few species (*Polyporus vernicosus* p. 182, and *Fomes pachyphloeus* Myc. Notes Pol. Issue, p. 34). The ligneous, colored, hyphae fibrils of the pore walls are pointed on the ends and project into the pores simulating the colored setae of many species, which are called cystidia, and which are distinct from the subhymenial tissue.

SECTION AMAURODERMUS 5.

RUDIS.—Specimens received from Australia are much larger than any in the museums of Europe. One specimen had a pileus eight inches in diameter.

SECTION AMAURODERMUS 5.

VARIABILIS (cfr. p. 111).—I think this is only known from the original collections of Spruce (No. 57 and 183 part) and Berkeley confused under this name two quite different species. Romell refers here and figures a plant from Brazil that must be different, first in its dark color (*atrocastaneus*). *Variabilis* is pale-colored. Second, in its much more obese habits (cfr. Romell's figure, Tab. 2, f. 31, with our figure 405 of the type). Third, in the spores described as "granulis," and they are smooth as far as I can note in the type.

SECTION LIGNOSUS 10.

DEALBATUS.—Few plants have been worse confused than dealbatus. It was originally collected by Ravenel and Curtis and named by Berkeley in 1853. These types are all I have seen. (There are cotypes also in Ravenel's collection in British Museum). They are found in a "Fomes" cover at Kew, but should be classed as a Polyporus.

At the same time Berkeley named mutabilis also from Ravenel's collection. It has little resemblance to dealbatus (cfr. figs. 422 and 446), and is a thin, zonate plant, a Polystictus as classed. Ravenel distributed (Fasc. 3, No. 10), Polystictus mutabilis as dealbatus, and Berkeley, when he made his resume in Grevillea, cites this distribution as being correct. Berkeley made so many "new species" he could not remember them himself.

Dealbatus is found in Saccardo, vol. 6, p. 159, as a "Fomes," and also page 218 as a "Polystictus," both with exactly the same description, word for word, and "Polystictus" has as much resemblance to "Fomes" as a piece of paper has to a lump of coal. It is a good example of the value of our "literature."

Murrill, in his half-hour studies in the principal museums, probably never saw the type specimen, for he gives mutabilis as a synonym for dealbatus. He uses the word dealbatus in keeping with the sacred principle of priority, it having been published in a "prior" position (the previous page) in the same article. Of course, that is much more important for the purpose of a juggle than the fact that the plants have little resemblance or relation to each other, and should not be classified in the same section. Then to make the matter more binding he discovers that mutabilis and unguicularius (which no one knows anything about) and a few others form a "new genus" and takes Polyporus dealbatus as his "type species."

SECTION LIGNOSUS 10.

PAULENSIS.—In a letter just received from Bresadola, he writes me that paulensis is a young specimen of angustus. I should never suspect it and the spores, according to my observations, were not the same.

SECTION LIGNOSUS 11 C.

Context colored. Setae present.

(See Musashiensis, page 135, Fig. 436. Also remarks on page 191).

SECTION LIGNOSUS 11B.

SCOPULOSUS.—This is a marked species with its black stem and smooth pale pileus crust. It was named by Berkeley from Australia fifty years ago, and the type is in good condition although "effete." Then Reichardt published a good figure of it under the name Trametes Rhizophorae. It grows in Australia and in the East abundantly, and has been known to European mycologists from abundant collections for years. It was sent to Murrill from the Philippines in quantities, and he referred it to a "new genus" that he had discovered under the specific name anebus, to which species it has no resemblance whatever. On his second visit to Kew he probably noted that the specimens that Cooke had referred to "anebus" were badly named and he "corrected" it, this time discovering it was another "new genus" and referred it to "Warburgianus," to which species it has less resemblance than to "anebus." It is curious how much easier it is to discover a "new genus" than it is to learn an old, common and well-known species, which is abundantly represented in the museums. We have specimens from A. D. Machardo, Perak; S. Hutchings, Bengal; and Bresadola, Philippines.

When in its prime the surface is smooth, but weathered specimens become scrobiculate. Such a specimen was the "type," and was named evidently from this "character." This is one of the misfortunes that plants often suffer from being named by those who have very scanty knowledge of them.

SECTION PETALOIDES 12.

OSSEUS (Fig. 496, page 188).—We have not included *Polyporus osseus* among the stipitate species, although as the pilei taper to short stem-like bases, it might be sought in this section. Its general habits of growth are imbricate, like many sessile species belonging to the section *Apus* of Fries. We know of no other species similar in this respect. *Polyporus osseus* is a rather rare plant in Europe as it is in the United States. The flesh when growing is firm and in drying becomes quite hard, hence it is not badly named. The European plant is white as far as I have seen specimens, but the American plant is gray. I have known it in America for years without a name for it, for I did not associate it with the white European species. However, Bresadola so refers my American specimens, and I believe correctly. (See Figure 496, page 188).

SECTION PETALOIDES 13.

FRACTIPES.—This has a *lateral stem* and is a *white* plant. The "type" specimen of *fractipes* may have had its stem broken, but the name has no application to the plant usually, and it seems a pity to have a plant so misnamed on account of an accidental feature of the type. It grows more common in the South, but has been found by Peck and called *Polyporus humilis*. It also reaches me from Rev. Rick, Brazil.

Polyporus Peckianus is a *yellow* plant with a *central* stem. It is very rare and I have but two collections (D. B. Griffin, Vermont, and A. S. Bertolet, Canada). It is given as a synonym for *fractipes* by Murrill, but differs entirely. He calls the plant "*fractipes* Berk." and draws the description from Peck's specimen. He puts it in the section *Merismus*, and neither *fractipes* nor *Peckianus* belongs there. One belonging to the section *Petaloides* with lateral stem, the other to *Lentus* with a central stem.

SECTION PETALOIDES 15.

MODESTUS AND RUBIDUS. The former is frequent in South America, the latter in the East, and I have been very much puzzled to decide if they are the same or different species. Both, I think, are rose-colored when fresh, but the old specimens I have seen are more brown, having lost the fresh color. Romell records both from Brazil (with different spores), but as he compares the color of *rubidus* with *vinosus* I judge the determination is doubtful. At Leiden I have seen old specimens determined as *rubidus* which are dimidiate and imbricate, but I do not know if correct or not.

Modestus is confused in the *Kunze exsiccatae*, the specimen at Kew not being the same as the type at Upsala. Berkeley at first had it right, but afterwards misled by the *Kunze* misdetermination he referred it usually to *albo-cervinus*, which is a synonym for *modestus*.

SECTION PETALOIDES 17.

MUSASHIENSIS (page 135, fig. 436).—This, I think would have been better placed in a section of *Lignosus* than *Petaloides*. It should go in Section 11c, being the only species in the section with setae. In its structure it is allied to *gilvus*, etc. My reference of Mr. Kawamura's specimen to Henning's species, on the description only is of course doubtful. The specimen was submitted to Bresadola and was unknown to him, and he probably is acquainted with *Musashiensis*.

SECTION PETALOIDES 18A.

PERVERSUS.—We have so indicated a form of *Polyporus grammocephalus*, which was collected by Copeland (No. 18) in the Philippines and recorded and distributed to three museums in Europe as being *coracinus*, as named by Mr. Murrill. This collection was probably so named by Mr. Murrill, but is quite different in its structure in having no cystidia whatever. The original *coracinus* has very peculiar cystidia, I am told by Bresadola, and is the same or close to *cinnamomco-squamosus* as illustrated, Fig. 441. I have not seen the original of *coracinus*, and until I learned of the mistake I took these Copeland specimens in good faith.

SECTION PETALOIDES 18 C.

It develops that there are several polyporoids with very peculiar setae, as illustrated in Figures 441 and 442. The species considered in this pamphlet are *cinnamomea-squamosus* (p. 138) *russiceps* (p. 138), *megaloporus* (p. 138), and *coracinus* (p. 146, unknown to me). In addition, *Favolus princeps* of Cuba has these same peculiar cystidia and perhaps other species of *Favolus*. Bresadola is inclined to refer them all to one species. They seem different to me in macroscopic characters, though the microscopic characters are quite close.

SECTION MERISMUS 28.

DISPANSUS (Fig. 498). Pileus submerismatoid, appearing to be borne irregular from a common base. Surface smooth. Color yellowish. Pores small, colored, reddish brown (about same color as those of *Polyporus rutilans*). Context thin, the pores almost reaching the cuticle. Spores abundant, globose, smooth, $3\frac{1}{2}$ -4 mic., hyaline.




Fig. 498.

Merismus dispansus.

Type from A. Yasuda (No. 7) from Sendai, Japan.

This is quite similar in its habits to *Polyporus Wynnei* of Europe, which is the only plant to my knowledge that it suggests. Both are doubtfully included in the section *Merismus*. The Japanese plant is abundantly different from the European in its colored pores, and spore shape. We have no plant in the United States that approaches either.

SECTION MERISMUS 29.

FIMBRIATUS.—The common plant in the American tropics which has the peculiar hymenial configuration, as shown in Figure 453, has been classed in six different genera. We consider it as a degenerate type of a polyporoid. We refer it to *fimbriatus* following Bresadola, though the type specimen of *fimbriatus* at Upsala has perfect pores. Otherwise it seems to be the same to us.

SECTION PELLOPORUS 36.

ORIENTALIS (Fig. 499).—Entire plant concolorous, ferruginous brown, turbinate, tapering to a short thick stipitate base. Surface smooth, faintly zoned, minutely velutinate, soft to the touch. Context ferrugineous. Hyphae colored. Pores minute. Setae colored, pointed, straight, thickened at the base. Spores 4 x 5, pale colored or sub hyaline.

Type from Jintaro Umemuro (No. 1) Akazaki, Japan.



Fig. 499.

Pelloporus orientalis.

This species is allied to *Polyporus tomentosus*, but differs in its form, setae, and surface, and has no spongy upper layer. The spores we have not found in numbers so that we are very sure about them, and possibly they are hyaline. In that case the plant in its context color, setae and spores is close to *Polyporus musashiensis* (page 135). While the plant is evidently more closely related to *Polyporus tomentosus* than to any in the section *Pelloporus*, we feel it can not be placed with *tomentosus* in the section *Spongiosus*.

SECTION PELLOPORUS 36A.

(Related to Amaurodermus).

TURBOFORMIS (Fig. 500).—Pileus depressed on top, turbinate, tapering to a short, thick stem at the base. Surface rugulose, reddish brown. Context light gilvous yellow, sublignous. Hypheae



Fig. 500

Pelloporus turboformis.

deep yellow. Pores brown, darker than the context, minute, with glancing mouths. Setae none. Spores globose, non-apiculate, deep colored, smooth, 5-6 mic. in diameter.

Type from G. H. Krumbiegel, Baroda, India.

This species is quite different from anything that I have found named in the museums. It resembles a *Ganoderma* in its general effect, but is quite different in its context color and spores. I am at a loss where to place it for it really should form a section of lignose plants close to *Amaurodermus*. However, I do not like to multiply the sections. It has a distant relationship to *Polyporus vallatus*, hence I place it (provisionally) in a related section, but according to our key characters it would be sought in *Amaurodermus*.

SECTION PELLOPORUS 36.

We have a collection from Prof. A. Yasuda (No. 12), Sendai, Japan, which is unknown to us and we believe unnamed. It seems to have an abortive and probably fictitious stipe. The color context and pores is ferrugineus. Setae none. Spores globose 4-5 white (or perhaps very pale color). We would prefer not to name it until we learn more as to its normal stipe characters.

SECTION PELLOPORUS 37C.

Context thin. Hymenium with Setae.

HAMATUS.—Mesopode, subcoriaceous, infundibuliform. Context thin (not spongy). Setae slender curved. Spores colored, elliptical, smooth, 5-6 x 8-9.

I know this only from Romell's excellent description and figure based on material from Brazil. It differs from all others of the section Pelloporus in having setae. In the nature of the setae (curved) it is close to *Polyporus circinatus* (p. 159) in the section *Spongiosus*.

SECTION OVINUS 39.

We have an unnamed specimen from Prof. Petch, Ceylon, that we would refer to this section, although it is thinner and not so fleshy as others in this section. It might be included in *Lentus*, although as it grows in the ground in its habits it is more allied to *Ovinus*. Prof. Petch tells me it is quite rare in Ceylon and grows in circles.

SECTION MELANOPORUS 44.

HARTMANNI.—The only collections are two at Kew and one recently received by me from Miss Margaret Flockton, Australia. The spores are 5 x 12, hyaline, smooth. The species is badly figured in the Handbook, as it is evidently not red as shown, but brown.

SECTION LENTUS 45 (d MICROPORUS).

INCOMPTUS.—This was given in my Synopsis as a synonym for *flabeliformis*, my view being based on a specimen named by Fries, which has a lateral stipe. In the sense of the figure given in *Reliquiae Afzelianae* it has a mesopodial stipe and is the same (and for me the correct name now) as what was called in my pamphlet *Polystictus Holstii*. Fries evidently did not attach any value to the mesopodial and pleuropodial characters in this section.

SECTION LENTUS 46A.

PARTITUS (cfr. p. 175).—Romell records and figures this plant from Brazil. He records the spores as elliptical, "hyaline-luteolae" 7 x 11-12.

SECTION MELANOPORUS 54.

GUILFOYLEI AND WARBURGIANUS.—These two species are not as close as might be inferred from our figures. *Guilfoylei* is a pale colored plant and *Warburgianus* is almost black. Murrill refers *Guilfoylei* as a synonym for *elegans*, which is almost as bad a reference as his determination of *scopulosus*, first as *anebus*, then as *Warburgianus*. The five species *Guilfoylei* and *elegans*, *anebus*, *scopulosus* and *Warburgianus*, which he has muddled have no resemblance and little relation one to the other.

INDEX SYNONYMS, MISTAKES, ETC.

Species which in my opinion, are synonyms, errors, mistakes or blunders, species that rest on some old vague description of which no material exists, also the species that have not been seen by me and of which no specimens have been found in the principal museums, together with the country from which they have been exploited, and the names of those who are responsible for this work.

	PAGE		
abortivus	United States	Peck	161
acanthoides	Europe	Bulliard	156
Adami	Ceylon	Cooke	145
aemulans	Cuba	Berkeley	176
agariceus	Ceylon	Berkeley	178
Agaricon	Java	Zollinger	145
albo-cervinus	Brazil	Berkeley	145
albo-cinctus	Africa	Patouillard	108
albo-luteus	Asia	Rostrup	145
alligatus	Europe	Fries	156
alpinus	Europe	Sauter	168
alveolaris	United States	Bosc	178
amygdalinus	United States	Berkeley	156
anax	United States	Cooke	156
anisoporus	Europe	Montagne	145
annularius	Java	Fries	145
apalus	Brazil	Berkeley	178
apophysatus	Europe	Rostkovius	145
arculariellus	United States	Murrill	176
arculariformis	United States	Murrill	176
argillaceus	Cuba	Murrill	108
Armitii	Australia	Cooke	178
asperulatus	Philippines	Murrill	108
asprellus	Europe	Léveillé	168
atratus	Mexico	Fries	187
atripes	Asia	Rostrup	187
atro-albus	Africa	Hennings	145
atro-cervinus	Brazil	Error Sacc	145
atro-fuscus	Brazil	Léveillé	187
atypus	Java	Léveillé	145
auriscalpioides	Brazil	Hennings	121
Aurora	Borneo	Cesati	145
avellaneus	Central America	Murrill	108
bambusicola	India	Hennings	145
Barrelieri	Europe	Viviani	156
bataanensis	Philippines	Murrill	121
Bataviensis	Java	Holterman	178
Baurii	Africa	Kalchbrenner	145
Beatiei	United States	Peck	156
Beccarianus	Borneo	Cesati	187
benquetensis	Philippines	Murrill	161
biennis	Europe	Bulliard	161
Binnedykei	Java	Cooke	178
Boleticeps	South America	Patouillard	121
boleticus	South America	B. S. Myc	121
bomfinensis	Brazil	Hennings	145
bonariensis	South America	Spegazzini	156
botryoides	"Incog"	Léveillé	156
Brasiliensis	Brazil	Spegazzini	172
bulbipes	Australia	Fries	166
bulbipes (bis)	Europe	Beck	168

INDEX SYNONYMS, MISTAKES, ETC.

			PAGE
Butignoti	Europe	Boudier	129
cachoeiracensis	Brazil	Hennings	178
cadaverinus	Europe	Schulze	168
caespitosus	South America	Cooke	156
calochrous	Incog.	Léveillé	178
Calyculus	South America	Patouillard	187
Campbelli	India	Berkeley	168
canalium	China	Loureiro	129
candidus	Europe	Roth	156
carbonarius	Europe	Fries	166
caryophyllaceus	South America	Cooke	146
casearius	Europe	Fries	156
cassiacolor	Brazil	Berkeley	121
caudicinus	Europe	Juggle	168
celebicus	East Indies	Hennings	146
Ceratoniac	Europe	Risso	156
cervicornis	West Indies	Cooke	146
cervino-nitens	South America	Berkeley	146
Cincinnatus	United States	Morgan	156
cinerascens	East Indies	Léveillé	146
Cladonia	Australia	Berkeley	166
Clemensiac	Philippines	Murrill	121
Clusianus	Europe	Britzelmayr	168
clypeatus	South America	Patouillard	178
cofeatus	West Indies	Berkeley	108
collybioides	Australia	Kalchbrenner	178
Columbiensis	United States	Berkeley	178
confundens	Borneo	Cesati	146
conglobatus	United States	Berkeley	156
conglobatus (bis)	Europe	Karsten	161
connatus	United States	Schwcinitz	166
conspicabilis	Europe	Britzelmayr	178
coracinus	Philippines	Murrill	146
cotyledoneus	South America	Spegazzini	145
Cowelli	West Indies	Murrill	178
crassipes	India	Curry	178
cremoricolor	India	Berkeley	176
crenatus	Ceylon	Berkeley	146
cretatus	United States	Cooke	146
cuneatiformis	Philippines	Murrill	145
cupreo-nitens	Australia	Kalchbrenner	178
cupuliformis	United States	Berkeley	146
Currani	Philippines	Murrill	108
Currani (bis)	Philippines	Murrill	146
Curtisii (bis)	United States	Berkeley (as Favolus)	178
cyathiformis	West Indies	Léveillé	178
cyathoides	Europe	Swartz	187
cycliscus	Chili	Montagne	187
declivis	Pacific Islands	Kalchbrenner	108
decolor	Brazil	Berkeley	146
decrescens	Java	Zollinger	145
decurrans	United States	Underwood	168
delicatus	United States	Berkeley	146
dendriticus	Mexico	Fries	146
depressus	South America	Patouillard	172
diabolicus (bis)	Brazil	Spegazzini	187
dibaphrus	United States	Berkeley	178
dictyoporus	West Indies	Patouillard	172
dilatatus	Java	Léveillé	145
dilatatus (bis)	Ceylon	Berkeley	146

INDEX SYNONYMS, MISTAKES, ETC.

		PAGE
diminutus	Australia	Massee 146
dimorphus	Malay	Cooke 187
discifer	Java	Patouillard 145
discolor	Mauritius	Klotzsch 156
dualis	United States	Peck 161
Earlei	United States	Underwood 168
Ehrenreichii	Brazil	Hennings 166
Elmerianus	Philippines	Murrill 121
criopus	Borneo	Cesati 146
esculentus	Europe	Britzelmayr 178
Euphorbiac.	China	Patouillard 166
eurocephalus	Ceylon	Berkeley 156
evanido-squamulosus	Africa	Hennings 145
fagicola	United States	Murrill 176
favularis	Pacific Islands	Fries 178
fibro-radians	South America	Montagne 146
fimbriatus	Europe	Bulliard 166
fissus	United States	Berkeley 187
flabellato-lobatum	Africa	Hennings 146
flabellatus	Europe	Bresadola 156
flavidus (bis)	United States	Peck 178
flaviporus	West Indies	Murrill 108
flavo-squamosus	United States	Underwood 168
flavo-virens	United States	Berkeley 168
flexipes	Brazil	Fries 178
floccopus	Europe	Rostkovius 178
formosissimus	South America	Spegazzini 109
Forquignoni	France	Quelet 168
fuegianus	South America	Spegazzini 172
fuligineo-albus	Europe	Trog 178
fuligineus	Europe	Fries 168
gallinaceus	Brazil	Berkeley 146
geminella	Brazil	Moeller 146
glaberimus	South America	Montagne (as Irpex) 145
glabratus	Australia	Kalchbrenner 187
Glaziovii	Brazil	Cooke 156
Glaziovii (bis)	Brazil	Hennings 156
Gregonii	Africa	Smith 145
gualaensis	South America	Patouillard 145
Guarapiensis	Brazil	Spegazzini 172
Haenslerianus	New Zealand	Hennings 109
hapalus	Brazil	var. spelling 178
helopus	Adventitious	Patouillard 156
hemibaphus	Brazil	Berkeley 129
hirto-lineatus	Java	Patouillard 145
hispidellus	United States	Peck 146
hispidoides	United States	Peck 161
holocyaneus	United States	Atkinson 168
holophaeus	Europe	Montagne 161
Hostii	Africa	Hennings 146
Hostmanni	South America	Berkeley 146
humilis	United States	Peck 146
Humphreyi	West Indies	Hennings 178
hydrophilus	Cuba	Berkeley 146
imbricatus	Europe	Bulliard 156
incendiarius	Europe	Fries 176
incompletus	Borneo	Cesati 145
inconspicuus	Africa	Miquel 146
incrustatus	Central America	Fries 109
infernalis	Brazil	Berkeley 187

INDEX SYNONYMS, MISTAKES, ETC.

	PAGE
intonsus	146
involutus	146
irregularis	156
Javanicus	109
Juranensis	187
Kalchbrenneri	161
Kansensis	168
Katui	178
Koenigii	134
Korthalsii	161
Kurzianus	146
labiatus	145
lacer	146
lacerus	146
lactifluus	157
laeticolor	170
languidus	146
leiodermus	145
lenzitoides	146
Leprieurii (bis)	147
Leprieurii var. juranensis	187
Léveilléi	147
Libum	147
licmophorus	147
Liebmanni	147
ligoniformis	147
liturarius	147
lobatus	157
luridus	178
luteoluteus	170
maculatus	176
maculosus	187
malacensis	147
manubriatus	145
maximus	161
meizoporus	178
melanocephalus	178
Meleagris	147
Memmingeri	161
Menziesii	147
Merrittii	157
Michelii	170
microloma	147
Mildbachii	178
minimus	188
minutissimus	147
Mollerianus	147
monachus	145
monochrous	147
Morganii	170
Muelleri	147
multiceps	157
multifidus	157
murinus	147
mycenoides	178
myclodes	170
nanus (bis)	147
neglectus	109
nephelodes	188
Nepalensis	145

INDEX SYNONYMS, MISTAKES, ETC.

			PAGE
nigrescens	Brazil	Cooke	147
nigripes	Brazil	Fries	121
nigripes (bis)	Africa	Massee	188
nodipes	India	Berkeley	170
notopus	Java	Léveillé	147
novo-guineensis	New Guinea	Hennings	170
nummularius	Europe	Bulliard	188
nutans	Central America	Fries	109
obesus	United States	Ellis	162
obolus	Central America	Ellis	178
obscura	China	Kalchbrenner	178
obsoletus	Brazil	Fries	129
occultus	Europe	Lasch	162
Olea	Europe	Panizzi	157
olivascens	Asia	Rostrup	145
olivaceo-fuscus	Ceylon	Berkeley	170
oxyporus	Europe	Sauter	157
Pala	South America	Léveillé	121
palensis	Philippines	Murrill	147
pallidus	Brazil	Berkeley	129
pallidus (bis)	Europe	Schulzer	170
paraguayensis	South America	Spegazzini	178
Parmula	Brazil	Berkeley	121
parvimaginatus	South America	Spegazzini	145
parvulus	British Columbia	Klotzsch	166
passerinus	Brazil	Berkeley	121
Pauletii	Europe	Fries	157
pauperculus	South America	Spegazzini	172
peltatus	Costa Rica	Fries	147
pendula	United States	Juggle	147
Penningtonii	South America	Spegazzini	176
Pentzkei	Australia	Kalchbrenner	145
perdurans	Tasmania	Kalchbrenner	166
peronatus	Europe	Schulzer	166
peroxydatus	Australia	Berkeley	147
pertenuis	Asia	Kalchbrenner	188
Perula	Africa	Palisot	178
perzonatus	Cuba	Murrill	109
petaloides (bis)	Europe	Fries	147
Pes-simiae	Brazil	Berkeley	109
phaeoxanthus	United States	Montagne	179
phlebophorus	New Zealand	Berkeley	147
pisiformis	Australia	Kalchbrenner	179
planus	Europe	Wallroth	179
platensis	South America	Spegazzini	179
platyporus	India	Berkeley	162
plumarius	Cuba	Berkeley (as Hydnum)	157
Pocos	Japan	Berkeley	145
polygrammus	Cuba	Berkeley	147
poripes	United States	Fries	170
praelongus	Cuba	Murrill	109
procerus	Brazil	Berkeley	121
proliferus	United States	Lloyd	166
prostratus	China	Patouillard	145
proteiporus	Australia	Cooke	162
pseudoboletus	South America	Spegazzini	170
pseudo-cinascens	New Guinea	Hennings	147
puellaris	Pacific Islands	Kalchbrenner	147
Puiggarii	Brazil	Spegazzini	179
pulcher	Africa	Fries	121

INDEX SYNONYMS, MISTAKES, ETC.

			PAGE
pullatus.	China.	Cooke.	121
pulverulentus.	West Indies.	Murrill.	109
punctiporus.	Europe.	Britzelmayr.	170
pusillus (bis)	Asia.	Rostrup.	188
pusillus.	West Indies.	Léveillé.	147
putidus.	Central America.	Fries.	147
quadrans.	Australia.	Berkeley.	179
ramosissimus.	Europe.	Juggle.	157
ramosus.	United States.	Schweinitz.	157
rasipes.	East Indies.	Berkeley.	147
Ravenelii (bis).	United States.	Berkeley.	147
retipes.	United States.	Underwood.	170
Rhizophorac.	East Indies.	Reichardt.	129
rhodophaeus.	Java.	Léveillé.	129
rigescens.	Perak.	Cooke.	147
Rostafinskii.	Europe.	Blonski.	157
rubricus.	India.	Berkeley.	157
rubripes.	Europe.	Rostkovius.	179
rubro-maculatus.	Europe.	Britzelmayr.	179
rufo-atratus.	Brazil.	Berkeley.	188
rufobadius.	South America.	Patouillard.	121
rufo-ochraceus.	Brazil.	Patouillard.	147
rugosus (bis).	Brazil.	Berkeley (as Porothelium).	121
russogramme.	East Indies.	Berkeley.	147
saccatus.	East Indies.	Persoon.	179
Sahranpurensis.	India.	Hennings.	162
Salpincta.	New Zealand.	Cooke.	166
saxatilis.	Europe.	Britzelmayr.	166
scabellus.	West Indies.	Patouillard.	188
scabriusculus.	Australia.	Berkeley.	157
Schweinfurthianus.	Africa.	Hennings.	170
scleropodius.	Africa.	Léveillé.	129
scobinaceus.	Europe.	Juggle.	170
scutellatus.	Siberia.	Borszczow.	166
scutiger.	Europe.	Kalchbrenner.	162, 166
semiclausus.	Brazil.	Berkeley.	129
seminigrita.	Brazil.	Berkeley.	188
sericellus.	Europe.	Saccardo.	162
similis.	Brazil.	Berkeley.	179
simillimus.	United States.	Peck.	166
Sistotrema.	Europe.	Juggle.	162
sparassioides.	South America.	Spegazzini (as Craterellus).	157
spathulatus.	South America.	Hooker.	166
speciosus.	Europe.	Juggle.	157
spectabilis.	United States.	Fries.	162
splendens.	United States.	Peck.	166
spongia.	Europe.	Fries.	162
squamaeformis.	Borneo.	Berkeley.	148
squamiger.	Australia.	Berkeley (as Favolus).	176
squamoso-maculatus.	India.	Saccardo.	179
stereoides (bis).	Cuba.	Berkeley.	148
sterinoides.	South America.	Hennings.	148
stipitarius.	Cuba.	Berkeley.	179
stipitatus.	Central America.	Murrill.	109
Strangerii.	Australia.	Mueller.	188
Stuckertianus.	South America.	Spegazzini.	148
subamboinensis.	Brazil.	Hennings.	109
subelegans.	West Indies.	Murrill.	188
subflabellus.	Africa.	Hennings.	148
subfornicatus.	Central America.	Murrill.	109

INDEX SYNONYMS MISTAKES ETC.

			PAGE
	United States	Berkeley	157
	Brazil	Spegazzini	145
	West Indies	Murrill	109
	United States	Arnsperg	145
	Cuba	Berkeley	148
	United States	Murrill	170
	Central America	Murrill	121
	Samoa	Berkeley	121
	United States	Peck	166
	Europe	Linnaeus	170
	Cuba	Murrill	145
	Europe	Rostkovius	179
	Philippines	Murrill	148
	Brazil	Hennings	148
	Australia	Cooke	148
	United States	Murrill	157
	United States	Berkeley	162
	South America	Montagne	188
	Europe	Fries	170
	Mexico	Cooke Miss	188
	Asia	Rostrup	145
	Europe	Inzenga	157
	East Indies	Hennings	148
	Africa	Saccardo	148
	United States	Montagne	188
	United States	Berkeley	157
	Europe	Quélet	129
	Europe	Fries	129
	United States	Murrill	109
	Europe	Karsten	188
	Europe	Quélet	179
	United States	Error Sacc.	162
	South America	Spegazzini	172
	Algeria	Patouillard	176
	Java	Junghuhn	145
	Java	Junghuhn	179
	Africa	Fries	121
	United States	Murrill	188
	Mexico	Fries	148
	West Indies	Murrill	176
	China	Patouillard	176
	Mexico	Cooke	166
	Tasmania	Berkeley	148
	India	Berkeley	188
	Africa	Fries	148
	China	Patouillard	170
	India	Berkeley	148
	Europe	Fries	170
	United States	Schweinitz	148
	Europe	Persoon	170
	Pacific Islands	Reichardt	145
	Europe	Kalchbrenner	179
	Brazil	Berkeley	157
	Brazil	Montagne	179
	United States	Murrill	170
	Europe	Rostkovius	170
	Brazil	Berkeley	121
	Africa	Hennings	179
	Java	Saccardo	179

INDEX.

Index of the species considered valid in this pamphlet, the section to which they belong, the country from whence they were exploited, and the personal name that may be added to them by those who favor this system of advertisement. Those species that appeal to me as being good and distinct and as having merit, I put in bold face type. Those that are better considered as varieties or which are based on more or less doubtful material are marked with a star in light face type.

			PAGE
acicula	Lentus	Cuba	Berkeley 177
admirabilis*	Melanopus	United States	Peck 180
affinis	Petaloides (Micro.)	Java	Nees 142
Africanus	Ganodermus	Africa	McOwan 103
albellus* (bis)	Petaloides	India	Massee 137
Albertinii	Spongiosus	Australia	Mueller 160
albiceps	Melanopus	United States	Peck 180
Alluandi	Ganodermus	Africa	Patouillard 107
Amboinensis	Ganodermus	East Indies	Fries 102
angustus	Amaurodermus	South America	Berkeley 114
annulatus	Petaloides	Java	Junghuhn 131
anthelminticus*	Spongiosus	India	Berkeley 158
anthracophilus	Merismus	Australia	Cooke 152
antilopus	Petaloides	Australia	Kalchbrenner 142
aquosus	Petaloides	Brazil	Hennings 130
aratoides*	Petaloides	New Caledonia	Patouillard 135
arcularius	Lentus	Europe	Batsch 175
arenatus	Lignosus	New Guinea	Patouillard 126
armenicolor	Petaloides	Cuba	Berkeley 142
asperulus*	Petaloides	New Caledonia	Patouillard 134
atropurpureus*	Lignosus	Brazil	Berkeley 126
Auriscalpium	Amaurodermus	Brazil	Persoon 113
basilapidoides*	Amaurodermus	Australia	McAlpine 115
Berkeleyi	Merismus	United States	Fries 148
biokoensis	Petaloides	Africa	Hennings 131
Blanchetianus*	Melanopus	Brazil	Montagne 180
Boninensis*	Ganodermus	Bonin Island	Patouillard 104
Boucheanus*	Ovinus	Europe	Klotzsch 168
brachyporus	Petaloides	South America	Montagne 132
brachypus*	Petaloides	West Indies	Léveillé 134
brumalis	Lentus	Europe	Persoon 170
brunneolus	Petaloides	Philippines	Berkeley 133
brunneo-maculatus	Petaloides	India	Currey 133
brunneo-pictus	Lignosus	Brazil	Berkeley 127
caeruliporus	Ovinus	United States	Peck 167
calcigenus	Amaurodermus	Brazil	Berkeley 111
camerarius	Lignosus	Brazil	Berkeley 126
candidus*	Petaloides	Europe	Persoon 132
carneo-niger	Petaloides	Australia	Cooke 143
Cayennensis*	Petaloides	Brazil	Montagne 136
Chaperi	Amaurodermus	Cuba	Patouillard 112
ciliaris*	Lentus	South America	Montagne 176
ciliatus	Lentus	Europe	Fries 171
cinnabarinus	Petaloides	Europe	Jacquin 144
cinnamomeo-squa-			
mulosus	Petaloides	Africa	Hennings 138
cinnamomeus	Pelleporus	Europe	Jacquin 164
circinatus	Spongiosus	Europe	Fries 159

INDEX VALID SPECIES.

				PAGE
cochlear	Ganodermus	Java	Nees	103
cochleariformis	Petaloides	India	Cooke	139
Colensoi	Merismus	New Zealand	Berkeley	152
conchifer	Petaloides	United States	Schweinitz	145
concinnum	Lentus	Africa	Palisot	173
confluens	Ovinus	Europe	Albertini	167
confusus	Lentus	United States	Massec	177
corrugis	Lignosus	Europe	Fries	122
corylinus*	Lentus	Europe	Viviani	170
Craterellus	Lentus	Cuba	Berkeley	177
cremeo-tomentosus	Merismus	Brazil	Hennings	152
cristatus	Ovinus	Europe	Persoon	167
cryptopus	Lentus	United States	Ellis	170
Cumingii	Pelleporus	Philippines	Berkeley	162
Curtisii	Ganodermus	United States	Berkeley	102
cuticularis (bis)	Pelleporus	United States	Morris	165
dealbatus	Lignosus	United States	Berkeley	124
decurrens*	Pelleporus	United States	Morris	164
dependens	Pelleporus	United States	Berkeley	165
diabolicus	Fomes (stipitatus)	Brazil	Berkeley	100
Dickinsii*	Merismus	Japan	Berkeley	149
dictyopus*	Melanopus	South America	Montagne	180
Didrichsenii	Petaloides	Pacific Islands	Fries	133
discipes*	Petaloides	Ceylon	Berkeley	135
discoideus	Ovinus	Cuba	Berkeley	167
dispansus	Merismus	Japan	Yasuda	192
distortus*	Spongiosus	United States	Schweinitz	158
dorcadideus	Petaloides	Australia	Berkeley	137
dubiopansus*	Lignosus	Brazil	Demazio	125
elegans	Melanopus	Europe	Bulliard	180
Ellisii	Ovinus	United States	Cooke	168
Emerici*	Petaloides	India	Cooke	137
Emini	Ganodermus	Africa	Hennings	105
exilis	Amurodermus	Brazil	Berkeley	121
fasciculatus	Amurodermus	Africa	Patouillard	117
favoloides	Lentus	Africa	Hennings	174
favoloides* (bis)	Petaloides	Africa	Hennings	137
fimbriatus	Merismus	Brazil	Fries	152
flabelliformis	Petaloides (Micro.)	Mauritius	Klotzsch	143
flexipes	Ganodermus	China	Patouillard	104
florideus*	Lentus (Micro.)	India	Berkeley	173
focicola	Pelleporus	Cuba	Berkeley	164
fornicatus	Ganodermus	Brazil	Fries	104
fractipes	Petaloides	United States	Berkeley	131
fragilissimus*	Spongiosus	South America	Montagne	160
frondosus	Merismus	Europe	Fl. Dan	150
fuscidulus	Lentus	Europe	Schraeder	171
fusco-lineatus*	Petaloides	Australia	Berkeley	137
fusco-maculatus	Petaloides	Samoa	Bresadola	130
gallo-pavonis	Petaloides	Australia	Berkeley	134
Gaudichaudii	Petaloides	Malay	Léveillé	134
Gayanus	Melanopus	Chili	Léveillé	185
giganteus	Merismus	Europe	Persoon	151
Glaziovii	Petaloides	Brazil	Berkeley	135
glutinifer*	Petaloides	Mauritius	Cooke	130
Goetzei	Ovinus	Africa	Hennings	166
gracilis	Amurodermus	Brazil	Berkeley (as Hexagona)	117
gracilis	Lentus	West Indies	Klotzsch	176
grammocephalus	Petaloides	Philippines	Berkeley	136

INDEX VALID SPECIES.

			PAGE
graveolens	Fomes (Conglobatus).....	United States.....	Schweinitz..... 154
griseus	Ovinus.....	United States.....	Peck..... 167
guaraniticus	Lentus.....	South America.....	Spegazzini..... 171
Guilfoylei	Melanopus.....	Australia.....	Berkeley..... 186
guyanensis	Melanopus.....	South America.....	Montagne..... 183
hamatus	Pelloporus.....	Brazil.....	Romell..... 195
Hartmanni	Ovinus.....	Australia.....	Cooke..... 168
hemicapnodes	Melanopus.....	Ceylon.....	Berkeley..... 182
Henningsii	Ganodermus.....	Africa.....	Stuhlman..... 105
heteromorphus	Amaurodermus.....	Brazil.....	Léveillé..... 120
heteroporus*.....	Spongiosus.....	Europe.....	Fries..... 158
Hildebrandi*.....	Ganodermus.....	Africa.....	Patouillard..... 107
hirtus	Petaloides.....	Europe.....	Quélet..... 130
holotephrus	Petaloides.....	Cuba.....	Berkeley..... 142
hydniceps	Melanopus.....	Cuba.....	Berkeley..... 183
hypoplastus	Lignosus.....	Brazil.....	Berkeley..... 126
hystriculus	Spongiosus.....	Australia.....	Cooke..... 158
incomptus	Lentus (Micro.).....	Africa.....	Fries..... 173
incrustans*.....	Ganodermus.....	United States.....	Langlois..... 102
incurvus	Petaloides.....	Malay.....	Cooke..... 134
indicus	Pelloporus.....	India.....	Massee..... 162
insularis	Amaurodermus.....	New Caledonia.....	Patouillard..... 117
intermedius	Amaurodermus.....	Brazil.....	Bresadola..... 112
irinus*.....	Lentus.....	South America.....	Patouillard..... 172
Janseanus	Petaloides.....	Java.....	Hennings..... 132
Japonicus*.....	Ganodermus.....	Japan.....	Fries..... 102
juriensis	Amaurodermus.....	Brazil.....	Hennings..... 121
lateratus	Melanopus.....	Rawak.....	Persoon..... 185
Lauterbachii*.....	Ganodermus.....	New Guinea.....	Hennings..... 102
lentinoides*.....	Ovinus (Melan.).....	South America.....	Hennings..... 168
lentus	Lentus.....	Europe.....	Berkeley..... 176
lepideus	Lentus.....	Europe.....	Fries..... 171
leporinus*.....	Spongiosus.....	Europe.....	Fries..... 160
Leprieurii	Melanopus.....	South America.....	Montagne..... 183
leprodes*.....	Melanopus.....	Europe.....	Rostkovius..... 180
leptocephalus*.....	Lentus.....	Europe.....	Jacquín..... 170
leptopus	Amaurodermus.....	Rawak.....	Persoon..... 115
leucomelas	Ovinus.....	Europe.....	Persoon..... 167
Lingua	Ganodermus.....	Java.....	Nees..... 104
lithophylloides*.....	Merismus.....	Japan.....	Patouillard..... 152
longipes	Amaurodermus.....	South America.....	Léveillé..... 115
lucidus	Ganodermus.....	Europe.....	Leysser..... 102
luteo-nitidus	Pelloporus.....	Brazil.....	Berkeley..... 162
luteus	Petaloides (Micro.).....	Java.....	Nees..... 142
macer*.....	Amaurodermus.....	Brazil.....	Berkeley..... 119
maculatus*.....	Petaloides.....	Malay.....	Berkeley..... 137
makuensis*.....	Petaloides.....	Africa.....	Cooke..... 142
maliencis	Petaloides.....	Malay.....	Cooke..... 135
malnominus	Melanopus.....	Mexico.....	La Salle..... 187
marasmioides*.....	Amaurodermus.....	Brazil.....	Berkeley..... 121
marasmioides* (bis).....	Melanopus.....	West Indies.....	Patouillard..... 184
Marianus*.....	Petaloides.....	Pacific Islands.....	Persoon..... 134
marmellosensis	Lentus.....	Brazil.....	Hennings..... 176
mastoporus	Ganodermus.....	East Indies.....	Léveillé..... 104
Makuensis	Petaloides (Micro.).....	Africa.....	Cooke..... 142
megaloporus	Petaloides.....	South America.....	Montagne..... 138
melanopus	Melanopus.....	Europe.....	Schumann..... 180
melanopus (bis)	Melanopus.....	South America.....	Montagne..... 186
miniatus*.....	Merismus.....	Java.....	Junghuhn..... 154

INDEX VALID SPECIES.

			Pz.
modestus	Petaloides	South America	Fries 135
Montagnei	Spongiosus	Europe	Fries 160
montanus	Merismus	Europe	Quélet 148
multiformis	Pelloporus	South America	Montagne 163
multipler	Merismus	India	Berkeley 152
musashiensis	Petaloides	Japan	Hennings 135
mutabilis	Petaloides	United States	Berkeley 141
Mylittae	Ovinus	Australia	Cooke 167
nanus	Lentus	Algeria	Montagne 174
nephridius	Melanopus	Brazil	Berkeley 185
nicolor	Petaloides	New Zealand	Colenso 131
oblectabilis	Pelloporus	Brazil	Ule 164
oblectans*	Pelloporus	Australia	Berkeley 164
obliquus	Petaloides	New Guinea	Masse 132
oblivionis	Pelloporus	Brazil	Spruce 164
obovatus	Petaloides	Java	Junghuhn 141
ocellatus	Amaurodermus	Brazil	Berkeley 119
ochrolaccatus	Ganodermus	Philippines	Montagne 105
omphalodes	Amaurodermus	Brazil	Berkeley 1
opacus	Ganodermus	Brazil	Montagne 106
orbicularis	Lentus	Europe	Sauter 176
orientalis	Pelloporus	Japan	Umemuro 193
osseus	Petaloides	Europe	Kalchbrenner 191
ovinus	Ovinus	Europe	Schaeffer 167
pachypus*	Spongiosus	Cuba	Montagne 159
palpebralis	Melanopus	South America	Leprieur 184
pansus	Lignosus	South America	Berkeley 125
partitus	Lentus	Brazil	Berkeley 175
Paucheri*	Melanopus	New Caledonia	Patouillard 180
Paulensis*	Lignosus	Brazil	Hennings 126
Peckianus	Lentus	United States	Cooke 171
penetralis	Petaloides	Exotic	Smith 132
perennis	Pelloporus	Europe	Linnaeus 164
perversus*	Petaloides	Philippines	Copeland 136
Pes caprae	Ovinus	Europe	Persoon 167
petaliformis	Petaloides	Cuba	Berkeley 142
petalodes	Petaloides	Brazil	Berkeley 133
picipes*	Melanopus	Europe	Fries 180
pictus	Pelloporus	Europe	Schulter 164
pisachapani*	Ganodermus	Java	Nees 107
placopus	Ganodermus	Java	Léveillé 105
platotis*	Petaloides	Australia	Berkeley 137
Pocula	Petaloides	United States	Schweinitz 140
podlachicus*	Melanopus	Europe	Bresadola 183
politus*	Ovinus	Europe	Fries 167
polydactylus*	Lignosus	Brazil	Berkeley 126
popanoides	Ovinus	Mauritius	Cooke 167
porphyritis	Petaloides (Micro.)	Brazil	Berkeley 142
praetervisus*	Amaurodermus	Brazil	Patouillard 113
Preussii	Lignosus	Africa	Hennings 124
pseudoperennis	Lentus	Africa	Holst 174
pterygodes*	Petaloides (Micro.)	Africa	Fries 143
pudens*	Lignosus	India	Berkeley 126
Puiggarianus	Spongiosus	Brazil	Hennings 160
pusillus (bis)	Melanopus	South America	Fries 185
pusiolus	Petaloides	East Indies	Cesati 140
Puttemansii	Melanopus	Brazil	Hennings 183
radiato-scruposus	Melanopus	Brazil	Hennings 187
radicatus	Ovinus	United States	Schweinitz 168
Ramosii*	Amaurodermus	Philippines	Murrill 111

INDEX VA.

regulicolor*	Ganodermus.		
renatus	Amaurodermus.	B.	
renidens	Amaurodermus.	Braz.	Bresadola
repando-lobatus*	Lentus.	Brazil.	Spagazzini
Repsoldi*	Spongiosus.	Brazil.	Moeller
retiporus*	Merismus.	Australia.	Cooke
rhinocerotis	Lignosus.	Malay.	Cooke
Rhipidium	Petaloides.	United States.	Berkeley
rhizomatophorus	Lignosus.	Brazil.	Hennings
rhizomorphus	Melanopus.	South America.	Montagne
rhizophilus.	Lentus.	Africa.	Patouillard
Ridleyi.	Merismus.	Tasmania.	Massee
rivulosus.	Amaurodermus.	Java.	Patouillard
Rostkovii*	Ovinus (Melan.)	Europe.	Fries
rubidus	Petaloides.	Ceylon.	Berkeley
rubro-castaneus.	Melanopus.	Perak.	Ridley
rudis	Amaurodermus.	Tasmania.	Berkeley
rufescens.	Spongiosus.	Europe.	Persoon
gosus	Amaurodermus.	Java.	Nees
ussiceps	Petaloides.	Ceylon.	Berkeley
rutrosus*	Petaloides.	Europe.	Rostkovius
Sacer	Lignosus.	Africa.	Fries
sanguineus.	Petaloides.	South America.	Linnaeus
Sapurema	Ovinus.	Brazil.	Moeller
scabriceps	Lentus.	Cuba.	Berkeley
Schomburgkii	Amaurodermus.	South America.	Berkeley
Schweinitzii	Spongiosus.	Europe.	Fries
scopulosus	Lignosus.	Australia.	Berkeley
sericatus.	Amaurodermus.	Africa.	Holland
sideroides.	Spongiosus.	Java.	Léveillé
siennaeolor*	Petaloides.	Ceylon.	Cooke
sordulentus*	Merismus.	Chili.	Montagne
Sprucei.	Amaurodermus.	Brazil.	Patouillard
			(orig. Berkeley)
squamatus	Ovinus.	Europe.	Kalchbrenner
squamosus	Ovinus (Melan.)	Europe.	Fries
stereinus.	Petaloides.	Cuba.	Berkeley
subbulbipes*	Spongiosus.	South America.	Hennings
sui fulvus	Petaloides.	Cuba.	Berkeley
subvirgatus.	Lentus.	India.	Cave
sulphureus	Merismus.	Europe.	Fries
superpositus	Lignosus.	Australia.	Berkeley
talpae.	Merismus.	Brazil.	Cooke
Tasmanicus.	Ovinus.	Tasmania.	Massee
Tiliae*	Lentus.	Europe.	Schulzer
tomentosus.	Spongiosus.	Europe.	Fries
tricholoma	Lentus.	Cuba.	Montagne
triqueter*	Spongiosus.	Europe.	Fries
			(Name orig. Persoon)
tristiculus.	Petaloides.	South America.	Montagne
Tuba.	Lentus.	Cuba.	Berkeley
tuberas ter.	Ovinus.	Europe.	Fries
tumulosus	Ovinus.	Australia.	Cooke
turboformis.	Pelloporus.	India.	Krumbiegel
umbellatus.	Merismus.	Europe.	Viviani
umbilicatus	Lentus.	India.	Berkeley
unilaterus.	Amaurodermus.	Brazil.	Spruce
(unnamed).	Ganodermus.	Africa (?).	
vadosus.	Melanopus.	West Indies.	Duss
valesiacus*	Ganodermus.	Europe.	Boudier

VALID SPECIES.

			PAGE
atus	Fenopodis	India	Berkeley 162
variabilis	Amaurodermus	Brazil	Berkeley 111
varius	Melanopus	Europe	Persoon 180
veluticeps	Melanopus	Africa	Cooke 182
vernalis	Lentus	Europe	Fries 171
vernicosus	Melanopus	Brazil	Berkeley 182
vernicipes	Petaloides	Japan	Berkeley 144
virgatus	Lentus	Cuba	Berkeley 172
Warburgianus	Melanopus	East Indies	Hennings 186
Wrightii	Melanopus	Cuba	Murrill 183
Wynnei	Merismus	Europe	Berkeley 150
Xanthopus	Lentus (Micro.)	Africa	Fries 173
Xerophyllus	Melanopus	New Zealand	Berkeley 186
Zambesianus	Lignosus	Africa	Buchanan 128
Zelandicus	Merismus	New Zealand	Cooke 149

UNIVERSITY OF MICHIGAN



3 9015 05978 1719

BOUND

FEB 21 1955

UNIV. OF MICH.
LIBRARY

